

ICOMAA-2023

**6th INTERNATIONAL HYBRID CONFERENCE
ON MATHEMATICAL ADVANCES
AND APPLICATIONS**

ABSTRACT BOOK

Editors
Yusuf ZEREN
Murat KIRIŞÇI
Adem Cengiz ÇEVİKEL



ISBN


10-13 MAY, ISTANBUL



website

6th International HYBRID Conference on Mathematical Advances and Applications

May 10-13, 2023, İstanbul / TÜRKİYE



ISBN:978-605-69387-4-0
Yıldız Technical University, İstanbul, TURKEY –2023



FOREWORDS

Dear Conference Participant,

Welcome to the International Hybrid Conference on Mathematical Development and Applications (ICOMAA-23) we organized the sixth. First of all, I would like to start my words by reminding one of G. H. Hardy's words:

"Mathematics, more than any other art or science, is a young man's game."

This phrase he expressed in his book "A Mathematician's Apology" is quite meaningful. Because Newton discovered his biggest ideas, fluxions and the law of gravitation, when he was just 24 years old. He found the 'elliptic orbit' at 37 years old. Also, Galois(at twenty-one), Abel(twenty-seven), Ramanujan(thirty-three), and Riemann(at forty) had passed away in their youth.

That's why we thought we should continue this series of conferences that brings together exciting and productive young mathematicians. So, we aim to bring together scientists and young researchers from all over the world and their work on the fields of mathematics and applications of mathematics, to exchange ideas, to collaborate and to add new ideas to mathematics in a discussion environment. With this interaction, functional analysis, approach theory, differential equations and partial differential equations and the results of applications in the field of Mathematics are discussed with our valuable academics, and in mathematical developments both science and young researchers are opened. We are happy to host many prominent experts from different countries who will present the state-of-the-art in real analysis, complex analysis, harmonic and non-harmonic analysis, operator theory and spectral analysis, applied analysis.

I would like to express my gratitude to those who see and appreciate our efforts and innovative steps that we have made to improve our conference every year, to our dear invited speakers and to all our participants. I owe a debt of gratitude to the Scientific committee, organizing committee, local organizing committee and for their efforts throughout this conference series.

The conference brings together about 203 participants and 9 invited speakers from 22 countries (Azerbaijan, India, Algeria, Bangladesh, Georgia, Greece, Iran, Iraq, Italy, Kazakhstan, Kosovo, Malaysia, Mexico, Morocco, Pakistan, Poland, Saudi Arabia, Turkey, United Arab Emirates, Uzbekistan, Yemen).

More than 50% of our participants participated from abroad. This shows that the conference meets the criteria of being international.

It is also an aim of the conference to encourage opportunities for collaboration and networking between senior academics and graduate students to advance their new perspective. Additional emphasis on ICOMAA-23 applies to other areas of science, such as natural sciences, economics, computer science, and various engineering sciences, as well as applications in related fields. The articles submitted to this conference will be addressed on the conference web sites and, in the journals, listed below:

- Miskolc Mathematical Notes,
- Türkiye Mathematical Sciences
- Sigma Journal of Engineering and Natural Sciences,
- Istanbul Commerce University Journal of Sciences,
- Journal of Nonlinear Sciences and Applications,
- *Special Issue "Symmetries of Difference Equations, Special Functions and Orthogonal Polynomials" in Symmetry,

This booklet contains the titles and abstracts of almost all invited and contributed talks at the **6th International E-Conference on Mathematical Advances and Applications**. Only some abstracts were not available at the time of printing the booklet. They will be made available on the conference website <http://2023.icomaas.com/> when the organizers receive them.

We wish everyone a fruitful conference and pleasant memories throughout the online conference.

Prof. Dr. Yusuf ZEREN
On Behalf of Organizing Committee
Chairman

It was a big excitement moment when Prof. Dr. Yusuf ZEREN discussed with me on the issue of "6th International Hybrid Conference on Mathematical Development and Applications" (ICOMAA-2023) in Yıldız Technical University, Istanbul. It is a great pleasure that this conference is going to take place now. As one of the organizers of the conference, I am delighted with all the delegates, distinguished mathematicians, speakers and young researchers in this international event. It is expected that delegates and participants will benefit from this conference experience and the legacy of information dissemination will continue.

I wish all of you to have a nice and enjoyable participation in the conference.

Prof. Dr. Necip ŞİMŞEK

SCIENTIFIC COMMITTEE

Abdel-Aty, M., Egypt
Abdeljawad, T., Arabia
Abdullayev, F., Türkiye
Agarwal, P., India
Akin, L., Türkiye
Aktosun, T., USA
Altomare, F., Italy
Bilalov, B., Azerbaijan
Bouhamidi, A., France
Braha, N., Kosova
Burenkov, V. I., U.K.
Celik, E., Türkiye
Cevikel, A.C., Türkiye
Chalabi, A., France
Cruz-Uribe, D., U.S.A.
Diening, L., Germany
Duman, O., Türkiye
Ersoy, B.A., Türkiye
Fiorenza, A., Italy
Gasymov, T., Azerbaijan
Gogatishvili, A., Czech
Gulcu, A., Türkiye
Guliyev, V.S., Azerbaijan
Guzel, N., Türkiye
Hästö, P.A., Finland
Hristov, J., Bulgaria
Huseynli, A., Azerbaijan
Isgenderoglu, M., Türkiye
Ismayilov, M., Azerbaijan
Izgi, A., Türkiye
Jbilou, K., France
Kalantarov, V., Türkiye
Kara, E.E., Türkiye
Karakaya, V., Türkiye
Klingler, B., France
Kokilashvili, V., Georgia
Kolatoglu, E., Türkiye
Lang, J., U.S.A.
Loeser, F., France

Mamedov, F., Azerbaijan
Mardanow, M.J., Azerbaijan
Mendez, O., USA
Monsurro, S., Italy
Mursaleen, M., India
Nieto, J. J., Spain
Nuray, F., Türkiye
Oleg, R., Russia
Ozger, F., Türkiye
Pascu, M., Romania
Phong, D. H., USA
Pinto, C., Portugal
Piskin, E., Türkiye
Rakotoson, J.M., France
Sajid, M., Saudi Arabia
Samko, S., Portugal
Sari, M., Türkiye
Savas, E., Türkiye
Secer, A., Türkiye
Serbetci, A., Türkiye
Sevli, H., Türkiye
Rezapour, S., Iranian
Sawano, Y., Japan
Shukurov, A., Azerbaijan
Simsek, N., Türkiye
Srivastava, H.M., Canada
Softova-Palagacheva, L., Italy
Soleimani-damaneh, M., Iranian
Sun-Sig, B., South Korea
Stan, A., USA
Tabachnikov, S., USA
Toprakseven, S., Türkiye
Ugur, T., Türkiye
Usta, F., Türkiye
Wang, W., China
Zeren, Y., Türkiye


ORGANIZING COMMITTEE

Yusuf ZEREN (Chairman)(Turkey)
Misir J. MARDANOV (Azerbaijan)
Necip SIMSEK(Turkey)
Lyoubomira SOFTOVA PALAGACHEVA (Italy)
Amiran GOGATISHVILI (Czech Republic)
Claudia CAPONE (Italy)
Naim L. BRAHA (Kosovo)
Sofiene TAHAR (Canada)
Fatih SIRIN(Turkey)
Seyma CETIN (Turkey)
Cemil KARAÇAM (Turkey)
Feyza Elif DAL (Turkey)

LOCAL ORGANIZING COMMITTEE

Kader Simsir Acar, Istanbul Commerce University, Türkiye
Selim Yavuz, Indiana University, USA
Melih Cinar, Yildiz Technical University, Türkiye
Elif Deniz, Concordia University, Canada
Kubra Aksoy, Concordia University, Canada
Faruk Özger, İzmir Katip Celebi University, Türkiye
Selmahan Selim, Yildiz Technical University, Türkiye
Yunus Atalan, Aksaray University, Türkiye
Canay Aykol, Ankara University, Türkiye
Erdem Kocakuşaklı, Sinop University, Türkiye
Mustafa Gezek, Namik Kemal University, Türkiye
Arash Ghorbanalizadeh, IASBS, Iran
Ersin Şener, Kırklareli Üniversitesi, Türkiye
Harun Baldemir, Cankiri Karatekin University, Türkiye
Burhan Tiryakioğlu, Marmara Üniversitesi, Türkiye
Serkan Onar, Yildiz Technical University, Türkiye
Tuğçe Ünver Yıldız, Kirikkale University, Türkiye
Gülşen Ulucak, Gebze Technical University, Türkiye
Hülya Öztürk, Gebze Technical University, Türkiye
Ibrahim Murat Turhan, Yildiz Technical University, Türkiye
Mohammed Alsharafi, Yıldız Technical University, Türkiye
Abuzer Gunduz, Sakarya University, Türkiye
Seher Kaya, Ankara University, Türkiye
Abdulhamit Küçükaskan, Yıldırım Beyazıt University, Türkiye
Lubos Pick, Charles University, Czech Republic
Sabina Sadigova, Azerbaijan National Academy of Sciences, Azerbaijan
Valdete Loku, University of Applied Sciences, Ferizaj, Kosovo
Salvatore Tramontano, University of Salerno, Italy
Rosamaria Rescigno, University of Salerno, Italy
Emilia Anna Alfano, University of Salerno, Italy
Irshaad Ahmed, Sukkur IBA University, Pakistan
Julio S. Neves, University of Coimbra, Portugal
Pankaj Jain, South Asian University, India
Abdu Alameri, University of Science and Technology, Yemen
Mohammed Mohsen, Sheba Region University, Yemen
Haifa Ahmed, Aden University, Yemen
Aydin Shukurov, Azerbaijan National Academy of Sciences, Azerbaijan
Rakif Efendiyev, Baku State University, Azerbaijan

Tengiz Kopaliani, Tbilisi State University, Georgia
Zahira Mamedova, Azerbaijan National Academy of
Sciences, Azerbaijan



ICOMAA 2023

6th International HYBRID Conference on Mathematical Advances and Applications May 10-13, 2023,
İstanbul / TÜRKİYE
<https://2023.icomaas.com/>

CONTENTS

FOREWORDS	2
SCIENTIFIC COMMITTEE	4
ORGANIZING COMMITTEE	5
LOCAL ORGANIZING COMMITTEE	5
INVITED TALKS	22
HOLMSTEDT'S FORMULA FOR THE K-FUNCTIONAL: THE LIMIT CASE $\theta_0 = \theta_1$	22
AMIRAN GOGATISHVILI.....	22
SOME REGULARITY RESULTS FOR A CLASS OF ELLIPTIC EQUATIONS WITH LOWER ORDER TERMS AND ORLICZ GROWTH	23
CLAUDIA CAPONE.....	23
MATRIX WEIGHTS IN VARIABLE LEBESGUE SPACES: THE CASE OF AVERAGING OPERATORS	24
DAVID CRUZ-URIBE, OFS.....	24
GENERALISED FRACTIONAL OPERATORS WITH APPLICATIONS IN BIOLOGY	25
DUMITRU BALEANU ^{1,2}	25
MAXIMAL REGULARITY FOR NON-AUTONOMOUS PROBLEMS WITH A PRIORI ASSUMPTIONS	26
PETER HÄSTÖ.....	26
SOLVABILITY IN THE SMALL OF HIGHER ORDER ELLIPTIC EQUATIONS IN WEIGHTED SOBOLEV-BANACH SPACES	27
B. BILALOV, S. SADIGOVA, L.SOFTOVA ¹ , Y. ZEREN.....	27
SOME NON-MATRIX SUMMABILITY METHODS AND KOROVKIN TYPE APPROXIMATION THEOREMS	28
M. MURSALEEN.....	28
CONES GENERATED BY A GENERALIZED FRACTIONAL MAXIMAL FUNCTION	29
NURZHAN BOKAYEV ¹ , AMIRAN GOGATISHVILI ² , AZHAR ABEK ³	29
WEIGHTED MORREY SPACES	30
YOSHIHIRO SAWANO.....	30
CONTRIBUTED TALKS	31
ON THE SOME RELATIONS SHIP BETWEEN POSITIVE AND POSITIVE P-SUMMING OPERATORS	31
ABDELMOUMEN TIAIBA ¹	31
AN ALGORITHM FOR RELAXING MULTICURVES ON SMALL GENUS NON-ORIENTABLE SURFACES	32
FERIHE ATALAN ¹ , ABDULLAH BAYKAL ² , ÖYKÜ YURTTAŞ ³	32
SOME PROPERTIES OF GRILL TOPOLOGICAL SPACES VIA $G^A_\Omega O(X)$	33
AMIN SAIF ¹ , A. MAHDI ²	33

BOUNDED FINITE-TIME STABILIZING POSITIONAL CONTROLS VIA THE CONTROLLABILITY FUNCTION AS MOTION TIME	34
ABDON E. CHOQUE-RIVERO ¹ ,	34
TWO-WEIGHTED INEQUALITIES FOR SOME CLASSICAL OPERATORS AND THEIR COMMUTATORS IN THE MODIFIED MORREY SPACES	35
ABDULHAMIT KUCUKASLAN	35
FEKETE-SZEGÖ PROBLEM FOR CERTAIN SUBCLASSES OF ANALYTIC FUNCTIONS ASSOCIATED WITH THE COMBINATION OF DIFFERENTIAL OPERATORS	36
ADEM KIZILTEPE ¹ , ERHAN DENİZ ² , SERCAN KAZIMOĞLU ³	36
INVERSE SCATTERING PROBLEM FOR THE SCHRÖDINGER EQUATION WITH AN ADDITIONAL GROWING POTENTIAL ON THE ENTIRE AXIS	37
DAVUD ORUCOV ¹ AND AGIL KHANMAMEDOV ^{1,2}	37
DOUBLY DIFFRACTED FIELDS STIMULATED BY NON-SYMMETRICAL EDGES AT THE JUNCTION	38
ALAA SERHAN ¹ , FERAY HACIVELIOĞLU ²	38
OPTIMAL LINEAR APPROXIMATION AND ISOMETRIC EXTENSIONS	39
ALEXANDER KUSHPEL	39
A NEW METHOD ABOUT FINDING SQUARE ROOT OF PERFECT SQUARE NUMBERS	40
ALI DEĞİRMENCI	40
ON COVERED QUASI HYPERIDEALS IN SEMIHYPERGROUPS	41
SHAHNAWAZ	41
EXPONENTIAL GRAPH RESOLVENT	42
ALI A. SHUKUR	42
ON BASICITY A SYSTEM OF EIGENFUNCTIONS OF A SECOND ORDER DISCONTINUOUS DIFFERENTIAL OPERATOR IN LEBESGUE SPACES	43
ALIRZA AKHMEDOV ¹ AND İLKIN FEYZULLAYEV ²	43
EXISTENCE OF SOLUTIONS FOR IMPULSIVE FRACTIONAL BOUNDARY VALUE PROBLEM WITH P-LAPLACIAN VIA CRITICAL POINT THEORY	44
AMINA BOUCENNA ¹ , AMIRA ROUAGHI ²	44
EXISTENCE AND UNIQUENESS OF SOLUTIONS FOR P-LAPLACIAN FRACTIONAL BOUNDARY VALUE PROBLEM ON THE HALF-LINE	45
AMIRA ROUAGHI ¹ , AMINA BOUCENNA ²	45
HARDY-LITTLEWOOD MAXIMAL FUNCTION ON LOCALLY COMPACT ABELIAN GROUPS	46
ANI OZBETELASHVILI ¹	46
NOVEL TEMPERED FRACTIONAL HERMITE-HADAMARD TYPE INEQUALITIES VIA n-POLYNOMIAL UNIFORMLY CONVEX FUNCTIONS WITH APPLICATION	47
ARTION KASHURI ¹ , ROZANA LIKO ²	47
APPROACHING GEOMETRIC PROBLEMS WITH THE PARTICLE SWARM OPTIMIZATION METHOD	48
KARAÇAM CEMİL ¹ , SEVGİ ATA ² , KARACAN EFE ATAKAN ³	48

KARTAL ARDA EREN ⁴	48
ON FRAME PROPERTIES OF ITERATES OF A MULTIPLICATION OPERATOR.....	49
A.SH.SHUKUROV	49
LEVINSON-TYPE FORMULA FOR STURM-LIOUVILLE OPERATOR WITH A RATIONAL FUNCTION OF SPECTRAL PARAMETER IN BOUNDARY CONDITION	50
AYNUR ÇÖL ¹	50
ON A SCATTERING PROBLEM FOR DISCONTINUOUS SECOND ORDER DIFFERENTIAL OPERATORS WITH HERGLOTZ FUNCTION OF SPECTRAL PARAMETER IN BOUNDARY CONDITION	51
AYNUR ÇÖL ¹ AND KHANLAR R. MAMEDOV ²	51
BASICITY OF THE DOUBLE EXPONENTIAL SYSTEM IN WEIGHTED LEBESGUE SPACE.....	52
AYSEL GULIYEVA	52
CONFORMAL MAPPING PRESERVING THE Z-SYMMETRIC TENSOR ON A RIEMANNIAN MANIFOLD	53
AYŞE YAVUZ TAŞCI ¹ ,FÜSUN ÖZEN ZENGİN ²	53
EXISTENCE AND DECAY OF SOLUTIONS FOR A FOURTH-ORDER HYPERBOLIC EQUATION WITH A VARIABLE COEFFICIENTS	54
AYŞE FIDAN ¹ ,ERHAN PIŞKIN ²	54
THE INVERSE STRUM-LIOUVILLE PROBLEM WITH ONE OF THE BOUNDARY CONDITIONS INCLUDING EIGENVALUE	55
SEDA KIZILBUDAK ÇALIŞKAN, AYŞEGÜL KILIÇ.....	55
FINITE ELEMENT BASED SCHEME FOR THE BURGERS EQUATION WITH FORCING EFFECTS.....	56
SELMAHAN SELİM ¹ ,AYSENUR BUSRA ÇAKAY ²	56
ON NON-INCREASING REARRANGEMENT OF GENERALIZED FRACTIONAL MAXIMAL FUNCTION	57
AZHAR ABEK	57
ON THE STRONG SOLVABILITY OF A NONLOCAL BOUNDARY VALUE PROBLEM FOR THE LAPLACE EQUATION IN A RECTANGULAR DOMAIN.....	58
TELMAN GASYMOV ¹ AND BEHERCHIN AKHMEDLI ²	58
STUDY OF NONLINEAR PARTIAL FRACTIONAL BOUNDARY VALUE PROBLEM.....	59
NOUREDDINEBOUTERAA ¹	59
RECURSIVE ESTIMATION OF THE CONDITIONAL DISTRIBUTION FUNCTION ON RIEMANNIAN MANIFOLDS	60
BOUADJEMI ABDELKADER ¹	60
ON THE RECURSIVE SEQUENCE	61
BURAKOĞUL ¹	61
DYNAMICS AND HOPF BIFURCATION OF A SUSTAINABLE TOURISM MODEL WITH TIME DELAY	62
PROF. DR. MUSTAFA TAYLAN ŞENGÜL ¹ AND BÜNYAMIN KURTKAYA ²	62
KOROVKIN-TYPETHEOREMSANDTHEIRSTATISTICALVERSIONSINTHEWEIGHTED GRAND-LEBESGUESPACES WITHOUT MUCKENHOPT CONDITION	63
YUSUF ZEREN ¹ , MIGDAD ISMAILOV ² , CEMIL KARACAM ³ AND FATİH SIRIN ⁴	63
HARMONIC TRINOMIAL WITH COMPLEX	64
PARAMETER	64

CHAHRAZED HARRAT ¹	64
NUMERICAL SOLUTION OF SECOND-ORDER PARTIAL VOLTEPRA INTEGRO DIFFERENTIAL EQUATIONS VIA TAYLOR COLLOCATION METHOD	65
CHEIMA KHENNAOUI ¹	65
EXISTENCE OF REAL MODELS OF K₃-SURFACES IN ARITHMETICAL TERMS	66
ÇISEM GÜNEŞ AKTAŞ	66
MAXIMAL-SIMULTANEOUS APPROXIMATION BY FABER SERIES IN BERGMAN SPACE	67
DANIYAL M. ISRAFILOV	67
THE APPLICATION OF EXPANSIVE MAPPINGS TO THE EXISTENCE PROBLEMS	68
DARIUSZ WARDOWSKI	68
BOUNDEDNESS AND COMPACTNESS OF COMMUTATORS FOR RIESZ POTENTIAL ON LOCAL MORREY-TYPE SPACES. ..	69
DAUREN MATIN ¹ , TALGAT AKHAZHANOV ¹ , AIDOS ADILKHANOV ¹	69
PRE & POST PANDEMIC FORMATIVE ASSESSMENTS METHODOLOGIES TO IMPROVE MATHEMATICS LEARNING FOR UNIVERSITY STUDENTS	70
DIANA AUDI ¹	70
PERIODICITY AND STABILITY IN THE KALDOR MODEL OF ECONOMICS WITH IMPULSE EFFECTS	71
DIĞDEM ŞEN ¹ , FATMA TOKMAK FEN ² ,	71
ADAPTATION OF SOME GRACEFUL GRAPH LABELING METHODS TO COMBINATORIAL BLOCK DESIGNS	72
DILARA ERDEMİR ¹ AND EMRE KOLOTOĞLU ²	72
EMBEDDING THE STATIONARY SPACETIMES INTO BRANS-DICKE COSMOLOGY VIA CONFORMAL TRANSFORMATIONS	73
DILEK KAZICI ¹ , ALI DEMİRÖZLÜ ²	73
ON A NEW TYPE GAUSSIAN SEQUENCES	74
ECE GULSAH COLAK ¹ , NAZMIYE GONUL BILGIN ² , YUKSEL SOYKAN ³	74
ON THE FACTORIAL OF SPLIT QUATERNIONS	75
EDANUR ERGÜL ¹ , SALIM YÜCE ²	75
SPECTRAL ANALYSIS OF A SINGULAR DISSIPATIVE THIRD-ORDER DIFFERENTIAL OPERATOR	76
EKİN UĞURLU ¹	76
DEPENDENCE OF EIGENVALUES OF DIRAC SYSTEMS ON THE DATA	77
EKİN UĞURLU ¹	77
SOME NEW RESULTS OF WEIGHTED GENERALIZED NORLUND-EULER STATISTICAL CONVERGENCE	78
EKREM ALJIMI ¹ , PENPARK SIRIMARK ² , AGON MAHMUTİ ³	78
ZERO-HOPF BIFURCATION OF FOURTH-ORDER QUADRATIC POLYNOMIAL DIFFERENTIAL SYSTEMS, VIA AVERAGING THEORY OF THIRD ORDER	79
BENDİB EL OUAHMA ¹ , MAKHLOUF AMAR ²	79
DISSIPATIVE EXTENSIONS OF ONE OPERATOR WITH POTENTIAL IN THE FORM OF A DIRAC FUNCTION	80
ELDAR M. MUSTAFAYEV	80

DRBEM SOLUTION OF MHD FLOW IN A RECTANGULAR DUCT UNDER AXIALLY-CHANGING EXTERNAL MAGNETIC FIELD	81
ELIF EBREN KAYA ¹ , MÜNEVVER TEZER-SEZGIN ²	81
ON K- ORESME HYBRID POLYNOMIALS	82
SERPİL HALICI ¹ AND ELIFCAN SAYIN ²	82
AN INVERSE COEFFICIENT PROBLEM FOR A TWO-DIMENSIONAL HYPERBOLIC EQUATION WITH INTEGRAL OVERDETERMINATION	83
ELVIN AZİZBAYOV ¹ AND GÜLNARA SHUKUROVA ²	83
NUMERICAL INVESTIGATION OF STIFF EPIDEMIOLOGICAL MODELS	84
EMEL KURUL ¹ , MURAT SARI ² , HÜSEYİN TUNÇ ³	84
SS-INJECTIVE MODULES	85
EMİNE ÖNAL KIR ¹	85
POWER SERIES STATISTICAL CONVERGENCE OF CHENEY-SHARMA OPERATORS	86
EMRE GÜVEN ¹ AND DİLEK SÖYLEMEZ ² ,	86
HIGH ORDER NUMERICAL APPROACH FOR SOLVING THE MODIFIED EQUAL WIDTH EQUATION	87
EMRE KIRLI	87
EFFECT OF CAPUTO-FABRIZIO DERIVATIVE ON SCABIES MODEL	88
ESMEHAN UÇAR ¹	88
ALTERNATIVE SOLUTION METHOD ON MATRIX GAMES WITH FUZZY GOALS	89
ADEM CENGİZ CEVİKEL ¹ , ESRA SOSLU ²	89
A NOTE ON SINGULAR INTEGRAL OPERATORS ASSOCIATED WITH LAPLACE-BESSEL DIFFERENTIAL OPERATOR IN VARIABLE LORENTZ SPACE	90
ESRA KAYA ¹ , CANAY AYKOL ²	90
A LOCAL TAYLOR SERIES METHOD FOR NON-LINEAR SINGULAR BOUNDARY VALUE PROBLEMS	90
ESRA ÇELİK ¹ , HÜSEYİN TUNÇ ² , MURAT SARI ³	90
MECHANISTIC MODELLING OF CONTACT TRACING AND ISOLATION POLICIES	91
ESRA ÖZGE ASAN ¹ , MURAT SARI ² , ENES SEYFULLAH KOTİL ³ , HÜSEYİN TUNÇ ⁴	91
THERMOELASTIC LAMINATED TIMOSHENKO BEAM WITH DISTRIBUTED DELAY: GLOBAL EXISTENCE AND STABILITY	92
FARES YAZID ¹ , FATİMA SİHAM DJARADI ² AND NOR-EL-HOUDA BEGHERSA ³	92
ON SOME BETA TYPE OPERATORS	93
FARUK ÖZGER	93
ON SOME BETA TYPE OPERATORS	94
FARUK ÖZGER	94
LIE GROUP PROPERTY OF GENERALIZED LINEAR DEFORMATION TRANSFORMATION	95
FATİH TUĞRUL ¹ , ŞENAY BAYDAŞ ² AND BÜLENT KARAKAŞ ³	95
A NOTE ABOUT PSEUDOSPECTRUM OF THE TENSOR PRODUCT OPERATOR	96
FATİH YILMAZ ¹	96

ON BASICITY OF THE SYSTEM OF EIGENFUNCTIONS FOR A DISCONTINUOUS SPECTRAL PROBLEM IN VARIABLE EXPONENT LEBESGUE SPACE	97
FATİH SIRIN ¹	97
CONSTRUCTING LYAPUNOV FUNCTIONS FOR DELAY REACTION DIFFUSION SYSTEMS AND APPLICATION	98
FATIHA NAJM ¹ , RADOUANE YAFIA ¹ , M. A. AZİZ ALAOUİ ²	98
SOLUTION OF COMPOSITE BEAM WITH PERIODICAL CURVATURE BY FINITE DIFFERENCE METHOD UNDER UNIFORM LOADINGS.....	99
FATMA SAYDAM AKMAN ¹ , ZAFER KÜTÜĞ ² , ONUR ÇINAR ³ , ÖZLEM VARDAR ⁴ AND YOUSAF ANWAR ⁵	99
MACHINE LEARNING MODELS IN THE DIAGNOSIS OF ALZHEIMER DISEASE	100
FATMA NUR BUDAK ¹ , MURAT SARI ² , HUSEYİN TUNC ³	100
A NEW FUNCTION TYPE IN GRILL MINIMAL SPACES AND ITS PROPERTIES	101
FERİT YALAZ.....	101
APPROXIMATIVE PROPERTIES OF PERTURBED EXPONENTIAL SYSTEM IN ORLICZ SPACES	102
BİLAL BİLALOV ^{1,2} , YUSUF ZEREN ¹ , FİDAN ALİZADEH ² , FEYZA ELİF DAL ¹	102
ALMOST SHORTEST PATHS: TO ALTERNATE ROUTES FOR SECURITY REASONS IN GRAPHS	103
GÖKÇE ÇAYLAK KAYATURAN.....	103
UPPER BOUNDS FOR BLOW UP OF SOLUTIONS FOR A FOURTH-ORDER PARABOLIC EQUATION WITH VISCOELASTIC TERM	104
GÜLISTAN BUTAKIN ¹ , ERHAN PIŞKIN ²	104
BLOW UP SOLUTIONS FOR A FOURTH-ORDER PARABOLIC EQUATION WITH VARIABLE EXPONENTS.....	105
GÜLISTAN BUTAKIN ¹ , ERHAN PIŞKIN ²	105
INVESTIGATION OF HADAMARD FRACTIONAL DIFFERENTIAL INCLUSIONS	106
HABİDJOURDEM ¹	106
NEUTROSOPHIC STATISTICAL ANALYSIS OF TEMPERATURES OF CITIES IN THE SOUTHEASTERN ANATOLIA REGION OF TURKEY	107
HACER ŞENGÜL KANDEMİR ¹ , NAZLIM DENİZ ARAL ² , MURAT KARAKAŞ ³ , MİKAIL ET ⁴	107
SEMI GLOBAL DOMINATION NUMBER IN PRODUCT FUZZY GRAPHS	108
HAİFA AHMED ¹ , MAHİOUB SHUBATAH ² , MOHAMMED ALSHARAFI ³	108
TRACE FORMULA FOR DIFFERENTIAL EQUATION WITH COEFFICIENT CONTAINING FIRST DEGREE POLYNOMIAL DEPENDING ON SPECTRAL PARAMETER	109
HAJAR MOVSUMOVA.....	109
SOME IMPROVEMENT OF BEREZIN RADIUS INEQUALITIES VIA SPECHT'S RATIO.....	110
HAMDULLAH BAŞARAN ¹ , MEHMET GÜRDAL ²	110
EXPONENTIAL, LOGARITHMIC AND TRIGONOMETRIC FUNCTIONS ON SPLIT QUATERNIONS AND THEIR CHARACTERIZATIONS	111
HAMİDE FEYZA AYKUT ¹ , SALIM YÜCE ²	111
SOME COMBINATORIAL RESULTS FOR THE HYPEROCTAHEDRAL GROUP	112
HASAN ARSLAN	112

A RELATION BETWEEN BOSBACH STATE OPERATOR AND VERY TRUE OPERATOR	113
IBRAHIM SENTURK.....	113
SOME EXISTENCE THEOREMS FOR SOLUTIONS OF SET-VALUED OPTIMIZATION PROBLEMS WITH RESPECT TO VARIOUS SET ORDER RELATIONS	114
İLKNUR ATASEVER GÜVENÇ ¹	114
REVENUE MAXIMIZATION FOR POLYNOMIAL DEMAND FUNCTION USING INTERVAL-VALUED TRAPEZOIDAL FUZZY NUMBERS	115
ISMAIL TEMBELO ¹ , SALIH AYTAR ²	115
MINKOWSKI DIFFERENCE AND SUM OF HALF PLANES	116
JALOLXON NURITDINOV.....	116
ON THE COMPLETENESS AND MINIMALITY IN THE SPACE $Lp(0, \pi)$ OF A WEIGHTED SYSTEM OF SINES WITH A POWER-LAW WEIGHT WITH AN EXCLUDED ARBITRARY ELEMENT.....	117
YUSUF ZEREN ¹ , MİGDAD I. ISMAILOV ² , KADER ŞİMŞİR ACAR ³	117
A NOVEL ALTERNATIVE ALGORITHM TO FIND ALL MULTIPLE SOLUTIONS OF GENERAL INTEGER LINEAR PROGRAMS	118
KADRIYE ŞİMŞEK ALAN.....	118
A NOVEL ALTERNATIVE ALGORITHM FOR SOLVING INTEGER LINEAR PROGRAMMING PROBLEMS	119
KADRIYE ŞİMŞEK ALAN.....	119
EFFECTS OF MUSICS COMPOSED USING MATHEMATICAL METHODS AND DNA ON THE EEG FREQUENCY BANDS OF HEALTHY INDIVIDUALS	120
CEMİL KARAÇAM ¹ , HALİL YAKIT ² , KAYRA EGE ALTUN ³ , ŞERİF EFE DARTAR ⁴	120
NEW TRIANGULAR Q-FIBONACCI MATRIX AND ITS APPLICATIONS	121
KORAY İBRAHİM ATABEY ¹ , MUHAMMED ÇINAR ² , MİKAIL ET ³	121
A NEW APPROACH TO THE GENERALIZATION OF HARDY-LITTLEWOOD INTEGRAL INEQUALITIES ON TIME SCALE	122
LÜTFİ AKIN ¹ , YUSUF ZEREN ²	122
SEVERAL CHARACTERIZATIONS OF THE BESSEL FUNCTIONS AND THEIR APPLICATIONS	123
MAHVİŞ ALI.....	123
PSEUDO-IMPLICATION AND FUZZY IDEALS IN BL-ALGEBRAS.....	124
MANAL A. ALQAHTANY ^{1,2} , DR TANİŞA BAGHWAN DALE ³	124
EXISTENCE OF SOLUTIONS TO FUZZY STOCHASTIC DIFFERENTIAL EQUATIONS	125
MAREK T. MALINOWSKI.....	125
A UNIFIED FENG-LIU TYPE RESULT IN RELATIONAL METRIC SPACES WITH AN APPLICATION.....	126
M.D. HASANUZZAMAN ¹ AND MOHAMMAD İMDAD ²	126
ANALYTICAL APPROXIMATE SOLUTIONS OF SOME NONLINEAR EVOLUTION EQUATIONS OF FRACTIONAL ORDER VIA A NOVEL FRACTIONAL VARIATIONAL ITERATION METHOD	127
M.D. ASADUZZAMAN ¹ AND FARUK ÖZGER ²	127
²DEPARTMENT OF ENGINEERING SCIENCES, İZMİR KATİP CELEBİ UNIVERSITY, 35620, İZMİR, TÜRKİYE,	127
STANCU-TYPE GENERALIZED Q-BERNSTEIN-KANTOROVICH OPERATORS INVOLVING BEZIER BASES.....	128

MD. NASIRUZZAMAN	128
THE MARCHENKO METHOD FOR THE DERIVATIVENLS EQUATION	129
MEHMET UNLU ¹ , RAMAZAN ERCAN ²	129
A NEW NOTE ON SUMMABILITY OF INFINITE SERIES AND FOURIER SERIES	130
HIKMET SEYHAN ÖZARSLAN ¹ AND MEHMET ÖNER ŞAKAR ²	130
A NOTE ON CONSTRAINED MULTIVARIATE LINEAR MODEL AND ITS REDUCED MODELS	131
MELEK ERIŞ BÜYÜKKAYA ¹ , NESRİN GÜLER ²	131
SOLITON SOLUTIONS OF THE (3+1)-DIMENSIONAL GENERALIZED KADOMTSEV-PETVIASHVILI EQUATION	132
MELİH CİNAR ¹	132
MODULAR GRAD-DIV STABILIZATION FOR LINEARLY EXTRAPOLATED BLENDED THREE-STEP DIFFERENTIATION SCHEME FOR THE INCOMPRESSIBLE NATURAL CONVECTION	133
MERVE AK ¹ , MINE AKBAŞ ²	133
SCATTERING ANALYSIS OF STURM-LIOUVILLE DIFFERENTIAL EQUATION WITH SIGN-CHANGING WEIGHT	134
MERVE GÖRGÜLÜ	134
A NEW TRANSFORM METHOD AND ITS APPLICATION	135
MERVE YÜCEL ¹ , OKTAY SH. MUKHTAROV ² , KADRIYE AYDEMİR ³	135
A FINITE ELEMENT METHOD FOR SOLVING 2D CONTACT PROBLEMS WITH TRESCA FRICTION LOW	136
DALAH MOHAMED ¹ , FADLIA BESMA ² , DERBAZI AMMAR ³	136
BERNSTEIN TYPE OPERATORS ON ELLIPTIC DOMAIN AND THEIR INTERPOLATION PROPERTIES	137
MOHAMMAD ILIYAS ¹ , ASIF KHAN ¹ AND MOHAMMAD MURSALEEN ²	137
FIXED POINT THEOREMS UNDER LOCALLY T -TRANSITIVE BINARY RELATIONS EMPLOYING MATKOWSKI CONTRACTIONS	138
MOHAMMAD ARIF ¹ , MOHAMMAD IMDAD ² , AFTAB ALAM ³	138
SELF ADAPTIVE INERTIAL YOSIDA APPROXIMATION ITERATIVE ALGORITHMS FOR SPLIT VARIATIONAL INCLUSION AND FIXED POINT PROBLEMS	139
MOHAMMAD DILSHAD	139
ON TWO SEQUENTIAL HYBRID FRACTIONAL Q-DIFFERENTIAL INCLUSIONS	140
MOHAMMAD ESMAEL SAMEI ¹	140
SOLVING A SIMPLE HARMONIC OSCILLATOR EQUATIONBY ADOMIAN DECOMPOSITION METHOD	141
MOHAMMED MOHSWN ¹ , YAHYA QAID ²	141
ADOMIAN DECOMPOSITION METHOD TO SOLVE THE SECOND ORDER ORDINARY DIFFERENTIAL EQUATIONS WITH CONSTANT COEFFICIENT.....	142
YAHYA QAID ¹ , MOHAMMED MOHSWN ²	142
COMPUTATIONAL ANALYSIS OF DEGREE-BASED TOPOLOGICAL DESCRIPTORS OF TRIPHENYLENE BENZENOID SYSTEM	143
MOHAMMED ALSHARAFI ¹ , YUSUF ZEREN ² , MUTLU AKAR ³ , ABDO ALAMERI ⁴	143
BLENDING TYPE APPROXIMATIONS OF SZ' ASZ-MIRAKJAN OPERATORS DEPENDING ON PARAMETERA.....	144

MOHD RAIZ ¹ , VISHNU NARAYAN MISHRA ² , NADEEM RAO ³	144
A NOVEL NUMERICAL MANNER FOR NON-LINEAR COUPLED VARIABLE ORDER REACTION-DIFFUSION EQUATION ...	145
MOHD KASHIF.....	145
APPROXIMATION BY NONLINEAR LUPAS TYPE BERNSTEIN OPERATOR OF MAX-PRODUCT KIND	146
MOHD SHANAWAZ MANSOORI ¹ , ASIF KHAN ²	146
NEW AUXILIARY FILLED FUNCTION WITHOUT PARAMETER FOR SOLVING OPTIMIZATION PROBLEMS	147
MUHAMMAD WASIM AWAN ¹ , AHMET ŞAHINER ² , TEMEL ERMIŞ ³ ,	147
RISK ASSESSMENT OF AUTONOMOUS VEHICLES IN TERMS OF PUBLIC HEALTH	148
MURAT KIRIŞCI ¹	148
CHILDHOOD CANCER RISK ANALYSIS WITH INTEGRATED DECISION-MAKING METHOD	149
MURAT KIRIŞCI ¹	149
SOME APPLICATIONS OF KRASNOSEL'SKII'S THEOREM FOR MULTI-VALUED OPERATORS	150
CESİM TEMEL ¹ , MÜBERRA SELAH ²	150
BASICITY OF DOUBLE SYSTEM OF EXPONENTS WITH LINEAR PHASE IN THE WEIGHTED LEBESGUE SPACE	151
NATAVAN NASIBOVA ¹ , NAZİR İSMAİLOV ²	151
ON HARMONIC SUMMABILITY OF DIFFERENCE SEQUENCES OF FRACTIONAL ORDER	152
NAZLİM DENİZ ARAL ¹ , HACER ŞENGÜL KANDEMİR ² , MİKAIL ET ³	152
EXISTENCE AND BLOW UP OF SOLUTIONS FOR A VISCOELASTIC PETROVSKY EQUATION WITH VARIABLE EXPONENTS	153
NEBİ YILMAZ ¹ , ERHAN PIŞKIN ²	153
STABILITY AND STABILIZATION OF THE IMPLICIT DIFFERENTIAL EQUATIONS IN HILBERT SPACES	154
NOR-EL-HOUDA BEGHERSA ¹ AND MEHDI BENABDALLAH ²	154
ON THE EXPONENTIAL STABILITY OF THE STATIONARY AND PERTURBED IMPLICIT EQUATIONS	155
NOR-EL-HOUDA BEGHERSA ¹ AND MEHDI BENABDALLAH ²	155
FINITE VOLUME METHOD FOR SEAWATER INTRUSION PROBLEM IN POROUS MEDIA	156
NOURA DJEDAİDİ ¹ , SOUAD GASMI ² AND FATMA ZOHRA NOURİ ³	156
EXISTENCE AND NONEXISTENCE RESULTS OF A FIFTH-ORDER BVP	157
NOURREDINE HOUARİ ¹ , FAOUZI HADDOUCHİ ^{1,2}	157
MULTIPLE PURSUERS AND ONE EVADER IN A PURSUIT DIFFERENTIAL GAME WITH GRONWALL-TYPE CONSTRAINTS	158
GAFURJAN İBRAGİMOV ¹ , OMONGUL EGAMBERGANOVA ² , RİSMAN MAT HASİM ³	158
GAUSS DECOMPOSITION OF \mathbb{Z}_3-GRADED QUANTUM GROUP $GL_q(2)$	159
SALİH ÇELİK ¹ , ÖZTÜRK SAVAŞ ²	159
INVERSE BOUNDARY PROBLEM FOR CONSTRUCTION OF THE RIGHT SIDE IN A THIRD ORDER TWO-DIMENSIONAL PSEUDOHYPERBOLIC EQUATION	160
YASHAR MEHRALİYEV ¹ , RENA SADIKHZADA ²	160


A NEW FAMILY OF SYMMETRIC FUNCTIONS OF BINARY PRODUCTS OF BIVARIATE MERSENNE LUCAS POLYNOMIALS WITH (P,Q)-NUMBERS	161
ROKIYA SAHALI ¹ AND ALI BOUSSAYOUD ²	161
QUASILINEAR PARABOLIC EQUATIONS WITH PARTIAL VMO COEFFICIENTS	162
ROSAMARIA RESCIGNO,LUBOMIRA SOFTOVA	162
IMPACT OF LARGE CLAIMS ON THE STABILITY BOUND OF A BIVARIATE RISK MODEL	163
SAFIA HOCINE ¹ , DJAMIL AÏSSANI ² , AÏCHA BARECHE ³	163
GLOBAL SOLVABILITY OF THE LAPLACE EQUATION IN WEIGHTED SOBOLEV SPACES	164
B. BILALOV, N.NASIBOVA, L.SOFTOVA, S. TRAMONTANO1	164
ON EVOLUTION EQUATIONS AND OPERATOR SEMIGROUPS IN BANACH SPACES	165
ERDAL GÜL ¹ ,SAMET KARASU ²	165
DATA DEPENDENCE RESULT FOR A NEW ITERATIVE ALGORITHM	166
SAMET MALDAR ¹ AND YUNUSATLAN ²	166
AN EXPLICIT MILSTEIN-TYPE SCHEME FOR INTERACTING PARTICLE SYSTEMS AND MCKEAN--VLASOV SDES WITH COMMON NOISE AND NON-DIFFERENTIABLE DRIFT COEFFICIENTS	167
SANI BISWAS ¹ ,CHAMAN KUMAR ¹ , NEELIMA ² , GONÇALO DOS REIS ³ , CHRISTOPH REISINGER ⁴	167
NUMERICAL INVESTIGATION OF FREE ROTOR AERODYNAMICS FOR SHIP-ROTOR INTERACTION	168
SARIH SARI ¹ ,ALI DOGRUL ² , SEYFETTIN BAYRAKTAR ³	168
SOME RESULTS ON b_3-SUBBALANCING NUMBERS.....	169
SELIN SARI ¹ ,GÜL KARADENİZ GÖZERİ ²	169
SPLIT QUATERNION SEQUENCES.....	170
SENA NUR AKTAŞ ¹ ,SALIM YÜCE ²	170
A BOUND SET OF CHAOTIC SATELLITE SYSTEM AND ITS APPLICATION IN CHAOS SYNCHRONIZATION	171
AYUB KHAN ¹ ,SHADAB ALI ² , ARSHAD KHAN ³	171
OPTIMAL POINTS IN $\mathbb{S}B\mathbb{S}$- METRIC SPACES ENDOWED WITH GRAPH	172
SHAGUN SHARMA AND SUMIT CHANDOK.....	172
SOLUTION OF NONLINEAR INTEGRAL EQUATIONS BY GENERALIZED CONTRACTIVE CONDITION.....	173
SHIV KANT TIWARI ¹	173
FRACTIONAL MATHIEU-DUFFING SYSTEM WITH THE STABILITY CONDITION AND CHAOS CONTROL	174
SHIVA ESHAGHI	174
DRBEM SOLUTIONS OF SINGULARLY PERTURBED MHD FLOW IN A SQUARE DUCT WITH VARIABLY CONDUCTING AND NO-SLIP/SLIPPING WALLS	175
SINEM ARSLAN ÖLÇER ¹ ,MÜNEVVER TEZER-SEZGIN ²	175
ASSESSMENT OF BANK LIQUIDITY RISK WITH ARTIFICIAL NEURAL NETWORKS	176
İBRAHİM DEMİR ¹ ,SINEM KÜBRA KÖSEN ²	176
ON MODIFICATION OF HIROTA METHOD THROUGH TRANSITION BETWEEN PARAMETERS	177
SUKRI KHARENG ¹ ,ÖMER ÜNSAL ²	177

STATISTICAL CONVERGENCE OF SEQUENCES OF SETS OF ORDER (A, B)	178
SUNIL K. SHARMA.....	178
PREDICTION OF HIV-1 NUCLEOSIDE REVERSE TRANSCRIPTASE RESISTANCE WITH ARTIFICIAL NEURAL NETWORKS ..	179
SUMEYYE YILMAZ ¹ ,MURAT SARI ² , HUSEYIN TUNC ³	179
PROPOSING A NEW DNA AND MUSIC POWERED CRYPTOLOGY TECHNIQUE	180
MUHARREM BAYRAK ¹ , HALIL YAKIT ² , KAYRA EGE ALTUN ³ , ŞERIF EFE DARTAR ⁴	180
A STUDY ON T-MAGNETIC CURVES	181
ŞERIFE NUR BOZDAĞ ¹ ,ŞEVVAL KUBLAY ²	181
SOLVABILITY IN THE SMALL AND SCHAUDER-TYPE ESTIMATES FOR HIGHER ORDER ELLIPTIC EQUATIONS IN BANACH-SOBOLEV SPACES	182
BILAL BILALOV ^{1,2} , YUSUF ZEREN ¹ ,SABINA SADIGOVA ^{2,3} , SEYMA CETIN ¹	182
ON LEFT DERIVATIONS OF GENERALIZED MATRIX ALGEBRAS	183
SEYMANUR AYDIN ¹ , NURCAN ARGAC ²	183
RESULTS ON EIGENVALUE PROBLEMS OF NONLINEAR CONFORMABLE FRACTIONAL DIFFERENTIAL EQUATIONS.....	184
ŞUAYIP TOPRAKSEVEN ¹	184
ON SOLUTION OF SOME DIFFERENTIAL EQUATIONS USING BICOMPLEX TRANSFORMATIONS	185
SERPİL HALICI ¹ ,SULE CURUK ²	185
GRAPHS OF BITONIC ALGEBRAS.....	186
REFET POLAT ¹ , ŞULE AYAR ÖZBAL ² AND SAADET ESKİİZMİRLİLER ³	186
INEQUALITIES OF HERMITE-HADAMARD-MERCER TYPE FOR TWICE DIFFERENTIABLE WITH APPLICATIONS.....	187
FRANCISCO MARTÍNEZ GONZÁLEZ ¹ , MUHAMMAD ZAFAR IQBAL ² , TALIB HUSSAIN ³	187
APPLIED MATHEMATICAL PHYSICS: SOLVING REAL-WORLD PROBLEMS.....	188
SEYYED ALI MOHAMMADIYEH ¹ AND HAMID PURHASANI ²	188
SOFT FRACTIONAL IDEALS AND SOFT STAR-OPERATIONS OF INTEGRAL DOMAINS	189
TASADUK RASHID MIR	189
NABLA CAPUTO TYPE Q-FRACTIONAL DIFFERENCE OPERATORS AND THEIR PROPERTIES	190
TAYLAN DEMİR ¹ ,MOHAMMED ESMAEL SAMEI ²	190
APPLICATIONS OF Q-MITTAG LEFFLER FUNCTION ON Q-FRACTIONAL DIFFERENTIAL OPERATORS	191
TAYLAN DEMİR ¹ ,MOHAMMED ESMAEL SAMEI ²	191
APPLICATIONS OF Q-MITTAG LEFFLER FUNCTION ON Q-FRACTIONAL DIFFERENTIAL OPERATORS	192
TAYLAN DEMİR ¹ ,MOHAMMED ESMAEL SAMEI ²	192
ON THE BASIS PROPERTY OF EIGENFUNCTIONS OF THE INDEFINITE STURM-LIOUVILLE PROBLEM IN LEBESGUE SPACES	193
TELMAN GASYMOV ¹ AND ULKER HASHIMOVA ¹	193
DESCRIPTION OF DOMAINS OF DEFINITION OF FRACTIONAL POWERS OF A SECOND –ORDER DIFFERENTIAL OPERATOR WITH INTEGRAL BOUNDARY CONDITIONS	194
TELMAN GASYMOV ¹ AND REYHAN TAGHIYEVA ¹	194

WEIGHTED HARDY'S INEQUALITY REVISITED	195
TUĞÇE ÜNVER YILDIZ1, UĞUR CANYILDIRIM2	195
SEQUENTIAL T_3 (REGULAR HAUSDORFF) SPACES	196
TUNÇAR ŞAHAN ¹ , ELANUR ÖZCAN ²	196
SOME BASIC RESULTS ON GROUPS UP TO CONGRUENCE RELATIONS	197
TUNÇAR ŞAHAN ¹ , AHMET EREN KILIÇ ²	197
ON BOUNDEDNESS OF THE RIESZ POTENTIAL AND ITS COMMUTATORS IN LOCAL ORLICZ-MORREY TYPE SPACES	198
MENDYBAY TURGUMBAYEV ¹ AND AIDOS ADILKHANOV ²	198
ON A CLASS OF NONLOCAL POROUS MEDIUM EQUATIONS OF KIRCHHOFF TYPE	199
UĞUR SERT	199
CONSTRUCTION OF THE ASYMPTOTICS OF THE SYSTEM OF FUNDAMENTAL SOLUTIONS FOR A CLASS OF UNUSUAL EQUATIONS	200
VAGIF MASTALIYEV ^{1,2} AND ULVIYA ISKANDAROVA ¹	200
A NEW TYPE ALGORITHM FOR FINDING A COMMON SOLUTION OF CAYLEY INCLUSION AND J-FIXED POINT PROBLEM IN HILBERT SPACES	201
UQBA RAFAT ¹	201
ON THE SOLUTION OF THE MIXED PROBLEM POSED FOR A CLASS OF EQUATIONS DEGENERATED BY THE TYPE	202
VAGIF MASTALIYEV ^{1,2} AND BAHRUZ AGARZAYEV ³	202
BLOW UP OF SOLUTIONS FOR A LOGARITHMIC PETROVSKY EQUATION WITH DAMPING TERM	203
YAVUZ DİNÇ ¹ , ERHAN PIŞKIN ² , CEMİL TUNÇ ³	203
REGULAR AND SEMIREGULAR HYPERMODULES	204
YILDIZ AYDIN ¹ , BURCU NIŞANCI TÜRKMEN ²	204
SOME FIXED-POINT THEOREMS FOR CONTINUOUS FUNCTIONS	205
YUNUS ATALAN ¹ AND SAMET MALDAR ²	205
ON SOME DIFFERENT REPRESENTATIONS OF Q-CATALAN NUMBERS AND THEIR SUMS	206
SERPİL HALICI ¹ , ZEHRA BETÜL GÜR ²	206
A TAUBERIAN THEOREM FOR THE WEIGHTED MEAN SUMMABILITY METHOD OF DOUBLE SEQUENCES IN ORDERED SPACES	207
ZERRİN ÖNDER ŞENTÜRK	207
FUNDAMENTAL CONCEPTS IN VARIABLE LEBESGUE SPACES ASSOCIATED WITH LAPLACE-BESSEL DIFFERENTIAL OPERATOR	208
ZEYNEP TERCAN ¹ , ESRA KAYA ²	208
THE EXACT SOLUTIONS OF SOME DIFFERENCE EQUATIONS VIA NARAYANA NUMBERS	209
ZEYNEP AKMEŞE ¹ , MELİH GÖCEN ²	209
STATISTICAL CONVERGENCE OF ASYMMETRIC METRIC VALUED SEQUENCES WITH DEFERRED CÉSÀRO MEAN	210
ZEYNEP HANDE TOYGANÖZÜ	210
THREE WEIGHTED INEQUALITY FOR A SUPERPOSITION OF SOME OPERATORS	211

YERNIYASHOVA ZHADYRA	211
ON THE BILINEAR RIESZ POTENTIAL AND ITS COMMUTATORS ON GENERALIZED ORLICZ-MORREY SPACES	212
ZHANAT ZHULDASSOV ¹ , TALGAT AKHAZHANOV ² , DAUREN MATIN ²	212
RECURSIVE ESTIMATION OF THE CONDITIONAL DISTRIBUTION FUNCTION ON RIEMANNIAN MANIFOLDS	213
BOUADJEMI ABDELKADER ¹	213
EXISTENCE AND UNIQUENESS OF THE STRONG SOLUTION OF THE TIME FRACTIONAL INTEGRO-DIFFERENTIAL EQUATION WITH INTEGRAL BOUNDARY CONDITIONS.....	214
AHCENE MERAD ¹ , OUARDA BENMANSEUR ²	214
BANACH HARDY CLASSES AND SOME PROPERTIES	215
VAFA ALILI	215
AN INVERSE COEFFICIENT PROBLEM FOR A TWO-DIMENSIONAL HYPERBOLIC EQUATION WITH INTEGRAL OVERDETERMINATION	216
ELVIN AZIZBAYOV ¹ AND GULNARA SHUKUROVA ²	216
AN INVERSE COEFFICIENT PROBLEM FOR A TWO-DIMENSIONAL HYPERBOLIC EQUATION WITH INTEGRAL OVERDETERMINATION	217
ELVIN AZIZBAYOV ¹ AND GULNARA SHUKUROVA ²	217
A COMPACT FINITE DIFFERENCE METHOD FOR FRACTIONAL EQUATION BASED ON CAPUTO DERIVATIVE: CODES MATLAB.....	218
BENDIB ISSAM ¹ , DALAH MOHAMED ² , BOUDJEDOUR ALLAOUA ³	218
RECURSIVE ESTIMATION OF THE CONDITIONAL DISTRIBUTION FUNCTION ON RIEMANNIAN MANIFOLDS	219
BOUADJEMI ABDELKADER ¹	219
NUMERICAL ANALYSIS APPLIED TO STIFFENED SHELL STRUCTURES.....	220
HAMADI DJAMAL ¹ , LEGOUIRAH MARIA ² AND ABDULRAHMAN AL-NADHARI ³	220
AN INTERIOR-POINT ALGORITHM FOR CQP BASED ON TRIGONOMETRIC KERNEL FUNCTION	221
EL AMIR DJEFFAL ¹ AND RANDA CHALEKH ²	221
ON QUALITATIVE PROPERTIES OF SOME NONUNIFORM ELLIPTIC EQUATIONS.....	222
FARMAN MAMEDOV ¹	222
CHELYSHKOV POLYNOMIALS WITH APPLICATIONS IN APPROXIMATION THEORY	223
IMRAN TALIB.....	223
ON Λ-FRACTIONAL VARIATIONAL CALCULUS	224
K.A.LAZOPOULOS	224
ON A BOUNDARY VALUE PROBLEM FOR NONLINEAR DIFFUSION EQUATION WITH INTEGRAL CONDITION OF THE SECOND KIND	225
MATLAB SALIMOV, AFARIM SALIMOVA	225
COMPILATION OF METHOD FOR SOLVING PARTIAL DIFFERENTIAL EQUATIONS OF THE FIRST-ORDER BY THE ADOMIAN DECOMPOSITION METHOD	226
MUBARK RAANA ¹ , YAHYA Q. HASAN ²	226

BLOW UP OF SOLUTIONS FOR A VISCOELASTIC WAVE EQUATIONS WITH BALAKRICHNAN-TAYLOR DAMPING AND A DELAY TERM	227
NOURI BOUMAZA ¹ , BILLEL GHERAIBIA ²	227
DYNAMICS OF DISCRETIZED ROSENZWEIG–MACARTHUR PREDATOR–PREY MODEL WITH A HOLLING TYPE II FUNCTIONAL RESPONSE FOR CONSTANT IMMIGRATION	228
PRADEEP MALIK	228
APPROXIMATION ON BIVARIATE OF DURRMEYER OPERATOR BASED ON BETA FUNCTION	229
MOHD RAIZ ¹ , RUCHI SINGH RAJAWAT ¹ , LAKSHMI NARAYAN MISHRA ³ , VISHNU NARAYAN MISHRA ¹	229
SOLUTION IN THE SMALL AND INTERIOR SCHAUDER-TYPE ESTIMATE FOR THE M-TH ORDER ELLIPTIC OPERATOR IN MORREY-SOBOLEV SPACES	230
BILAL BILALOV ^{1,2} , YUSUF ZEREN ² , SABINA SADIGOVA ^{1,3}	230
PROPERTIES OF SOLUTIONS TO LINEAR DIFFERENTIAL EQUATIONS NEAR A SINGULAR POINT	231
MAZOUZ SAID ¹ , SAADA HAMOUDA ²	231
STUDY OF ONEDIMENSIONAL MIXED PROBLEM FOR ONE CLASS OF THIRD ORDER NONLINEAR PSEUDOPARABOLIC EQUATIONS	232
SAMED ALIYEV AND ARZU ALIYEVA	232
SOME QUESTIONS OF HARMONIC ANALYSIS IN WEIGHTED MORREY TYPE SPACES	233
SAMIA BIBI	233
A MATHEMATICAL MODEL FOR THE BREAST CANCER DIAGNOSIS AND ITS PROGNOSIS USING FUZZY EXPERT SYSTEM	234
SANJEEV KUMAR	234
SOME NUMERICAL PRESENTATION FOR A QUADRATIC MAP	235
SELMANI WISSAME ¹ , DJELLIT ILHEM ²	235
PROBABILITY IN MATHEMATICAL PHYSICS: THEORY AND APPLICATIONS	236
SEYYED ALI MOHAMMADIYEH ¹ AND HAMID PURHASANI ²	236
PROBLEMS OF MATHEMATICAL EDUCATION IN THE CURRICULUM IN THE CAUCASIAN COUNTRIES	237
Shahin Aghazade ¹ , Gunel Novruzova ² , Atilla Bingol ³	237
ON STRONG SOLVABILITY OF THE DIRICHLET PROBLEM FOR A CLASS OF SEMILINEAR ELLIPTIC EQUATIONS	238
SH. YU. SALMANOVA	238
PARABOLIC FRACTIONAL INTEGRAL OPERATORS WITH ROUGH KERNELS IN PARABOLIC LOCAL GENERALIZED MORREY SPACES	239
SHEMSIYYE MURADOVA ^{1,2}	239
BIFURCATION ANALYSIS AND COMPLEX DYNAMICS OF A NEW FRACTIONAL SYSTEM	240
SHIVA ESHAGHI	240
BASICITY OF SINE-COSINE SYSTEMS IN GRAND LEBESGUE SPACE	241
TURAL HAGVERDI	241
DIFFERENT VERSIONS OF DIRICHLET PROBLEMS FOR LAPLACE EQUATION IN NON-STANDARD FUNCTION SPACES	242
BILAL BILALOV ^{1,2} , NIGAR AHMADZADEH ¹ , TARLAN GARAYEV ^{1,3}	242



ICOMAA 2023

INVITED TALKS

Holmstedt's formula for the K-functional: the limit case $\theta_0 = \theta_1$

Amiran Gogatishvili

(joint work with Irshaad Ahmed and Alberto Fiorenza)

Institute of Mathematics of the Czech Academy of Sciences
Prague, Czech Republic
E-mail: gogatish@math.cas.cz

Abstract

We consider K-interpolation spaces involving slowly varying functions, and derive necessary and sufficient conditions for a Holmstedt-type formula to be held in the limiting case $\theta_0 = \theta_1 \in \{0,1\}$. We also study the case $\theta_0 = \theta_1 \in (0,1)$. Applications are given to Lorentz-Karamata spaces, generalized gamma spaces and Besov spaces.

Keywords: K-functional, slowly varying functions, K-interpolation spaces, Holmstedt's formula, reiteration

References:

1. I. Ahmed, A. Fiorenza and A. Gogatishvili, Holmstedt's formula for K-functional: the limit case $\theta_n = \theta_1$. Institute of Mathematics CAS, Preprint No. 60-2022, Prague 2022, Preprint arXiv:2210.12377 22 pages, accepted in Math. Nachr.

This research was supported by Czech Academy of Sciences (RVO 67985840), by Czech Science Foundation (GAČR), grant no: 23-04720S, and by Shota Rustaveli National Science Foundation of Georgia (SRNSFG), grant no: FR21-12353.

Some regularity results for a class of elliptic equations with lower order terms and Orlicz growth

Claudia Capone

Istituto per le Applicazioni del Calcolo "M. Picone" - C.N.R

c.capone@na.iac.cnr.it

Abstract

Let us consider the Dirichlet problem

$$\begin{cases} \operatorname{div} A(x, Du) + b(x) \frac{\phi'(|u|)}{|u|} u = f \\ u = 0 \end{cases} \quad \partial\Omega$$

with $\Omega \subset \mathbb{R}^n, n > 2$, a bounded open set and $A(x, \xi)$ a function satisfying growth conditions, with respect to the second variable, expressed through an N-function ϕ . We study the regularity properties of the weak solutions $u: \Omega \rightarrow \mathbb{R}$ to these problems. More precisely, if $\phi(t) = t^p, p=2$, we establish higher differentiability of solutions to our problem under a Sobolev assumption on the partial map $x \rightarrow A(x, \xi)$. In the case of a general N-function ϕ , we prove higher differentiability and higher integrability of the gradient, under mild assumptions on the data, by assuming $A(x, \xi)$ depending on x through an Hölder continuous function. The novelty, in both case, is that we take advantage from the regularizing effect of the lower order term, due to the interplay between $b(x)$ and $f(x)$.

Keywords: Higher differentiability, higher integrability, Orlicz spaces, Besov-Orlicz spaces, regularizing effect, boundedness of solutions.

References:

1. C. Capone, T. Radice, Higher differentiability for solutions to a class of elliptic equations with lower order terms, *Journal of Elliptic and Parabolic Equations*, 6 (2), (2020), 751--771.
2. C. Capone, On the regularity of solutions of degenerate PDEs with lower order terms, *J. Math. Anal. Appl.* 504, (2021) 125394.
3. C. Capone, A. Passarelli di Napoli, Regularity results to a class of elliptic equations with explicit ϕ -dependence and Orlicz growth, (submitted 2022).

C. Capone, A. Passarelli di Napoli, Regularity results to a class of elliptic equations with explicit ϕ -dependence and Orlicz growth, (submitted 2022).

Matrix weights in variable Lebesgue spaces: the case of averaging operators

David Cruz-Uribe, OFS
The University of Alabama

Abstract

In the 1990s, Nazarov, Treil and Volberg introduced a generalization of the Muckenhoupt A_p weights, replacing the scalar weights w with matrix weights W : that is, measurable, $d \times d$ matrix valued functions whose values are self-adjoint and positive semi-definite almost everywhere. Beginning with their results, there has been a great deal of work done to prove inequalities of the form

$$\|W(T\vec{f})\|_p \leq C\|W\vec{f}\|_p,$$

where T is a Calderón-Zygmund singular integral or some other classical operator of harmonic analysis.

A natural question is to extend these ideas to the scale of variable Lebesgue spaces. These are spaces where the constant exponent p is replaced by an exponent function $p(\cdot)$, with the norm given by the Luxemburg norm:

$$\|f\|_{p(\cdot)} = \inf \left\{ \lambda > 0 : \int_{\mathbb{R}^n} \left(\frac{|f(x)|}{\lambda} \right)^{p(x)} dx \leq 1 \right\}.$$

The scalar theory of Muckenhoupt weights was extended to these spaces by Cruz-Uribe, Fiorenza, and Neugebauer; Diening and Hästö, and others; however, no work to date has been done on matrix weights in this setting.

We will define a new class of matrix $\mathcal{A}_{p(\cdot)}$ weights which reduces to the scalar class $A_{p(\cdot)}$ defined previously when $d = 1$, and when $p(\cdot) = p$ is constant to the class of matrix weights \mathcal{A}_p defined by Nazarov, Treil and Volberg.

The ultimate goal of this project is to prove that singular integrals are bounded on the space $L^{p(\cdot)}(W)$ when $1 < p_- \leq p_+ < \infty$ and $p(\cdot)$ satisfies the log-Hölder continuity conditions. We begin with an important first step: averaging operators. We prove that this class is necessary and sufficient for averaging operators to be bounded on $L^{p(\cdot)}(W)$. As an application we prove that approximations of the identity converge in this space, and then use this to show that the $H = W$ theorem of Meyers and Serrin can be extended to this setting. This generalizes work by Cruz-Uribe, Moen and Rodney in the constant exponent case.

Results in this talk are joint work with my student Michael Penrod.

Generalised fractional operators with applications in biology


Dumitru Baleanu^{1,2}

¹Department of Mathematics, Cankaya University, Ankara, Turkiye

²Institute of Space Sciences, Magurele-Bucharest, Romania

Abstract

Fractional calculus deals with the study of fractional order integral and derivative operators over real or complex domains, and their real world applications. In my talk I will provide the applications of the generalized operators in biology. Several illustrative examples will be presented.



ICOMAA 2023

Maximal regularity for non-autonomous problems with a priori assumptions

Peter Hästö
University of Turku

Abstract

Vector-valued generalized Orlicz spaces can be divided into anisotropic, quasi-isotropic and isotropic. In isotropic spaces, the Young function depends only on the length of the vector, i.e. $\Phi(v) = \varphi(|v|)$. In the quasi-isotropic case $\Phi(v) \approx \varphi(|v|)$ so the dependence is via the length of the vector up to a constant. In the anisotropic case, there is no such restriction, and the Young function depends directly on the vector. Jihoon Ok and I obtained maximal local regularity results of weak solutions or minimizers of

$$\operatorname{div} A(x, Du) = 0 \quad \text{and} \quad \min_u \int_{\Omega} F(x, Du) dx,$$

when A or F are general quasi-isotropic Young functions. We established local $C^{1,\alpha}$ -regularity for some $\alpha \in (0, 1)$ and C^α -regularity for any $\alpha \in (0, 1)$ of weak solutions and local minimizers with varying a priori assumptions such as $u \in L^s$. Previously known regularity results are included as special cases.

Solvability in the small of higher order elliptic equations in weighted Sobolev-Banach spaces

B. Bilalov, S. Sadigova, L. Softova, Y. Zeren

Department of Mathematics, University of Salerno
lsoftova@unisa.it

Abstract

We consider m -th order linear, uniformly elliptic equations $Lu=f$ with non-smooth coefficients in Banach-Sobolev spaces generated by weighted general Banach Function Spaces (BFS) on a bounded domain.

Supposing boundedness of the Hardy-Littlewood Maximal and Calderón-Zygmund singular operators in weighted BFS we obtain solvability in the small in Sobolev-Banach weighted spaces and establish interior Schauder type a priori estimates for the corresponding elliptic operator.

Keywords: Linear higher order elliptic equations, Banach weighted function spaces, Sobolev spaces, a priori estimates.

References:

1. Bennett C., Sharpley R., Interpolation of Operators, Academic Press, 1988, 469 p.
2. Bers L., John F., Schechter M., Partial Differential Equations, Lectures in Applied Mathematics, 3A. Providence, R.I.: AMS XIII, 343 p., 1979.
3. Bilalov B.T., Sadigova S.R., On solvability in the small of higher order elliptic equations in grand-Sobolev spaces, *Compl. Var. Elliptic Equ.*, 66, No. 12 (2021), 2117--2130.
4. Bilalov B., Sadigova S., Softova L., Zeren Y., Solvability in the small and interior Schauder type estimates for higher order elliptic equations in weighted Banach spaces, [arXiv:2212.11227](https://arxiv.org/abs/2212.11227)

Some non-matrix summability methods and Korovkin type approximation theorems

M. Mursaleen

Department of Mathematics

Aligarh Muslim University

Aligarh-202002 India

E-mail:mursaleenm@gmail.com

Abstract

Korovkin type approximation theorems are useful tools to check whether a given sequence $(L_n)_{n \geq 1}$ of positive linear operators on $C[0, 1]$ of all continuous functions on the real interval $[0, 1]$ is an approximation process. That is, these theorems exhibit a variety of test functions which assure that the approximation property holds on the whole space if it holds for them. Such a property was discovered by Korovkin in 1953 for the functions $1, x$ and x^2 in the space $C[0, 1]$ as well as for the functions $1, \cos$ and \sin in the space of all continuous 2π -periodic functions on the real line. In this paper, we use the notion of almost convergence and statistical convergence to prove the Korovkin type approximation theorems for the test functions $1, e^{-x}, e^{-2x} ..$

Cones generated by a generalized fractional maximal function

Nurzhan Bokayev¹, Amiran Gogatishvili², Azhar Abek³

^{1,3}Department of Mathematics, Eurasian National University, Astana, Kazakhstan,

²Institute of Mathematics of the Czech Academy of Sciences, Prague,

bokayev2011@yandex.ru

gogatish@math.cas.cz

azhar.abekova@gmail.com

Abstract

In this work we consider the space of generalized fractional-maximal function, constructed on the basis of a rearrangement-invariant space. Four types of cones generated by a nonincreasing rearrangement of a generalized fractional-maximal function and equipped with positive homogeneous functionals are constructed. Questions of mutual covering of such cones are considered. The question of embedding the space of generalized fractional-maximal function in a rearrangement-invariant space is investigated. The cones from nonincreasing rearrangements of generalized Riesz potentials were previously considered in the papers [1],[2].

Keywords: Rearrangement-invariant spaces, non-increasing rearrangements, cones generated by generalized fractional-maximal function, covering of cones.

References:


1. Goldman M.L, On the cones of rearrangements for generalized Bessel and Riesz potentials, Complex Variables and Elliptic Equations. Vol. 55, No.8-10 (2010), 817--832.
2. Bokayev N. A., Goldman M. L., & Karshygina G. Zh. Cones of functions with monotonicity conditions for generalized Bessel and Riesz potentials, Mathematical Notes. 104, No.3 (2018), 356--373.

Weighted Morrey spaces

Yoshihiro Sawano
Department of Mathematics
Chuo University

Abstract

Weighted function spaces arise naturally in many contexts of mathematics. Roughly speaking, weighted measure spaces, which generate weighted function spaces, are useful although they break some symmetries. We note that a group can be used to determine the symmetry of a space. In fact, a group action allows the symmetry of the spaces to be measured. For example, $O(n)$, the group of all orthonormal matrices, acts naturally on R^n , and R^n acts on itself as a translation. Hence, we can say that R^n is symmetric enough.



ICOMAA 2023

CONTRIBUTED TALKS

On the some relations ship between positive and positive p-summing operators

Abdelmoumen TIAIBA¹

¹Department of Physics, Mohamed Boudiaf University of (M'sila), Algeria
abdelmoumen.tiaiba@univ-msila.dz

Abstract

In this paper , with a new study of some relations between positive, regular and positive p- summing operators we show as first contribution a theorem which give a necessary and sufficient conditions that the KB -Banach space Y has the Radon-Nkodym propreity is that all additive operator frome $L_{p'}(\gamma)$, to Y be representable by a function $f \in L_p^+(\gamma, Y)$ such that $\frac{1}{p} + \frac{1}{p'} = 1$. In the second part , we prove some results to positive and increasing sublinear operators.

Keywords: Banach lattice, Hahn Banach theorem, linear operator, sublinear operator, summing linear operator.

References:

1. O. Blasco, Boundary values of vector valued harmonic functions considered as operators, Studio Math. 86 (1987), 19-33.
2. O. Blasco. Positive p-summing operators on $L_{\{p\}}$ -spaces. Proc. Amer. Math. Soc. 100, 2 (1987), 275-280.
3. L. Mezrag and A. Tiaiba, On the sublinear operators factoring through $L_{\{q\}}$. Int. J. Math. Math. Sci. 50 (2004), 2695-2704.

ICOMAA 2023

An Algorithm for Relaxing Multicurves on Small Genus Non-Orientable Surfaces

Ferihe ATALAN¹, Abdullah BAYKAL², Öykü YURTTAŞ³

¹Department of Mathematics, Atilim University

²Department of Mathematics, Dicle University

³Department of Mathematics, Dicle University

ferihe.atalan@atilim.edu.tr

baykal@dicle.edu.tr

saadet.yurttas@dicle.edu.tr

Abstract

A multicurve on a given surface S is a disjoint union of essential simple closed curves on S up to isotopy. In this talk, we consider multicurves in the case where S is an n -punctured non-orientable surface of genus one with one boundary component, and present an algorithm for solving an interesting combinatorial problem regarding multicurves on S . In particular, we describe an efficient algorithm for relaxing a multicurve in S making use of generalized Dynnikov Coordinates and update rules which describe the action of the mapping class group (group of isotopy classes of homeomorphisms) of S in terms of generalized Dynnikov Coordinates.

Keywords: Non-orientable surface, Generalized Dynnikov coordinates, Multicurves

References:

1. Artin, E., *Theorie der Zöpfe*, Abh. Math. Sem. Univ. Hamburg, **4**,(1925),47–72.
2. Artin, E., *Theory of braids*, Ann. of Math., **48**(2),(1947),101–126.
3. Dynnikov, I., *On a Yang-Baxter mapping and the Dehornoy ordering*, Uspekhi Mat. Nauk, **57**(3(345)),(2002),151–152.
4. Dehornoy, P., Dynnikov, I., Rolfsen, D., and Wiest, B., *Ordering braids*, Mathematical Surveys and Monographs, vol. 148, American Mathematical Society, Providence, RI, 2008
5. Hall, T., Yurttaş, S.Ö., *On the topological entropy of families of braids*, Topology Appl., **156**(8),(2009),1554–1564.

ICOMAA 2023

SOME PROPERTIES OF GRILL TOPOLOGICAL SPACES VIA G^{α}_{ω} $O(X)$

Amin Saif¹, A. Mahdi²

¹ Department of Mathematics, Faculty of Sciences, Taiz University, Taiz, Yemen.

² Department of Mathematics, Faculty of Education, University of Saba Region, Mareb, Yemen.

alsanawyamin@yahoo.com

ahidere@gmail.com

Abstract

The main idea is introducing and investigating a new class of open sets in grill topological spaces, namely G^{α}_{ω} -sets, which is considered as a strong form of class of G^{α}_{ω} -open sets and it is induced topologies by the collection of G^{α}_{ω} -open sets. Next, we study the separation axioms in the collection of G^{α}_{ω} -open sets.

Keywords: Grill topological space, Induced topology, Separation axioms.

References:

1. B. Roy and M. N. Mukherjee, On a typical topology induced by a grill, Soochow J. Math., Vol. 33, No. 4, (2007) 771-786.
2. A. Saif, A. Mahdi and K. M. Hamadi., G^{α}_{ω} -open Open sets in Grill Topological Spaces, International Journal of Computer Applications, Vol. 183, No.13, (2021) 1-5.
3. A. Saif, A. Mahdi and K. M. Hamadi., (2021), On Continuity Property of Grill Topological Spaces into Topological Spaces via G^{α}_{ω} -open sets, International Journal of Applied Mathematical Sciences, Vol. 14, No. 1, (2021) 1-16.
4. A. Saif, M. Al-Hawmi and B. Al-Refaei, On G^{α}_{ω} -open sets in grill topological spaces, Journal of Advance in Mathematics and Computer Science, 35(6),(2020) 132-143.

ICOMAA 2023

Bounded finite-time stabilizing positional controls via the controllability function as motion time

Abdon E. Choque-Rivero¹,

¹Instituto de Física y Matemáticas, Universidad Michoacana de San Nicolás de Hidalgo, México
abdon.choque@umich.mx

Abstract

For the canonical system, a set of bounded finite-time stabilizing positional controls is constructed. This set depends on a parameter which belong to a maximal interval whose terminal points are explicitly calculated. We use the Korobov's controllability function which is Lyapunov type function. In our investigation, the controllability function is exactly the motion time from a given initial x_0 to the origin.

Keywords: Bounded controls, finite-time stabilization, controllability function.

References:

1. A.E.Choque-Rivero, Korobov's Controllability function as motion time. Extension of the solution set of the synthesis problem, accepted to appear in *Mat. Fiz. Anal. Geom.* (2023) (31 pages).
2. A.E.Choque-Rivero, Extended set of solutions of a bounded finite-time stabilization problem via the controllability function, *IMA J. Math. Control Inf* 38(4) (2021) 1174–1188.
3. A.E.Choque-Rivero, V.I. Korobov, V.O. Skoryk, Controllability function as time of motion I. *Mat. Fiz. Anal. Geom.* (in Russian) 2(11) (2004) 208–225.



ICOMAA 2023

Two-Weighted Inequalities for Some Classical Operators and Their

Commutators in the Modified Morrey Spaces

Abdulhamit Kucukaslan

Department of Aerospace Engineering, Ankara Yıldırım Beyazıt University

a.kucukaslan@aybu.edu.tr

Abstract

In this talk, we prove the boundedness of some classical operators as the generalized fractional maximal operator and the generalized Riesz potential operator and their commutators in the weighted modified Morrey spaces and the weighted weak modified Morrey spaces by using two-weighted inequalities. That is, we give sufficient conditions for the boundedness of the generalized Riesz potential and the generalized fractional maximal operator and also their commutators in the weighted modified Morrey spaces and weighted weak modified Morrey spaces. Our weighted functions belong to the Muckenhoupt-Wheeden classes and the function b is in the Bounded Mean Oscillation BMO spaces. The generalized Riesz potential operator was initially investigated in [1]. The boundedness of our two-operators in the Morrey-type spaces is studied by many authors, for instance, see [2, 3, 4, 5].

Keywords: *Weighted Inequalities, Generalized Riesz potential operator, Generalized fractional maximal operator, Commutators, Modified Morrey spaces.*

References:

1. Gadjiev A.D., On generalized potential-type integral operators, *Funct. Approx. Comment. Math.* 25: 37-44, 1997.
2. Guliyev V.S., Hasanov J.J., and Zeren Y., Necessary and sufficient conditions for boundedness of the Riesz potential in modified Morrey spaces, *J. Math. Inequalities*, 5(4): 491-506, 2011.
3. Guliyev V.S., Ismayilova A.F., Kucukaslan A., and Serbetci A., Generalized fractional integral operators on generalized local Morrey spaces, *J. Funct. Spaces*, Article ID 594323, 8 pages, 2015.
4. A. Kucukaslan, Equivalence of norms of the generalized fractional integral operator and the generalized fractional maximal operator on generalized weighted Morrey spaces, *Ann. Funct. Anal.* 11: 1007-1026, 2020.
5. E. Nakai, On generalized fractional integrals, *Taiwanese J. Math.*, 5: 587-602, 2001.

Fekete-Szegő Problem for Certain Subclasses of Analytic Functions Associated with The Combination of Differential Operators

Adem Kızıltepe¹, Erhan Deniz², Sercan Kazımoğlu³

^{1,2,3}Department of Mathematics, Faculty of Science and Letters, Kafkas University, Kars-Turkey

admzkzlt@gmail.com

edeniz36@gmail.com

srcnkzmglu@gmail.com

Abstract

In this work, we introduce and study some new subclasses of analytic functions defined by the combination of Noor and Deniz-Özkan differential operators, and obtain coefficient estimates and Fekete-Szegő inequalities for these new subclasses.

Keywords: Fekete-Szegő problem, Analytic functions, Starlike and convex functions of complex order, Noor and Deniz-Özkan differential operator.

References:

1. Çağlar, M. and Orhan, H. (2021). Fekete-Szegő problem for certain subclasses of analytic functions defined by the combination of differential operators. Bol. Soc. Mat. Mex., 27, 41.
2. Deniz, E. and Orhan, H. (2010). The Fekete-Szegő problem for a generalized subclass of analytic functions. Kyungpook Math. J., 50, 37–47.
3. Fekete, M. and Szegő, G. (1933). Eine Bemerkung über ungerade schlichte Funktionen. J. Lond. Math. Soc., 8, 85–89.
4. Kazımoğlu, S. and Deniz, E., (2020). Fekete-Szegő problem for generalized bi-subordinate functions of complex order. Hacettepe Journal of Mathematics and Statistics, 49(5), 1695-1705.
5. Orhan, H., Deniz, E. and Çağlar, M. (2012). Fekete-Szegő problem for certain subclasses of analytic functions. Demonstr. Math., 45(4), 835–846.

ICOMAA 2023

Inverse Scattering Problem for the Schrödinger Equation with an Additional Growing Potential on the Entire Axis

Davud Orucov¹ and Agil Khanmamedov^{1,2}

¹*Department of Mathematics, Baku Engineering University,*
dorudjov@gmail.com

²*Department of Applied Mathematics, Baku State University*
agil_khanmamedov@yahoo.com

Abstract

The Schrödinger equation with an infinitely growing potential at the right end and disappearing at the left end is considered. The method of transformation operators is used to study the direct and inverse problems of scattering theory. The main integral equations of the inverse problem are obtained. The unique solvability of the main equations is proved.

Keywords: Schrödinger equation, harmonic oscillator, scattering data, inverse scattering problem, main integral equations.

References:

1. Chelkak D., Kargaev P., Korotyaev E., An Inverse Problem for an Harmonic Oscillator Perturbed by Potential: Uniqueness, *Lett. Math. Phys.* **64**:1 (2003), 7–21.
2. Chelkak D., Kargaev P., Korotyaev E., Inverse problem for harmonic oscillator perturbed by potential, characterization, *Comm. Math. Phys.*, **249**:1(2004), 133-196.
3. Orucov D.H., On the transformation operator for the Schrodinger equation with an additional increasing potential, *Proceedings of the Institute of Mathematics and Mechanics of NAS of Azerbaijan*, **48**:2(2022), 311-322.
4. Guseynov G.M., Khanmamedov A.Kh., Mamedova A.F., Inverse spectral problem for the Schrodinger equation with an additional linear potential, *Theoretical and Mathematical Physics*, **202**(1)(2018), 58–71.

ICOMAA 2023

Doubly Diffracted Fields Stimulated by Non-symmetrical Edges at the Junction

Alaa Serhan¹, Feray Hacivelioglu²

^{1,2}Department of Mathematics, Gebze Technical University

a.serhan2020@gtu.edu.tr

f.dagidir@gtu.edu.tr

Abstract

This study investigates the doubly diffracted fields generated by the strip that has half soft and half hard boundary conditions on its upper face, and totally soft boundary condition on its lower face. The spectral iteration technique with the Wiener-Hopf method is employed in the study to examine the doubly diffracted fields at the junction on the upper face of the strip. The effect of the non-symmetrical end edges of the strip on the doubly diffracted field at the edge (junction) on the upper face of the strip is presented graphically for the various parameters of the problem.

Keywords: Doubly diffracted field, soft and hard boundary conditions, spectral iteration technique, Wiener–Hopf method.

References:

1. T. B. A. Senior, “Scattering by resistive strips,” *Radio Sci.* 14 (5), 911–924 (1979).
2. A. Büyükaksoy and G. Uzgoren, “Secondary diffraction of a plane wave by a metallic wide strip residing on the plane interface of two dielectric media,” *Radio Sci.* 22 (2), 183–191 (1987).
3. A. Büyükaksoy and A. Alkumru, “Multiple diffraction of plane waves by a soft/hard strip,” *J. Eng. Math.* 29, 105–120 (1995).
4. G. Cinar, “Wiener–Hopf analysis of a plane wave diffraction by an impedance strip attached on a perfectly conducting half-plane,” *Electromagnetics* 29 (2), 165–184 (2009).
5. F. Hacivelioglu, “Diffraction of Sound Plane Waves by a Strip with Mixed Soft-Hard Faces” *Computational Mathematics and Mathematical Physics*, 62 (6) 1020-1031 (2022).

This work is supported by TÜBİTAK. Project No: 1002 – Short Term R&D Funding Program, 122F238

ICOMAA 2023

Optimal linear approximation and isometric extensions

Alexander Kushpel

Department of Mathematics, Çankaya University
kushpel@cankaya.edu.tr

Abstract

n -Widths of a compact origin-symmetric sets A , in a Banach space X were introduced in 1936 by Kolmogorov [2] to compare the efficiency of numerical algorithms. It was noticed by Kolmogorov and demonstrated on a concrete example by Tikhomirov [3] that the linear n -width may decrease in an isometric extension of X . Hence it is natural to consider absolute n -widths introduced by Ismagilov [1] in 1974. It was shown by Ismagilov [1] that absolute linear n -widths can be attained in the space of bounded functions on the unit ball in the dual space X^* . We show that n -dimensional extensions of X are sufficient to attain absolute linear n -widths and construct these extensions explicitly.

Keywords: Linear approximation, n -width, isometric extension.

References:

1. R. S. Ismagilov, Diameters of sets in normed linear spaces and the approximation of functions by trigonometric polynomials. Russ. Math. Surv. 29 (1974) 169.
2. A. N. Kolmogorov, Über die beste annäherung von funktionen einer gegebenen funktionklasse. Annals of Mathematics 37 (1936) 183–107.
3. V. M. Tikhomirov, Diameters of sets in functional spaces and the theory of best approximations. Russian Math. Surveys. 15 (1960) 75-111.

ICOMAA 2023

A New Method About Finding Square Root of Perfect Square Numbers

Ali Değirmenci

*Department of Industrial Engineering, Yildiz Technical University,
ali.degirmenci@yildiz.edu.tr*

Abstract

There is a relationship between the squares of two numbers whose sum or difference is multiple of fifty. There is a certain number between zero and twenty-five that will round each number to the nearest multiple of fifty. Therefore, every integer number has a related number between zero and twenty-five. In this method, a smaller perfect square number, which is square of the related number between zero and twenty-five, are used to calculate square root of a bigger perfect square number.

Keywords: Square Root, Square, Perfect Square

References:

1. Gazali, W. (2017). An algorithm to find square root of quadratic residues over finite fields using primitive elements. *Procedia computer science*, 116, 198-205.
2. Zalaket, J., & Hajj-Boutros, J. (2011). Prime factorization using square root approximation. *Computers & Mathematics with Applications*, 61(9), 2463-2467.
3. Zhu, Y. (2016). Different cultures of computation in seventh century China from the viewpoint of square root extraction. *Historia Mathematica*, 43(1), 3-25.

ICOMAA 2023

On covered quasi hyperideals in Semihypergroups

Shahnawaz

Department of Mathematics *Jamia Millia Islamia*, New Delhi-110025

Email: shahnawazali57@gmail.com

Abstract: In this article, covered quasi hyperideal (CQ-hyperideal) are developed in semihypergroups. Covered quasi hyperideal is the generalization of covered quasi ideal. The greatest covered quasi hyperideal and quasi hyperbase of a semihypergroup are described. Various properties of covered quasi hyperideals are investigated. Also, we have established a relationship between covered hyperideal with covered quasi hyperideal and greatest hyperideal with covered quasi hyperideal.

Keywords: Hyperideal; Covered quasihyperideal; Greatest hyperideal; Quasi hyperbase.

2010 AMS Subject Classification: 18B40; 20M12.

References:

1. I. Fabrici, One sided bases of semigroup, *Matematicky Cusopis*, 22 (1972), 284–290.
2. I. Fabrici, On bases and maximal ideal in semigroup, *Mathematica Slovaca*, 31 (1984), 115–120.
3. I. Fabrici, Semigroups containing covered one-sided ideals, *Mathematica Slovaca*, 31 (1981), 225–231.
4. I. Fabrici, Semigroups containing covered two-sided ideals, *Mathematica Slovaca*, 34 (1984), 355–363
5. M. Firoj Ali and Noor Mohammad Khan, Characterization of Ordered Semihypergroup by Covered Hyperideals, *Kragujevac Journal of Mathematics*, 47 (2023), 417–429.

ICOMAA 2023

Exponential Graph Resolvent

Ali A. Shukur

^{1,2,3}Department of Mathematics, University of Kufa,
Shukur.math@gmail.com


Abstract

Let G be a simple graph of order n and $A(G)$ be an adjacency matrix of G , the spectra of G is the set of eigenvalues of $A(G)$. In other side, the resolvent matrix is a matrix with property that all of its eigenvalues are outside the spectra. In this talk, we present an exponential growth to the graph resolvent.

Keywords:Spectrum, resolvent, power matrix.

References:

1. A. Shukur, Exponential Growth of Graph Resolvent, Iranian J. Math. Chem. 12 (3) December (2021) 187 – 195.
2. A. Shukur & H. Shelash, Pseudospectrum energy of graph, Iranian J. Math. Chem. 11 (2) July (2020) 83 – 93.
3. A. Shukur & I. Gutman, Energy of monad graphs, Bull. Int. Math. Virtual Inst., Vol. 11(2)(2021), 261-268.



ICOMAA 2023

On basicity a system of eigenfunctions of a second order discontinuous differential operator in Lebesgue spaces

Alirza Akhmedov¹ and Ilkin Feyzullayev²

¹Baku State University,
ehmedov-elirza@mail.ru

²Institute of Mathematics and Mechanics
ilkin0704@mail.ru

Abstract

Consider the following spectral problem with a discontinuity point

$$\begin{aligned} y''(x) + \lambda y(x) &= 0, \quad x \in (0, 1/3) \cup (1/3, 1), \quad (1) \\ y'(0) = y'(1) &= 0, \quad (2) \end{aligned}$$

$$\begin{cases} y(1/3 - 0) = y(1/3 + 0), \\ y'(1/3 - 0) - y'(1/3 + 0) = \lambda m y(1/3 - 0), \end{cases} \quad (3)$$

where λ is a spectral parameter, and m is an arbitrary complex number other than zero. Such spectral problems arise in the study of various problems of mathematical physics by the Fourier method [1]. In this paper, using the method of works [2,3], we prove theorems on the basis property of the eigenfunctions of problem (1)-(3) in the spaces $L_p(0,1) \oplus C$ and $L_p(0,1)$, $1 < p < \infty$. Note that similar questions for the spectral problem with other boundary conditions were studied in [4,5].

Keywords: spectral problem with a discontinuity point, eigenfunctions, basicity.

References:

1. Atkinson, F.V., Discrete and Continuous Boundary Problems, New York–London: Academic Press, 1964.
2. B.T.Bilalov, T.B.Gasymov, On bases for direct decomposition, Doklady Mathematics, v. 93, No 2 (2016), 183-185.
3. B.T.Bilalov, T.B.Gasymov, On basicity a system of eigenfunctions of second order discontinuous differential operator, Ufa Mathematical Journal, v. 9, No 1 (2017), 109-122.
4. T.B.Gasymov, G.V.Maharramova, N.G.Mammadova, Spectral properties of a problem of vibrations of a loaded string in Lebesgue spaces, Trans. of NAS of Azerb., Ser. Phys.-Tech. Math. Sci., 38(1) (2018), 62-68.
5. B.T.Bilalov, T.B.Gasymov, G.V.Maharramova, Basis property of eigenfunctions in Lebesgue spaces for a spectral problem with a point of discontinuity. Differential Equations, v.55, No.12 (2019), 1544-1553.

Existence of solutions for impulsive fractional boundary value problem with p-Laplacian via critical point theory

Amina Boucenna¹, Amira Rouaghi²

¹Laboratory of fixed point theory and applications, Departements of mathematics, ENS-Kouba, Algiers, Algeria

²Departements of mathematics, University of Blida 1, Algeria

amina.boucenna@g.ens-kouba.dz

rouaghiamira4@gmail.com

Abstract

In this work, we consider the boundary value problem for impulsive fractional differential equations with p-Laplacian type. Applying a minimization principal method to prove the existence of solution for this problem.

Keywords: Impulsive fractional differential equation, Boundary value problem, critical point, p-Laplace operator, minimization principal.

References:

1. Boucenna, T. Moussaoui. (2015), Existence Results for a fractional boundary value problem via critical point theory, Acta univ. Palacki. Olomuc. Fac. Rerum. Natur. Math, (54) (1), 47-64.
2. Boucenna, T. Moussaoui, Existence of a positive solution for a boundary value problem via a topological-variational theorem, J. Fract. Calc. Appl, (5) (2014), 1-9.
3. César E. Torres Ledesma, Nemat Nymoradi, Impulsive fractional boundary value problem with p-Laplace operator, Korean Society for Computational and Applied Mathematics 2016.
4. E.W.C. Van Groesen. Variational methods for nonlinear operator equations, Mathematisch Centrum. Amsterdam, 1979.

ICOMAA 2023

Existence and uniqueness of solutions for p-Laplacian fractional boundary value problem on the half-line

Amira Rouaghi¹, Amina Boucenna²

¹Departements of mathematics, University of Blida 1, Algeria

²Laboratory of fixed point theory and applications, Departements of mathematics, ENS-Kouba, Algiers, Algeria

rouaghiamira4@gmail.com

amina.boucenna@g.ens-kouba.dz

Abstract

The goal of this work is to study the existence and uniqueness of the solutions to a boundary value problem with p-Laplacian type, associated of caputo fractional differential equation. By using the fixed point theorem of Banach and Krasnosel'skii's theorem that guarenties the solutions of our problem.

Keywords: Fractional differential equation, unbounded intervals, existence solutions, fixed point theory, fractional derivative, p-Laplacian, cone.

References:

1. F. Haddouchi, Positive solutions of p-Laplacian fractional differential equations with fractional derivative boundary condition, Arxiv preprint arXiv: 1908.03966, 2019-arXiv.org.
2. A. Guezane-Lakoud, A.Kılıçman, Unbounded soluyyion for a fractional boundary value problem, Advances in Difference Equations 2014, 2014:154.
3. A.A. Kilbas, H.M. Srivastava and J.J. Trujillo, Theory and applications of fractional differential equations, Elsevier science, Amesterdam, 2006.
4. I. Podlubny, Fracional differential equations, Academic press, San Diego, 1999.

ICOMAA 2023

Hardy-Littlewood maximal function on Locally compact Abelian groups

Ani Ozbetelashvili¹

(Joint work with Tengiz Kopaliani²)

^{1,2}Department of Mathematics, Tbilisi State University,

anaozbetelashvili5677@gmail.com

tengiz.kopaliani@tsu.ge

Abstract

In this paper we investigate a boundedness of the Hardy-Littlewood maximal operator M in the variable Lebesgue spaces in the context locally compact abelian group. We show that the local Muckenhoupt condition implies the local boundedness of M .

Keywords: Locally compact abelian groups, maximal function, variable L^p spaces, Muckenhoupt weights.

References:

1. D.Cruz-Uribe and A. Fiorenza, Variable Lebesgue Spaces, Foundations and Harmonic Analysis. Birkhäuser, Basel (2013).
2. L. Diening, P. Harjulehto, P. Hästö, M. Růžička, Lebesgue and Sobolev spaces with variable exponents. Lecture Notes in Mathematics, 2017. Springer, Heidelberg (2011).
3. J. Duoandikoetxea, F.J. Martín-Reyes and S. Ombrosi, On the A_∞ conditions for general bases, Math. Z. 282 (2016), no. 3-4, 955–972.

There research was supported by Shota Rustaveli National Science Foundation of Georgia (SRNSFG), grant no: FR-22 17770.

ICOMAA 2023

Novel Tempered Fractional Hermite-Hadamard Type Inequalities via n -polynomial Uniformly Convex Functions with Application

Artion Kashuri¹, Rozana Liko²

^{1,2}Department of Mathematics, Faculty of Technical and Natural Sciences, University Ismail Qemali, 9400 Vlora, Albania,

artionkashuri@gmail.com

rozanaliko86@gmail.com

Abstract

In this paper, we obtain some novel tempered fractional Hermite-Hadamard type inequalities for n -polynomial uniformly convex functions. Moreover, using a new identity as an auxiliary result, we deduce several inequalities for n -polynomial convex functions with modulus ψ via tempered fractional integrals, and some special cases are given as well. To validate the accuracy of our main results, we present an application to special means.

Keywords: Hermite-Hadamard type inequalities, n -polynomial uniformly convex functions, Tempered fractional integral operators, Hölder's inequality, Power-mean inequality, Special means.

References:

1. Barsam, H., Sattarzadeh, A.R. Hermite-Hadamard inequalities for uniformly convex functions and its applications in means. *Miskolc Math. Notes* 21(2) 2020 621-630.
2. Kashuri, A., Liko, R. Some new Hermite-Hadamard type inequalities and their applications. *Stud. Sci. Math. Hung.* 56(1) 2019 103-142.
3. Niezgoda, M. An extension of Levin-Stečkin's theorem to uniformly convex and superquadratic functions. *Aequationes Math.* 94(4) 2020 303-321.
4. Barsam, H., Sayyari, Y. On some inequalities of differentiable uniformly convex mapping with applications. *Numer. Funct. Anal. Optim.* 44(4) 2023.
5. Zălinescu, C. On uniformly convex functions. *J. Math. Anal. Appl.* 95(2) 1983 344-374.

ICOMAA 2023

APPROACHING GEOMETRIC PROBLEMS WITH THE PARTICLE SWARM OPTIMIZATION METHOD

KARAÇAM CEMİL¹, SEVGİ ATA², KARACAN EFE ATAKAN³

KARTAL ARDA EREN⁴

¹ Mathematics Department, Köyceğiz Science and Art Center, Muğla/Turkey.

^{2,3,4} Galatasaray High School, İstanbul/Turkey.

Abstract

In our project, which we initiated due to the lack of application in the literature, it is aimed to develop different methods by approaching 7 different analytical geometry problems with the particle swarm optimization method. These problems include a problem which aims to divide the area between the x and y lines of a parabolic equation into 4 areas that will be proportional to a 4-element number array, and to find the points where two lines perpendicular to each other intersect the x-axis and the slope of one line. Another problem is to find the points where two lines intersect each other at right angles, which divide a triangle whose one point is on the origin into 4 equal areas, intersect the x-axis at two different points, and find the slope of one line. The method used in our project has been transformed into a design that can be used in many complex systems and problems that involve many variables.

Keywords: Analytical geometry, Particle Swarm Optimization, Area problems, Optimization by coding

References:

1. ADITYA, J. , ADITYA K. , A New Method of Solving the Bernoulli Quadrisection Problem, Synge Street CBS, 2022.
2. Eberhart, C. 2018. Revisiting the Quadrisection Problem of Jacob Bernoulli. Forum Geometricorum, Volume 18
3. ÖZSAĞLAM, M. Y. , ÇUNKAŞ M. , Optimizasyon Problemlerinin Çözümü için Parçacık Sürü Optimizasyonu Algoritması, Politeknik Dergisi, Cilt:11 Sayı: 4 s.299-305, 2008
4. Kwon Y.D., Kwon S.B., Kim J., “Convergence enhanced genetic algorithm with successive zooming method for solving continuous optimization problems”, Comput. Struct. 81:1715–1725, 2003.
5. Wilke D. N., Analysis of the particle swarm optimization algorithm, Master Thesis, 2005, Mechanical and Aeronautical Engineering, University of Pretoria.

ICOMAA 2023

On frame properties of iterates of a multiplication operator

A.Sh.Shukurov

Institute of Mathematics and Mechanics, Ministry of Science and Education, Republic of Azerbaijan; F.Agayev 9, Baku, Az1141, Azerbaijan,
ashshukurov@gmail.com

Abstract

This note is dedicated to the study of frame properties of iterates of a multiplication operator $T_\varphi f(t) = \varphi(t) \cdot f(t)$, $f \in L_2(a, b)$.

Theorem 1. Let $\varphi(t)$ be any measurable function and $f(t)$ any square summable function on (a, b) . The system $\{T_\varphi^n f\}_{n=0}^\infty$ cannot be a frame in $L_2(a, b)$.

Theorem 2. If $\{\varphi^n(t)\}_{n=-\infty}^\infty$ is a frame in $L_2(a, b)$, then $|\varphi(t)| = 1$ a.e. on (a, b) , i.e. $\varphi(t) = e^{i\alpha(t)}$, where $\alpha(t)$ is a real-valued function.

Theorem 3. Suppose that $\alpha(t)$ is an invertible function, inverse $\xi: [p, q] \rightarrow [a, b]$ of which satisfies the following conditions:

- 1) $\xi(t)$ is absolutely continuous, strictly increasing function on $[p, q]$; $\xi(p) = a$ and $\xi(q) = b$;
- 2) $[p, q] \subset [0, 2\pi]$ and there are constants $A, B > 0$ such that $A \leq \xi'(t) \leq B$ for almost all $t \in [p, q]$.

Then the system $\{e^{in\alpha(t)}\}_{n=-\infty}^\infty$ is a frame in $L_2(a, b)$.

Acknowledgement. The author is grateful to Professor B.T. Bilalov for discussions.

Keywords: Dynamical sampling, operator orbit, frame, system of powers, Lebesgue spaces.

References:

1. Shukurov, A. Sh., Kasumov Z.A. On frame properties of iterates of a multiplication operator, Results Math. 74 (2019), no. 2, 74:84.
2. Shukurov, A. Sh. Necessary condition for Kostyuchenko type systems to be a basis in Lebesgue spaces. Colloq. Math. 127 (2012), no. 1, 105-109.
3. Shukurov, A. Sh. Addendum to "Necessary condition for Kostyuchenko type systems to be a basis in Lebesgue spaces". Colloq. Math. 137 (2014), no. 2, 297-298.
4. Shukurov, A. Sh. The power system is never a basis in the space of continuous functions. Amer. Math. Monthly 122 (2015), no. 2, p.137.
5. Shukurov, A. Sh. Impossibility of power series expansion for continuous functions. Azerb. J. Math. 6 (2016), no. 1, 122-125.

Levinson-Type Formula for Sturm-Liouville Operator with a Rational Function of Spectral Parameter in Boundary Condition

Aynur Çöl¹

¹Department of Mathematics and Science Education, Sinop University
aynurcol@sinop.edu.tr

Abstract

We consider the Sturm-Liouville operator on the half line $[0, \infty)$ with a rational function of spectral parameter in the boundary condition and investigate the continuity of scattering function $S(\lambda)$. The increment of the argument of scattering function is obtained.

Keywords: Sturm-Liouville operator, scattering function, Levinson-type formula.

References:

1. Marchenko V.A. Sturm-Liouville Operators and Applications, Birkhauser Verlag, Basel, 1986.
2. Çöl, A. (2022). On a scattering problem for Sturm-Liouville equation with a rational function of spectral parameter in boundary condition. Journal of Contemporary Applied Mathematics, 12(2), 73-82.
3. Çöl, A. (2023). Inverse scattering problem for Sturm-Liouville equation with a rational function of spectral parameter in boundary condition. Journal of Contemporary Applied Mathematics, 13(1), 3-10.

ICOMAA 2023

On a Scattering Problem for Discontinuous Second Order Differential Operators with Herglotz Function of Spectral Parameter in Boundary Condition

Aynur Çöl¹ and Khanlar R. Mamedov²

¹Department of Mathematics and Science Education, Sinop University
aynurcol@sinop.edu.tr

²Department of Mathematics, Iğdır University
hanlar.residoglu@igdir.edu.tr

Abstract

In this work, we consider a boundary value problem of scattering theory for second order differential operators on the half line ($0 \leq x < \infty$) with discontinuous coefficient and Herglotz function of the spectral parameter in the boundary condition. By using the new representation of solution of differential equation, scattering data is provided and its properties are investigated. The Marchenko-type fundamental equation is derived in order to discuss inverse scattering problem.

Keywords: Second order differential operator, boundary value problem, scattering data, spectral parameter, discontinuous coefficient.

References:

1. Marchenko V.A. Sturm-Liouville Operators and Applications, Birkhauser Verlag, Basel, 1986.
2. Mamedov Kh.R. (2010) On an inverse scattering problem for a discontinuous Sturm-Liouville equation with a spectral parameter in the boundary condition. Boundary Value Problems, 2010:171967.
3. Çöl, A. (2015). Inverse spectral problem for Sturm-Liouville operator with discontinuous coefficient and cubic polynomials of spectral parameter in boundary condition. Advances in Difference Equations, 2015:132.

ICOMAA 2023

Basicity of the double exponential system in weighted Lebesgue space

Aysel Guliyeva

Ganja State University, Ganja, Azerbaijan,
aysel_guliyeva20@mail.ru

Abstract

This work considers a double exponential system with complex-valued coefficients, which is a generalization of the well-known exponential system $\{e^{i(n-\alpha \operatorname{sign} n)t}\}_{n \in \mathbb{Z}}$, where α – is, in general, some complex parameter. The study of basis properties of this system has a deep history which dates back to the works of Paley, Wiener and N. Levinson. The well-known Kadets 1/4 theorem also refers to this range of questions. In this work, it is proved that the double exponential system forms a basis for the weighted Lebesgue space $L_{p,w}(-\pi, \pi)$, $1 < p < +\infty$, if and only if it is isomorphic to the classical exponential system $\{e^{int}\}_{n \in \mathbb{Z}}$ in it and the weighted function $w(\cdot)$ satisfies the Muckenhoupt condition.

Keywords: weighted Lebesgue spaces, exponential system, basis property, Muckenhoupt condition.

References:

1. A.E.Guliyeva, Isomorphism of Two Bases of Exponentials in Weighted Lebesgue Spaces. Azerbaijan Journal of Mathematics V. 12, No 1, 2022, 183-194
2. B.T. Bilalov, T.R. Muradov, On equivalent bases in Banach spaces, Proc. ins. Math. and Mech., NAS Az., XXIII (XXXI), 2006, 3-8
3. S.R. Sadigova, A.E. Guliyeva, Bases of the perturbed system of exponents in weighted Lebesgue space with a general weight, Kragujevac Jour. Math., 46(3), 2022, 477-486

ICOMAA 2023

Conformal Mapping Preserving the Z-Symmetric Tensor on a Riemannian Manifold

Ayşe Yavuz Taşcı¹, Füsun Özen Zengin²

¹ Piri Reis University,

² Department of Mathematics, Istanbul Technical University,

aytasci@pirireis.edu.tr

fozen@itu.edu.tr


Abstract

In this paper, we denote the conformal mapping of the Z-symmetric tensor on a Riemannian manifold. After that, the properties of the conformal mapping that preserves this tensor are examined.

Keywords: Conformal mapping, Z-symmetric tensor, conharmonic transformation.

References:

1. H.W. Brinkmann, Einstein spaces which are mapped conformally on each other, Math. Ann., 94(1), (1925), 119-145.
2. O. Chepurna, V. Kiosak and J. Mikes, Conformal mappings of Riemannian spaces which preserve the Einstein tensor, Aplimat-Journal of Applied Mathematics, 3(1), (2010), 253-258.
3. L.P. Eisenhart, Riemannian geometry, Princeton Univ. Press, 1926.



ICOMAA 2023

Existence and decay of solutions for a fourth-order hyperbolic equation with a variable coefficients

Ayşe Fidan¹, Erhan Pişkin²

¹Department of Mathematics, Dicle University

²Institute of Natural and Applied Sciences, Dicle University

afidanmat@gmail.com

episkin@dicle.edu.tr

Abstract

In this presentation, we consider a fourth-order hyperbolic equation with a variable coefficients. Problems about the mathematical behavior of solutions for PDEs with time delay effects have become interesting for many authors mainly because time delays often appear in many practical problems such as chemical, physical, thermal, biological, economic phenomena, electrical engineering systems, mechanical applications and medicine. We prove the existence and decay of solutions.

Keywords: Existence, decay, variable coefficients.

References:

1. Benaissa, A. Benguessoum and S. A. Messaoudi, Energy decay of solutions for wave equation with a constant weak delay and a weak internal feedback, *Electronic Journal of Qualitative Theory of Differential Equations*, 11(2014) 1-13.
2. E. Pişkin, A. Fidan, Nonexistence of global solutions for the strongly damped wave equation with variable coefficients, *Universal Journal of Mathematics and Applications*, 5 (2) (2022) 51-56.
3. X. Zheng, Y. Shang, X. Peng, Blow up of Solutions for a Nonlinear Petrovsky Type Equation with Time-dependent Coefficients, *Acta Mathematicae Applicatae Sinica, English Series*, 36(4) (2020) 836-846.

ICOMAA 2023

The Inverse Sturm-Liouville Problem With One Of The Boundary Conditions Including Eigenvalue

Seda Kızılbudak Çalışkan, Ayşegül Kılıç
Department of Mathematics, Yıldız Technical University
skizilb@yildiz.edu.tr
aysegullklc@gmail.com

Abstract

Spectral analysis of differential operators has an important place in applied sciences and especially in some physics problems. Spectral analysis is studied under two headings: straight problems and inverse problems.

In our study, we considered the Inverse Sturm-Liouville Problem.

Based on the spectral properties of the Sturm-Liouville Problem with an eigenvalue in one of the boundary conditions we have considered, we have obtained the necessary and sufficient conditions for the potential function “q” in our equation and its solvability by using the eigenvalue sequences and norm constants of all the parameters in our boundary conditions.

Keywords: Inverse Problem, Sturm-Liouville Equation.

References:

1. Naming J Guliyev 2005 Inverse eigenvalue problems for Sturm-Liouville equations with spectralparameter lenearly contained in one of the boundary conditions
2. Levitan B M and Gasymov M G 1964 Determination of a differential equation by two of its spectra (Russian) Uspekhi Mat. Nauk 19 3-63
3. Marchenko V A 1977 The Sturm-Liouville operators and their applications (Russian) (Kiev: Naukova Dumka) English transl.: (Basel:Birkhäuser Verlag, 1986)
4. Yurko V A 2001 Inverse spectral problems and their applications (Russian) (Saratov: Izd-vo Saratovskogo pedinstituta) See also: Freiling G and Yurko V A 2001 Inverse Sturm-Liouville problems and their applications (Huntington, NY: Nova Science Publishers)
5. Yurko V A 2002 Method of spectral mappings in the inverse problem theory (Utrecht: VSP)

ICOMAA 2023

FINITE ELEMENT BASED SCHEME FOR THE BURGERS EQUATION WITH FORCING EFFECTS

Selmahan Selim¹, Aysenur Busra Cakay²
^{1,2}Department of Mathematics, Yildiz Technical University,
sselim@yildiz.edu.tr
bsr.cakay@hotmail.com

Abstract

This study concentrates on investigation of numerical solutions of the Burgers Equation with source effects. To achieve that aim, the article offers three-point discretization in time and Galerkin Finite Element Method (TDFEM3). In the present method the backward finite difference in time and the finite element method in space are applied to solve the Burgers equation, and then the resulting system of the nonlinear ordinary differential equations obtained at each time step is solved by using computer codes generated in MATLAB. To show the efficiency of the presented method, the numerical solutions evaluated for various values of viscosity at different times is stated in terms of the error norms. These methods are seen to be a very good choice to obtain a high degree of accuracy for the numerical solution of the Burgers equation.

Keywords: Burgers equation, forcing term, time discretization, the backward finite difference, Galerkin finite element method

References:

1. Aksan, E. N., A Numerical Solution of Burgers' Equation by Finite Element Method Constructed on the Method of Discretization in Time, Appl. Math. Comput, 170 (2005), 2, 895-904.
2. H. P. Langtangen. Introduction to Computing with Finite Difference Methods, Simula Research Laboratory and University of Oslo, 2013.
3. J.N.Reddy, Applied Functional Analysis and Variational Method in Engineering, McGraw-Hill Book Com. New York (1986).

ICOMAA 2023

On non-increasing rearrangement of generalized fractional maximal function

Azhar Abek

L.N. Gumilyov Eurasian National University, Astana, Kazakhstan,
azhar.abekova@gmail.com

Abstract

We give a sharp pointwise estimates of the non-increasing rearrangement of the generalized fractional maximal function via an expression involving the non-increasing rearrangement of f . It is shown that the obtained estimate is more sharp than the inequality which is followed from the estimate for the generalized Riesz potential.

Similar questions for the space of classical fractional maximal function are considered in [1] and [2]. And questions of non-increasing rearrangements of generalized Riesz potentials were previously considered in [3].

Keywords: generalized fractional maximal function, non-increasing rearrangements, generalized Riesz potential

References:

1. N.A.Bokayev, M.L.Goldman, G.Zh.Karshygina, Cones of functions with monotonicity conditions for generalized Bessel and Riesz potentials. *Mathem. Notes.* 104 (3) (2018), 356-373.
2. A.Cianchi, R. Kerman, B. Opic, L. Pick, A sharp rearrangement inequality for the fractional maximal operator. *Studia Mathematica.* 138 (3) (2000), 277-284.
3. M.L.Goldman, On the cones of rearrangements for generalized Bessel and Riesz potentials. *Complex Variables and Elliptic Equations.* 55 (8-10) (2010), 817-832.

ICOMAA 2023

On the strong solvability of a nonlocal boundary value problem for the Laplace equation in a rectangular domain

Telman Gasymov¹ and Beherchin Akhmedli²

¹Department of Mathematics and Mechanics, Baku State University,
telmankasumov@rambler.ru

²Institute of Mathematics and Mechanics, Baku, Azerbaijan
a_beberchin@mail.ru

Abstract

The following nonlocal problem for the Laplace equation in a rectangular domain is considered:

$$u_{xx} + u_{yy} = 0, \quad 0 < x < 2\pi, \quad 0 < y < h, \quad (1)$$

$$u(x, 0) = \varphi(x), \quad u(x, h) = \psi(x), \quad 0 < x < 2\pi, \quad (2)$$

$$u_x(0, y) = 0, \quad u(0, y) = u(1, y), \quad 0 < y < h. \quad (3)$$

Such problems have specific features in comparison with problems with local conditions. For the Laplace equation in an unbounded domain, a similar problem was considered in [1,2], where the classical solution of the problem is studied. Earlier in [3], problems with nonlocal boundary conditions for a shifted equation were considered. For elliptic equations, nonlocal problems were considered in [4], and in [5], a boundary value problem with a nonlocal condition for multidimensional parabolic equations was solved.

In this paper, we study problem (1)-(3) in a weighted Sobolev space with a weight from the Mackenhaupt class. The notion of a strong solution of this problem is defined. Using the Fourier method, under certain conditions on the functions $\varphi(x)$ and $\psi(x)$, we prove the correct solvability of this problem.

Keywords: Laplace equation, nonlocal problem, weighted Sobolev space, strong solution.

References:

1. E.I. Moiseev, On the solution of a nonlocal boundary value problem by the spectral method. *Differents. Uravneniya*, v. 35, No. 8 (1999), 1094-1100.
2. M.E. Lerner, O.A. Repin, On Frankl'-type problems for some elliptic equations with degeneration of various types. *Differents. Uravneniya*, v. 35, No. 8 (1999), 1087-1093.
3. Frankl F.I. *Selected works on gas dynamics*, Moscow, 1973, 711 p.
4. Bitsadze A.V., Samarsky A. A. On some simplest generalizations of linear elliptic boundary value problems // *Reports of the USSR Academy of Sciences*, v. 185, No. 4 (1969), 739-740.
5. N.I. Ionkin and E.I. Moiseev, A Problem for Heat Transfer Equation with Two-Point Boundary Conditions, *Differents. Uravneniya*, v.15, No. 7(1979), 1284-1295.

Study of Nonlinear Partial Fractional Boundary Value Problem

Noureddine Bouteraa¹

Laboratory of Fundamental and Applied Mathematics of Oran (LMFAO), University of Oran 1, Ahmed Ben Bella, Algeria.

¹Oran Graduate School of Economics, Algeria

bouteraa-27@yahoo.fr

Abstract

In this piece writing, to resolve fractional stochastic equation with initial data, the fractional Duhamel principle is applied and use it to establish the well-boundedness and stability of a mild solution of our problem.

Keywords: Partial differential equations, existence, Duhamel principle, stability.

References:

1. J. Banas and K. Goebel, Measure of Noncompactness in Banach Space, Lecture Notes in Pure and Applied Mathematics, Marcel Dekker, New York, 1980.
2. T.E. Duncan, B. Maslowski, B. Pasik-Duncan, Semilinear stochastic equations in a Hilbert space with a fractional Brownian motion, SIAM J. Math. Anal. 40(6) (2009) 2286-2315. 335-201.
3. R. Metzler, K. Joseph, The random walk's guide to anomalous diffusion: a fractional dynamics approach, Physics reports, 339, 1(2000), 1-77.
4. Y. S. Mishura, Stochastic calculus for fractional Brownian motion and related processes, Springer (2008).
5. A. Viorel, Contribution to the Study of Nonlinear Evolution Equations, Ph.D. Thesis, Babes-Bolyai Univ. Cluj-Napoca, Department of Mathematics, 2011.

ICOMAA 2023

Recursive Estimation of the Conditional Distribution Function on Riemannian Manifolds

BOUADJEMI Abdelkader¹

¹ *Department of Mathematics,
Amine Elokkel El Hadj Moussa University of Tamanerasset
Tamanerasset,
Algeria.*

bouadjemi@gmail.com

Abstract

The main of this paper is to treat the estimation of conditional distribution function on Riemannian manifolds. We defined the recursive version of the Nadaraya–Watson estimator. Under some assumptions in Riemannian Manifolds data analysis, we study the properties of a recursive family kernels regression, the asymptotic normality of estimator is established.

Keywords: Riemannian Manifold, asymptotic normality, conditional cumulative distribution function.

References:

1. Bouadjemi, A. Asymptotic normality of the recursive kernel estimate of conditional cumulative distribution function. Journal of probability and statistical sciences, (2014). 12, Pages 117-126.
2. Salah khardani ; Wiem NefziAnne ; Françoise Yao. Nonparametric Kernel Spatial Density Estimation on Riemannian manifolds. January 2022. SSRN Electronic Journal; DOI: 10.2139/ssrn.4222942
3. Salah KhardaniAnne Françoise Yao. Nonparametric recursive regression estimation on Riemannian Manifolds. Statistics & Probability Letters October 2021,182(2):109274
4. Bruno Pelletier. (2006). Non-parametric regression estimation on closed Riemannian manifolds. Journal of Nonparametric Statistics 18(1):57-67.

ICOMAA 2023

On the recursive sequence

Burak Oğul¹

¹Department of Management Information Systems, Istanbul Aydin University,
burakogul@aydin.edu.tr

Abstract

Difference equations appear naturally as discrete analogs and as numerical solutions of differential and delay differential equations, having applications in biology, ecology, physics. Difference equations are used in a variety of contexts, such as in economics to model the evolution through time of variables such as gross domestic product, the inflation rate, the exchange rate, etc.

In this paper, we explore the dynamics of adhering to rational difference formula

$$x_{n+1} = \frac{x_{n-26}}{1 + x_{n-2}x_{n-5}x_{n-8}x_{n-11}x_{n-14}x_{n-17}x_{n-20}x_{n-23}},$$

where initial values are nonnegative real numbers.

Keywords: Difference equation, equilibrium points, recursive sequence.

References:

1. V.L. Kocic, G. Ladas, Global behavior of nonlinear difference equations of higher order with applications, Mathematics and its Applications, Kluwer Academic Publishers Group, Dordrecht, (1993).
2. E.M. Elsayed, K.N. Alharbi, The expressions and behavior of solutions for nonlinear systems of rational difference equations, Journal of Innovative Applied Mathematics and Computational Sciences (JIAMCS), 2(1), 78-91 (2022).
3. R. Karatas, C. Cinar, D. Simsek, On Positive Solutions of the Difference Equation $x_n = x_{n-5}$, Int. J. Contemp. Math. Sci, 1(10), 495-500(2006).
4. D. Şimşek, A. Doğan, On a class of recursive sequence, Manas Journal of Engineering, 2(1), 16-22(2014).

ICOMAA 2023

Dynamics and Hopf Bifurcation of A Sustainable Tourism Model with Time Delay.

Prof. Dr. Mustafa Taylan Şengül¹ and Bünyamin Kurtkaya²

^{1,2}*Department of Mathematics, Marmara University,*

taylan.sengul@marmara.edu.tr

bunyaminkurtkaya@marun.edu.tr

Abstract

In this work, a model describing the relations between the tourists visiting a protected area, environmental pollution and the capital stock is proposed and studied. The model consists of a system of three nonlinear delay ordinary differential equations. The time delay in the model accounts for the spread of public praise and plays a crucial role in the dynamics.

We show that the system possesses a positive equilibrium point. By studying the stability of this equilibrium, we find that as the time delay parameter exceeds a critical threshold, a Hopf bifurcation occurs which leads to the emergence of a time-periodic solution. We also analyze the stability of the Hopf bifurcation by applying the normal form theory and the center manifold theorem.

Finally we provide several numerical simulations and describe the results in terms of its consequences in terms of sustainable tourism.

Keywords: Sustainability, Stability Analysis, Hopf Bifurcation, Equilibrium Point, Time Delay.

References:

1. Russu P. Hopf bifurcation in an environmental defensive expenditures model with time delay. *Chaos Solitons Fractals*, 2009, 42, 3147-3159.
2. Becker R. Intergenerational equity: The capital-environment trade-off. *J. Environ. Econ Manage*, 1982, 9, 165-185.
3. Casagrandi R, Rinaldi S, A theoretical approach to tourism sustainability. *Conserv Ecol* 2002, 6
4. Hassard B, Kazarinoff D, Wan Y. *Theory and Applications of Hopf Bifurcation*. Cambridge, UK: Cambridge University Press, 1981.

ICOMAA 2023

Korovkin-type theorems and their statistical versions in the weighted grand-Lebesgue spaces without Muckenhoupt condition

Yusuf ZEREN¹, Migdad Ismailov², Cemil Karacam³ and Fatih Sirin⁴

¹Department of Mathematics, Yildiz Technical University
yzeren@yildiz.edu.tr

²Institute of Mathematics and Mechanics of the NAS of Azerbaijan,
Baku State University
migdad-ismailov@rambler.ru

³Koycegiz Science and Art Center
cemil-karacam@hotmail.com

⁴Department of Mathematics, Halic University,
fatihSirin@halic.edu.tr

Abstract

In this paper, we investigate Korovkin-type theorems and their statistical versions in the weighted grand Lebesgue spaces $L_{p,\rho}$ based on shift operator. We define the subspace $G_{p,\omega}(-\pi, \pi)$ of the space $L_{p,\omega}(-\pi, \pi)$, where continuous functions are dense, and study some properties of the functions belonging to this space. The analogs of Korovkin theorems are proved. $1 < p < +\infty$, when weight function $\omega \in L_1(-\pi, \pi)$, $\exists \epsilon_0 \in (0, p-1)$ and $\omega^{-1}(t) \in L_{\frac{1}{p-\epsilon_0-1}}(-\pi, \pi)$.

Keywords: weighted grand-Lebesgue space, Korovkin theorems, statistical convergence

References:

1. R.E.Castilo,H.Rafeiro,An Introductory Course in Lebesgue Spaces, Springer International Publishing Switzerland, 2016.
2. R.A.Hunt,B.Muckenhoupt,R.L.Wheeden,Weighted norm inequalities for the conjugate function and Hilbert transform, Trans. of Amer. Math. Soc., 176(1973), Soc., 176(1973), 227-251.
3. Zeren Y. İsmailov M. Karacam Cemil. Korovkin-type theorems and their statistical versions in grand Lebesgue spaces Turk J Math(2020)44: 1027 –1041 ©TÜBİTAK doi:10.3906/mat-2003-21

ICOMAA 2023

HARMONIC TRINOMIAL WITH COMPLEX

PARAMETER

CHAHRAZED HARRAT¹

¹Department of Mathematics, Faculty of Mathematics and Computer Sciences, University of Sciences and Technology "M.B" of Oran, Laboratory of, Fundamental and Applied Mathematics of Oran (LMFAO), University Oran 1, Algeria,

harratchahrazed@gmail.com

Abstract

Much work has been done examining the similarities and differences between analytic and harmonic functions. Many familiar results for analytic functions hold for complex-valued harmonic functions with only slight modifications. The harmonic analog of the Argument Principle for Analytic Functions. In this paper, we study the number of zeros of harmonic trinomials with complex coefficients of the form $h(z) = z^n + az^k - 1$, $n > k$, and $\gcd(n; k) = 1$; we determine how the number of zeros depends on the complex parameter a .

Keywords:

Polynomial, Harmonic trinomial, Wilmschurst's conjecture .

References:

1. Michael A. Brilleslyper and Lisbeth E. Schaubroeck, Locating unimodular roots, *College Math. J.* 45 (2014), no. 3, 162--168, DOI 10.4169/college.math.j.45.3.162. MR3207562.
2. Michael A. Brilleslyper and Lisbeth E. Schaubroeck, Counting interior roots of trinomials, *Math. Mag.* 91 (2018), no. 2, 142--150, DOI 10.1080/0025570X.2017.1420332. MR3777918
3. Daoud Bshouty, Walter Hengartner, and Tiferet Suez, The exact bound on the number of zeros of harmonic polynomials, *J. Anal.Math.* 67 (1995), 207--218, DOI 10.1007/BF02787790. MR1383494.
4. Dmitry Khavinson and Grzegorz Świątek, On the number of zeros of certain harmonic polynomials, *Proc. Amer. Math. Soc.* 131 (2003), no. 2, 409--414, DOI 10.1090/S0002-9939-02-06476-6. MR1933331.
5. Marie M. Johnson, Book Review: *Analytic Geometry* by Charles H. Lehmann, *Natl. Math. Mag.* 17 (1943), no. 6, 280. MR1570119.

ICOMAA 2023

Numerical Solution of Second-order Partial Volterra Integro Differential Equations Via Taylor Collocation Method

Cheima Khennaoui¹

¹Department of Mathematics and Informatiques, University center AbdelHafid boussouf Mila, Algeria.
c.khennaoui@centre-univ-mila.dz

Abstract

The goal of this work is to propose a method to solve a particular type of partial Volterra integro-differential equations (PVIDEs) numerically. These equations describe the behavior of systems that involve both waves and particles, making them particularly useful in studying phenomena like fluid dynamics, acoustics, and electromagnetic radiation.

The PVIDEs under consideration in this work are second kind, two-dimensional linear partial Volterra integro-differential equations of second-order

$$\frac{\partial^2 u(x, y)}{\partial x \partial y} + \alpha \frac{\partial u(x, y)}{\partial x} + \beta \frac{\partial u(x, y)}{\partial y} = \gamma u(x, y) + g(x, y) + \int_0^x H(x, y, t) u(t, y) dt,$$

subject to initial value conditions.

The proposed technique relies on utilizing two-dimensional Taylor polynomials and explicit schemes to obtain an approximate solution. The accuracy and efficiency of the method are illustrated through various examples using error estimation and convergence testing.

Keywords: Partial Volterra integro-differential equation, Hyperbolic partial differential equation, Collocation method, Two-dimensional equations, Taylor polynomials.

References:

1. A. Rivaz, S. J. Ara and F. Yousefi. Two-dimensional Chebyshev polynomials for solving two-dimensional integro-differential equations, Cankaya University Journal of Science and Engineering 12 (2); 2015.
2. H. J. Dobner. Bounds for the solution of hyperbolic problems, Computing (Wien. Print). 38(3), 209–218; 1987.
3. E. Yalçın and M. Sezer. A compatible Hermite–Taylor matrix-collocation technique with convergence test for second-order partial integro-differential equations containing two independent variables with functional bounds. Mathematical Sciences, 16(1), 59-69; 2022.

Existence of Real Models of K3-Surfaces in Arithmetical Terms

Çisem Güneş Aktaş

Department of Engineering Sciences, Abdullah Gül University,
cisem.gunesaktas@agu.edu.tr


Abstract

Although it is quite common that a real variety may have no real points, very few examples of equisingular deformation classes of projective models of K3-surfaces with this property are known. Clearly, any class of equisingular strata containing a real model is real. However, the converse is not true, but the only known counterexample is the stratum of the space of sextics found in [1]. In this research, we study phenomena of this kind, considering from a more general perspective, for all projective models of K3-surfaces and restate the existence of real models in arithmetical terms.

Keywords: K3-surfaces, real models, arithmetical reduction, projective models.

References:

1. Ayşegül Akyol and Alex Degtyarev, Geography of irreducible plane sextics, Proc. Lond. Math. Soc. (3) 111(2015) 1037–1337.
2. Alex Degtyarev and Ilia Itenberg, On real determinantal quartics, Proceedings of the Gökova Geometry-Topology Conference 2010, Int. Press, Somerville, MA (2011), 110–128.
3. Çisem Güneş Aktaş, Classification of simple quartics up to equisingular deformation, Hiroshima Math J. 47 (2017) 87–112.



ICOMAA 2023

Maximal-simultaneous approximation by Faber series in Bergman space

Daniyal M. Israfilov

Department of Mathematics, Balikesir University,
mdaniyal@balikesir.edu.tr

Abstract

Let G be a bounded simple connected domain in the complex plane C , bounded with a quasiconformal boundary L . We recall that L is a quasiconformal curve if there exists a quasiconformal homeomorphism of the complex plane C onto itself that maps a circle onto L . For the functions f analytic in G we consider the Bergman space defined as: $A^2(G) := \{f: \iint_G |f(z)|^2 d\sigma_z\}$.

We study maximal-simultaneous approximation of the Faber series produced by the integral representation given by V. I. Belyi in the domains with a quasiconformal boundary. The error of this approximation in the norm of the space $A^2(G)$ with dependence of the best approximation number and also of the parameters of the considered canonical domains are estimated.

Keywords: Faber series, maximal convergence, simultaneous approximation, Bergman space.

References:

1. Belyi V.I. Conformal mappings and approximation of functions in the domains with the quasiconformal boundary. Mat. Sb. 1977; 102 (3): 331-361 (Russian); English trans. in Math. USSR-Sb. 1977; 31: 289-317.
2. Israfilov D.M., Gürsel E., Aydin E. (2018), Maximal convergence of Faber Series in Smirnov Classes with Variable Exponent, Bull Braz Math Soc, New Series 2018; 49: 955–963.
3. Israfilov D.M., Testici A. (2017). Approximation problems in the Lebesgue spaces with variable exponent. Journal of Mathematical Analysis and Applications. 2017; 459(1), 112-123.

ICOMAA 2023

The application of expansive mappings to the existence problems

Dariusz Wardowski

*Faculty of Mathematics and Computer Science, Department of Nonlinear Analysis,
University of Lodz*

dariusz.wardowski@wmii.uni.lodz.pl

Abstract

We consider a certain class of expansive mappings defined on a closed and bounded subset of a Banach space. For such mappings there will be presented some results of Krasnosel'skii type concerning the existence of fixed points for compact operators. The results will be applied to the existence problems of nonlinear type.

Keywords: Expansive mapping, Banach space, Compact operator, Fixed point, Nonlinear integral equation.

References:

1. Krasnosel'skii, M.A.: Some problems of nonlinear analysis. Am. Math. Soc. Transl. Ser. 2(10), 345–409 (1958).
2. Liu, Y., Li, Z.: Krasnoselskii type fixed point theorems and applications. Proc. Am. Math. Soc. 136, 1213–1220 (2008).
3. N.-A. Secelean, D. Wardowski, Expansive mappings on bounded sets and their application to rational integral equations. RACSAM 114:134 (2020).
4. Wang, F., Wang, F.: Krasnosel'skii type fixed point theorem for nonlinear expansion. Fixed Point Theory 13(1), 285–291 (2012).
5. Xiang, T., Yuan, R.: A class of expansive-type Krasnosel'skii fixed point theorems. Nonlinear Anal. 71, 3229–3239 (2009).

ICOMAA 2023

Boundedness and Compactness of Commutators for Riesz Potential on Local Morrey-type spaces.

Dauren Matin¹, Talgat Akhazhanov¹, Aidos Adilkhanov¹

¹L. N. Gumilyov Eurasian National University, Astana, Kazakhstan .,

d.matin@mail.ru

talgat-a2008@mail.ru

adilkhanov-kz@mail.ru

Abstract

The paper considers Morrey-type local spaces from $LM_{p\theta}^w$. The main work is the proof of the commutator compactness theorem for the Riesz potential $[b; I_\alpha]$ in local Morrey-type spaces from $LM_{p\theta}^{w_1}$ to $LM_{q\theta}^{w_2}$.

We also give new sufficient conditions for the commutator to be bounded for the Riesz potential $[b; I_\alpha]$ in local Morrey-type spaces from $LM_{p\theta}^{w_1}$ to $LM_{q\theta}^{w_2}$. In the proof of the commutator compactness theorem for the Riesz potential, we essentially use the boundedness condition for the commutator for the Riesz potential $[b; I_\alpha]$ in local Morrey-type spaces $LM_{p\theta}^w$, and use the sufficient conditions from the theorem of precompactness of sets in local spaces of Morrey type $LM_{p\theta}^w$ [1], [2]. In the course of proving the commutator compactness theorem for the Riesz potential, we prove lemmas for the commutator ball for the Riesz potential $[b; I_\alpha]$. Similar results were obtained for global Morrey-type spaces $GM_{p\theta}^w$ [3] and for generalized Morrey spaces M_p^w [4].

Keywords: Compactness, Commutators, Riesz Potential, Local Morrey-type spaces.

This work is funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan (grant no. AP14969523).

References:

1. Bokayev, N.A., Burenkov, V.I., Matin, D.T. (2017). On precompactness of a set in general local and global Morrey-type spaces. *Eurasian mathematical journal*, 8 (№3), 109 - 115.
2. Bokayev, N.A., Burenkov, V.I., Matin, D.T. (2018). Sufficient conditions for the precompactness of sets in Local Morrey-type spaces. *Bulletin of the KARAGANDA UNIVERSITY*, 92 (№4), 54-63.
3. Bokayev, N.A., Matin, D.T. & Baituyakova, Zh.Zh. (2018). A sufficient condition for compactness of the commutators of Riesz potential on global Morrey-type space. *AIP Conference Proceedings*, (№1997), 020008. doi: 10.1063/1.5049002.
4. Bokayev, N.A., Matin, D.T. (2017). A sufficient condition for compactness of the commutators of Riesz potential on global Morrey-type space. *Herald L.N. Gumilyov Eurasian National University*, 121 (№6), 18-24.

Pre & Post Pandemic Formative Assessments Methodologies to Improve Mathematics Learning for University Students

*Diana Audi*¹

¹*Department of Mathematics & Statistics, American University of Sharjah, UAE*

Daudi@aus.edu

Abstract

Formative assessments help teachers identify concepts that students are struggling to understand, skills they are having difficulty acquiring, or learning standards they have not yet achieved so that adjustments can be made to lessons, instructional techniques, and academic support. This paper focuses on a study at the American University of Sharjah, Mathematics Department of integrating the use of technology with formative assessments in order to identify students who are struggling and focus on specific points prior to exams rather than doing a general review. More specifically, the study was conducted over two different semesters. The first semester (before the pandemic), classroom response systems with PollEverywhere was used while in the second semester (after the pandemic), ILearn Formative Feedback was used. The uniqueness of this research study is the integration of formative assessments and feedback with technology in the delivery of Mathematics Subjects in Higher Education. Both quantitative and qualitative results were collected and there was evident and significant improvements in students' performance when the proposed formative feedback prior to exams was used. It was also apparent that after the pandemic, the use of Ilearn formative feedback was very beneficial for the students and helped improve their performances significantly in Mathematics.

Keywords: Ilearn, LMS, Polleverywhere, Formative Feedback, Mathematics Teaching, Technology in Education, Lecture Capture, Learning Objectives, Formative Assessments.

References:

1. J. Daniel, V. Preast, H. Bender, and L. Hassall, "Is the effectiveness of lecture capture related to teaching approach or content type?," *Computers & Education*, vol. 72, pp. 121–131, 2014.
2. J. de Boer, P. A. Kommers, and B. de Brock, "Using learning styles and viewing styles in streaming video," *Comput. Educ.*, vol. 56, no. 3, pp. 727–735, Apr. 2011.
3. Stephen Cummins, Alastair R. Beresford, and Andrew Rice, "Investigating Engagement with In-Video Quiz Qs in a Programming Course," *IEEE Trans On Lear. Tech.*, V. 9, No. 1, 2016
4. J. Campbell, D. Horton, M. Craig, and P. Gries, "Evaluating an inverted CS1," in *Proc. 45th ACM Tech. Symp. Comput. Sci. Educ.*, Mar. 2014, pp. 307–312.

Periodicity and Stability in the Kaldor Model of Economics with Impulse Effects

Diğdem ŞEN¹, Fatma TOKMAK FEN²

^{1,2}Department of Mathematics, Gazi University, 06560, Ankara, Turkey

digdem.sen@gazi.edu.tr

fatmatokmak@gazi.edu.tr

Abstract

The Kaldor model in economics possessing impulse effects is under investigation. It is shown that the model has a unique periodic solution under certain conditions. The asymptotic stability is rigorously proved.

Keywords: Kaldor model, Asymptotic stability, Periodic solution

Acknowledgement: D. Şen has been supported by the 2210/A program of The Scientific and Technological Research Council of Turkey (TÜBİTAK).

References:

1. A. M. Samoilenko, N.A. Perestyuk, Impulsive Differential Equations, World Scientific (1995).
2. W. B. Zhang, Differential Equations, Bifurcations, and Chaos in Economics, World Scientific, Singapore (2005)
3. H. W. Lorenz, Nonlinear Dynamical Economics and Chaotic Motion, Springer, New York (1993).
4. L. I. Dobrescu, D. Opris, Neimark-Sacker bifurcation for the discrete-delay Kaldor-Kalecki model, Chaos, Solitons and Fractals 41 (2009) 2405-2413.
5. J. Yua, M. Penga, Stability and bifurcation analysis for the Kaldor-Kalecki model with a discrete delay and a distributed delay, Physica A460 (2016) 66-75.

ICOMAA 2023

Adaptation of Some Graceful Graph Labeling Methods to Combinatorial Block Designs

Dilara Erdemir¹ and Emre Kolotođlu²

^{1,2}Department of Mathematics, Yildiz Technical University, Istanbul, Turkey

dilaraer@yildiz.edu.tr

kolot@yildiz.edu.tr

Abstract

In 1985, S. Lo introduced the notion of edge-graceful graphs. A graph $G = (V, E)$ is said to be edge-graceful if there exists a bijection f from E to $\{1, 2, \dots, |E|\}$ such that the induced mapping f^+ from V to $\{0, 1, \dots, |V| - 1\}$ given by $f^+(x) = (\sum f(xy)) \pmod{|V|}$ taken over all edges xy is a bijection. We adapt the edge-graceful graph labeling definition into block designs and define a block design (V, B) with $|V| = v$ and $|B| = b$ as block-graceful if there exists a bijection $f: B \rightarrow \{1, 2, \dots, b\}$ such that the induced mapping $f^+: V \rightarrow \mathbb{Z}_v$ given by $f^+(x) = \sum_{A \in B} f(A) \pmod{v}$ is a bijection in this work. We find a necessary condition and prove some basic results on the existence of block-graceful (v, k, λ) -BIBDs. Moreover, we study the existence problem of block-graceful Steiner triple systems, affine and projective geometries.

Keywords: Block-graceful design, Steiner triple system, affine geometry, projective geometry.

References:

1. C.C. Lindner, C.A. Rodger, Design Theory, 2nd edn., CRC Press, New York, (2008).
2. Charles J. Colbourn and Jeffrey H. Dinitz, Handbook of Combinatorial Designs, 2nd Edition. New York: Chapman and Hall/CRC, (2007).
3. D.R. Stinson, Combinatorial Designs: Constructions and Analysis, Springer-Verlag, New York, (2003).
4. J.A. Gallian, A Dynamic Survey of Graph Labeling, Twenty-second edn., The Electronic Journal of Combinatorics, December 15, (2019).
5. S. Lo, On edge-graceful labelings of graphs, Congr. Numer., 50 (1985) 231-241.

ICOMAA 2023

Embedding the stationary spacetimes into Brans-Dicke cosmology via conformal transformations

Dilek Kazıcı¹, Ali Demirözlü²

^{1,2}*Department of Physics, Tekirdağ Namık Kemal University,*
dkazici@nku.edu.tr
demirozluali@gmail.com

Abstract

A conformal transformation of a static or stationary spacetime by a time dependent conformal scale factor $S(\tau)^2$ is one of the methods of producing a cosmological spacetime. Using this knowledge and Brans-Dicke (BD) field equations, we investigate two time dependent metrics, including Friedmann-Lemaitre-Robertson-Walker (FLRW) spacetime and conformally transformed Kerr-Newman black hole, and we obtain solutions that allow different expansion rates for each geometry. These expansion rates depend on the matter content of the conformally transformed geometry. We state that the BD scalar field yields accelerated expansion of the conformal spacetime if the original metric has vacuum geometry, and no acceleration if the original spacetime has some curvature or matter content in it.

Keywords: Cosmology, FLRW spacetime, Kerr-Newman black hole, Scalar tensor theories.

References:

1. C.H. Brans, R.H. Dicke, Mach's Principle and a Relativistic Theory of Gravitation, Phys. Rev., vol. 124 (1961), no. 925.
2. R. M. Wald, General Relativity, Chicago University Press, Chicago, 1984.
3. M.P. Dabrowski, J. Garecki, Conformal transformations and conformal invariance in gravitation, Annalen Phys. (Berlin), vol. 18 (2009), pp. 13-32.
4. S.N.G. Thakurta, Kerr metric in an expanding universe, Indian J. Phys. 55B (1981), pp. 304-310.
5. H. Moradpour, A. Dehghani, M.T. Mohammadi Sabet, Dynamic black holes in an FRW background: Lemaitre transformations, Mod. Phys. Lett. A, vol. 30 (2015), 1550207.

ICOMAA 2023

On a New Type Gaussian Sequences

Ece Gulsah Colak¹, Nazmiye Gonul Bilgin², Yuksel Soykan³

^{1,2,3}Department of Mathematics, Zonguldak Bulent Ecevit University,

egsavkli@beun.edu.tr

nazmiyegonul@beun.edu.tr

yuksel_soykan@hotmail.com

Abstract

In this work, we investigate several properties of a new complex sequence, such as Binet formula, generating function, summation formulas, identities and matrix formulation. We also examine Lucas generalization of these generalized Gaussian special numbers.

Keywords: Tribonacci numbers, Tribonacci-Lucas numbers, Gaussian Tribonacci Numbers, Gaussian Tribonacci-Lucas numbers.

References:

1. A. F. Horadam, Complex Fibonacci Numbers and Fibonacci Quaternions, American Mathematics Monthly, 70, 289-291, (1963).
2. S. Pethe and A. F. Horadam, Generalized Gaussian Fibonacci Numbers, Bull. Austral. Math. Soc., 33, 37-48, (1986).
3. S. Halıcı and S. Öz, On Some Gaussian Pell and Pell-Lucas Numbers, Ordu Üniversitesi Bilim ve Teknoloji Dergisi, 6 (1), 8-18, (2016).
4. Y. Soykan, E. Taşdemir, İ. Okumuş, M. Göcen, Gaussian Generalized Tribonacci Numbers, Journal of Progressive Research in Mathematics, 14 (2), (2018).
5. Y. Soykan, Generalized John Numbers, Journal of Progressive Research in Mathematics, 19 (1), (2022).

ICOMAA 2023

On the Factorial of Split Quaternions

Edanur Ergül¹, Salim Yüce²

¹Department of Mathematics, Marmara University and Yildiz Technical University,

²Department of Mathematics, Yildiz Technical University,

edanur.ergul@marmara.edu.tr

sayuce@yildiz.edu.tr

Abstract

In the literature, the Gamma function is defined for complex number z with a positive real part and its domain can be extended to complex plane with the exception of the values zero and negative integers, [2-3]. Moreover, the Gamma function is also defined for real quaternion q with a positive real part, [1]. In addition, the real quaternionic binomial coefficient and the inverse of real quaternionic binomial coefficient are defined and their properties are investigated.

In this study, we have used the set of split quaternions with the metric $(+, +, -, -)$. The aim of this study is to contribute to the literature by defining the split quaternionic gamma function, split quaternionic binomial coefficient and inverse of split quaternionic binomial coefficient for split quaternions and examining their properties.

Keywords: Real Quaternion, Split Quaternion, Factorial, Gamma Function, Binomial Coefficient, Inverse of Split Quaternionic Binomial Coefficient.

References:

1. Hogan, J. A., Massopust, P. (2017). Quaternionic B-splines. Journal of Approximation Theory, 224, 43-65. <https://doi.org/10.1016/j.jat.2017.09.003>
2. Davis, P. J. (1959). Leonhard Euler's Integral: A Historical Profile of the Gamma Function: In Memoriam: Milton Abramowitz. The American Mathematical Monthly, 66(10), 849-869. <https://doi.org/10.2307/2309786>
3. Shorey, T. N. (2021). Complex Analysis with Applications to Number Theory. Springer Singapore. <https://doi.org/10.1007/978-981-15-9097-9>
4. Alagöz, Y., Oral, K. H., Yüce, S. (2012). Split Quaternion Matrices. Miskolc Mathematical Notes, 13(2), 223-232. <https://doi.org/10.18514/MMN.2012.364>
5. Spivey, M.Z. (2019). The Art of Proving Binomial Identities (1st ed.). Chapman and Hall/CRC. <https://doi.org/10.1201/9781351215824>

Spectral Analysis of a Singular Dissipative Third-Order Differential Operator

Ekin Uğurlu¹

¹*Department of Mathematics, Çankaya University,
ekinugurlu@cankaya.edu.tr*

Abstract

In this talk, we investigate the spectral properties of a singular dissipative third-order differential operator with deficiency indices $(3,3)$ at singular point with the aid of the resolvent operator. In particular, after showing that the resolvent operator is a trace-class operator, using Lidskii's theorem we introduce some completeness theorems. Moreover we share the trace of the resolvent operator as well as some additional results on the resolvent operator.

Keywords: Dissipative operators, Spectral analysis, Resolvent operators.

References:

1. C.L. Dolph, Recent developments in some non-self-adjoint problems of mathematical physics, Bull. Amer. Math. Soc. 67 (1961), 1-69.
2. M.S. Livsic, On spectral decomposition of linear non-sel-adjoint operators, Transl. Amer. Math. Soc. II 5 (1957), 67-114.
3. E. Bairamov and A.M. Krall, Dissipative operators generated by the Sturm-Liouville expression in the Weyl limit circle case, J. Math. Anal. Appl. 254 (2001), 178-190.
4. D.B. Hinton, Deficiency indices of odd-order differential operators, Rocky Moun. J. Math. 8 (1978), 627-640.
5. J.C. Ferreira, V.A. Menegatto, C.P. Olivera, On the neclearity of integral operators, Positivity, 13 (2009), 519-541.

ICOMAA 2023

Dependence of Eigenvalues of Dirac Systems on the Data

Ekin Uğurlu¹

¹*Department of Mathematics, Çankaya University,
ekinugurlu@cankaya.edu.tr*


Abstract

In this talk, we show that the eigenvalues of a regular boundary value problem generated by a Dirac system depend continuously on the data. Moreover, we share some derivatives of the eigenvalues with respect to some elements of the data.

Keywords: Dirac systems, Frechet derivatives, Eigenvalues of boundary-value problems.

References:

1. J. Poeschel and E. Trubowitz, Inverse Spectral Theory, Academic Press, New York, 1987.
2. Q. Kong and A. Zettl, Eigenvalues of regular Sturm-Liouville problems, J. Differ. Eq. 131 (1996), 1-19.
3. Q. Kong, H. Wu, A. Zettl, Dependence of the nth Sturm-Liouville eigenvalue on the problem, J. Differ. Eq. 156 (1999), 328-354.
4. B.M. Levitan and I.S. Sargsjan, Sturm-Liouville and Dirac Operators, London, UK, Kluwer, 1991.



ICOMAA 2023

Some New Results of Weighted Generalized Norlund-Euler Statistical Convergence

Ekrem Aljimi¹, Penpark Sirimark², Agon Mahmuti³

¹ Faculty of Computer Science, Public University "Kadri Zeka", 60000 Gjilan, Kosovë,

² Department of Science and Mathematics, Rajamangala University of Technology Isan Surin Campus, Surin 32000, Thailand,

³ Department of Mathematics, University of Sarajevo, Zmaj od Bosne 33-35, 71000 Sarajevo, Bosnia and Herzegovina,

ekrem.halimi@uni-gjilan.net

penpark.si@rmuti.ac.th@gmail.com

agon.mahmuti@pmf.unsa.ba

Abstract

In this paper, we defined generalized weighted Norlund-Euler statistical convergence. We also proved some properties of this type of statistical convergence by applying generalized weighted Norlund-Euler summability method. Moreover, we prove Korovkin's theorem using generalized weighted Norlund-Euler summability theorem for functions on general. We also study rate of generalized weighted Norlund-Euler statistical convergence and some sequences spaces defined by Orlicz functions.

Keywords: Statistical convergence; Generalized weighted Norlund-Euler summability method; Korovkin's type; approximation theorem; Rate of generalized weighted Norlund-Euler statistical convergence; Orlicz functions.

References:

1. Fast, H. Sur la convergence statistique. Colloq. Math. 1951, 2, 241–244.
2. Aljimi, E.; Sirimark, P.; Ramizi, A.; Mahmuti, A. Weighted $(E\lambda, q)(C\lambda, 1)$ Statistical Convergence and Some Results Related to This Type of Convergence. Symmetry 2022, 14, 2363. <https://doi.org/10.3390/sym14112363>.
3. Braha, N.L.; Srivastava, H.M.; Et, M. Some weighted statistical convergence and associated Korovkin and Voronovskaya type theorems. J. Appl. Math. Comput. 2021, 65, 429–450.
4. Özger, F. Weighted statistical approximation properties of univariate and bivariate λ -Kantorovich operators. Filomat 2019, 33, 3473–3486.

ICOMAA 2023

Zero-Hopf bifurcation of fourth-order quadratic polynomial differential systems, via averaging theory of third order

Bendib El Ouahma¹, Makhlof Amar²

¹ Department of Mathematics, University of Skikda 20 August 1955, Algeria,

² Department of Mathematics, University of Badji Mokhtar Annaba, Algeria,

bendib_e@yahoo.fr, ou.bendib@univ-skikda.dz
makhlofamar@yahoo.fr

Abstract

The goal of this work is to study the maximum number of limit cycles that can be bifurcate from a zero-Hopf equilibrium point of a quadratic polynomial differential system in \mathbb{R}^4 , by using the averaging theory of third order.

Keywords: Zero-Hopf bifurcation, averaging theory, quadratic polynomial differential system.

References:

1. N.N. Bautin, On the number of limit cycles which appear with the variation of the coefficients from an equilibrium position of focus or center type, *Translations of the Amer. Math. Soc.* 1 (1954), 396-413.
2. E. Bendib, S. Badi and A. Makhlof, On the 3-Dimensional Hopf bifurcation via averaging theory of third order, *Turk J Math* (2017) 41 : 1053-1071.
3. D. Djedid, E. Bendib and A. Makhlof, Four-dimensional Zero-Hopf Bifurcation of Quadratic Polynomial Differential System, via Averaging Theory of Third Order, *Journal of Dynamical and Control Systems* 28(4) (2021).
4. J.A. Sanders and F. Verhulst, *Averaging methods in nonlinear dynamical systems*, Applied Mathematical Sciences, New York, Springer-Verlag, 1985.

ICOMAA 2023

Dissipative extensions of one operator with potential in the form of a Dirac function

Eldar M. Mustafayev
Baku State University, Baku, Azerbaijan
eldarmustafa59@gmail.com

Abstract

The paper describes all dissipative extensions of the minimal operator generated by the differential expression $-\Delta + \sum_{s=1}^n \delta(x - \xi_s)$, where $\delta(x)$ – the Dirac function, $\xi_s \in E_3$, $\xi_s \neq \xi_{s'}, s \neq s'$.

Keywords: the Dirac function, minimal operator, dissipative extensions of the operator.

References:

1. A.V. Strauss, On extensions and the characteristic function of a symmetric operator. Izv. USSR Academy of Sciences, Ser. Mat., 32 (1), (1968).
2. Naimark M.A. Linear differential operators. M.: Nauka. (1969).
3. M.L. Gorbachuk, A.N. Kochubey, M.A. Rybak, Dissipative extensions of differential operators in the space of vector-functions. Doklady AN SSSR, (1972), 205 (5), 1029-1032.



ICOMAA 2023

DRBEM Solution of MHD Flow in a Rectangular Duct under Axially-changing External Magnetic Field

Elif Ebr̄en Kaya¹, M̄inevver Tezer-Sezgin²

¹The Scientific and Technological Research Council of Turkey,

²Department of Mathematics, Middle East Technical University

elifebren90@gmail.com

munt@metu.edu.tr

Abstract

The laminar, fully-developed magnetohydrodynamic (MHD) pipe flow of a viscous and incompressible fluid has been considered between consecutive magnets placed on the pipe-axis. The flow is under the effect of an axial-dependent applied magnetic field. The MHD flow equations are transformed to three nonlinear Poisson type equations in terms of velocity, induced magnetic field and electric potential, and they are solved by using the dual reciprocity boundary element method (DRBEM) with the fundamental solution of Laplace's equation. The study shows that, axially-changing magnetic field makes the flow to turn its direction at a certain position of the axis. The problem parameters, Hartmann number M and magnetic Reynolds number R_m effects on the flow behavior contrast with each other in the sense that, the lengths of the intervals on the pipe-axis on which the flow is reversed are increasing as M increases, however, they are getting shorter as R_m increases.

Keywords: DRBEM, MHD duct flow, axially-changing magnetic field.

References:

1. P.W. Partridge, C.A. Brebbia and L.C. Wrobel, The Dual Reciprocity Boundary Element Method, Computational Mechanics Publications, (1992), Southampton Boston.
2. U. Muller and L. Buhler, Magnetohydrodynamics in Channels and Containers, (2001), Springer.
3. A. Sterl, Numerical Simulation of Liquid-metal MHD Flows in Rectangular Ducts, Journal of Fluid Mechanics. 216 (1990) 161-191.

ICOMAA 2023

On k - Oresme Hybrid Polynomials

Serpil Halıcı¹ and Elifcan Sayın²

^{1,2}*Department of Mathematics, Pamukkale University,*

shalici@pau.edu.tr

elifcan7898@gmail.com

Abstract

In this study, we examined both k - Oresme hybrid polynomials and the types of these polynomials with negative indices. Given all the remarkable properties of the polynomial sequence, we also derived some fundamental identities provided by the elements of these polynomials.

Keywords: Oresme polynomials, Hybrid numbers, Recurrence relations.

References:

1. Oresme, N. Quaestiones super geometriam Euclidis (Vol. 3) (1961) *Brill Archive*.
2. Horadam, A. F. Oresme numbers. *Fibonacci Quarterly*, 12(3) (1974) 267-271.
3. Cerda-Morales, G. Oresme polynomials and their derivatives. (2019) arxiv preprint arXiv:1904.01165
4. Halıcı, S., Sayın, E. On Some k - Oresme Hybrid Numbers. *Utilitas Mathematica*, 120 (2023) 1-11.
5. Szynal-Liana, A., Wloch, I. Oresme Hybrid Numbers and Hybrationals. *Kragujevac Journal of Mathematics*, 48(5) (2024) 747-753.

ICOMAA 2023

An Inverse Coefficient Problem for a Two-Dimensional Hyperbolic Equation with Integral Overdetermination

Elvin Azizbayov¹ and Gulnara Shukurova²

¹Department of Control for Intelligent Systems,
Academy of Public Administration under the President of the Republic of Azerbaijan,
eeazizbayov@dia.edu.az

²Department of Higher Mathematics, Baku State University
shukurova-gulnara@rambler.ru

Abstract

The paper considers a classical solvability of a nonlinear inverse boundary value problem for a two-dimensional hyperbolic equation with an integral overdetermination condition. To investigate the solvability of the stated problem, we first consider an auxiliary inverse boundary value problem and prove its equivalence to the original problem. Then we prove the existence and uniqueness theorem for the auxiliary problem by the contraction mappings principle. Furthermore, based on the equivalency of these problems, the existence and uniqueness theorem for the classical solution of the original inverse problem is proved.

Note that the related inverse problems for one-dimensional hyperbolic equation was studied in the papers [1-3] and the references therein.

Keywords: Inverse problem, hyperbolic equation, classical solution, existence, uniqueness.

References:

1. Azizbayov E.I. Inverse coefficient identification problem for a hyperbolic equation with nonlocal integral condition. Turk. J. Math. Vol. 46, No.4 (2022), 1243-1255.
2. Mehraliyev Y.T., Azizbayov E.I. A time-nonlocal inverse problem for a hyperbolic equation with an integral overdetermination condition. Electron. J. Qual. Theory Differ. Equ. Vol. 2021, No.29 (2021), 1-12.
3. Mehraliyev Ya.T., Isgenderova Q.N. Inverse boundary value problem for a second-order hyperbolic equation with integral condition of the first kind. PFMT. No.1(26) (2016), 42-47.

ICOMAA 2023

Numerical Investigation of Stiff Epidemiological Models

Emel Kurul¹, Murat Sari², Huseyin Tunc³

¹*Department of Mathematics, Yildiz Technical University,*

²*Department of Mathematics Engineering, Istanbul Technical University*

³*Department of Biostatistics and Medical Informatics, Bahcesehir University*

emel.kurul@std.yildiz.edu.tr

muratsari@itu.edu.tr

huseyin.tunc@bau.edu.tr

Abstract

Mathematical models play an important role in understanding natural laws and predicting how they shape the real world. In epidemiology, these models are used to simulate the spread of infectious diseases and forecast the number of individuals who will become infected over time. An essential characteristic of ordinary differential equation based epidemiological models is the presence of stiffness, which can arise due to different parameter settings. Stiff models are difficult to solve using standard numerical methods, as they require small time steps and significant computational resources. This study investigates the suitability of MATLAB numerical solvers, such as ode15s, ode23s and ode45, for simulating stiff epidemiological models and evaluates their performance in terms of computational efficiency.

Keywords: Mathematical epidemiology, SIR model, SEIR model, Time integration, Stiff ODEs.

References:

1. D'Ambrosio, R., Diaz de Alba, P., Giordano, G., & Paternoster, B. (2022). A Modified SEIR Model: Stiffness Analysis and Application to the Diffusion of Fake News. [Computational Science and Its Applications, ICCSA 2022](#)
2. [Hass, F., Kröger, M.](#) & Schlickeiser, R. (2022). Multi-Hamiltonian structure of the epidemics model accounting for vaccinations and a suitable test for the accuracy of its numerical solvers. [Journal of Physics A: Mathematical and Theoretical](#), 55(22).
3. Higham, DJ. & Trefethen, LN. (1993). Stiffness of ODEs. *Bit Numerical Mathematics*, Springer, 33 (2).
4. Byrne, G. D. & Hindmarsh, A. C. (1987). Stiff ODE Solvers: A Review of Current and Coming Attractions. *Journal of Computational Physics*, 70, 1-62.
5. Enrighte, W. H. & HULL, T. E. (1976). Comparing Numerical Methods for the Solution of Stiff Systems of ODES Arising in Chemistry. *Numerical Methods for Differential Systems*, edited by L. Lapidus and W. E. Schiesser (Academic Press, New York, p. 45.).

***ss*-Injective Modules**

Emine Önal Kır¹

¹*Department of Mathematics, Kırşehir Ahi Evran University,*
emine.onal@ahievran.edu.tr

Abstract

In this study, we define the concept of *ss*-injective modules, which lies between the class of injective modules and almost injective modules. We say that a module M is *ss*-injective provided M is an *ss*-supplement submodule of every module which contains M . We compare this concept with other injective module generalizations. We show that *ss*-injective property is inherited by direct summands. We also investigate *ss*-injective modules over various ring types.

Keywords: *ss*-injective modules, *ss*-supplement submodules, left V -ring, left small ring.

References:

1. E. Büyükaşık and R. Tribak, Rad-supplements in injective modules, *Algebra and Discrete Mathematics* 22(2) (2016) 171-183.
2. J. Clark, D. Keskin Tütüncü and R. Tribak, Supplement submodules of injective modules, *Communications in Algebra* 39 (2011) 4390-4402.
3. E. Kaynar, H. Çalışıcı and E. Türkmen, *ss*-Supplemented modules, *Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics* 69(1) (2020) 473-485.
4. D. W. Sharpe and P. Vámos, *Injective Modules*, Cambridge University Press, 1972.
5. R. Wisbauer, *Foundations of Module and Ring Theory*, Gordon and Breach, Philadelphia, 1991.

ICOMAA 2023

Power Series Statistical Convergence of Cheney-Sharma Operators

Emre GÜVEN¹ and Dilek SÖYLEMEZ²,
^{1,2}Department of Mathematics, Selcuk University,
emreguven88@gmail.com

Abstract

In this presentation, we investigate some Korovkin type approximation properties of the Cheney-Sharma operators via power series statistical convergence. We also compute the rates of convergence by modulus of continuity and the elements of Lipschitz class.

Keywords: Power series statistical convergence; Cheney-Sharma operators; rate of convergence

References:

1. Altomare F. and Campiti, M. Korovkin-type approximation theory and its applications, Walter de Gruyter, Berlin-New York, 1994.
2. Cheney, E.W., Sharma A., On a generalization of Bernstein polynomials. Riv. Mat. Univ. Parma 2(5) (1964) 77–84
3. Soylemez, D., Ünver, M.(2017) Korovkin type theorems for Cheney-Sharma Operators via summability methods. Results Math. 72(3),1601–1612
4. Söylemez, D., Ünver, M. (2021). “Rates of Power Series Statistical Convergence of Positive Linear Operators and Power Series Statistical Convergence of -Meyer–König and Zeller Operators”, Lobachevskii Journal of Mathematics, 42 (2), 426-434.
5. Ünver, M. and Orhan, C.(2019). Statistical convergence with respect to power series methods and applications to approximation theory. Journal Numerical Functional Analysis and Optimization. 40(5), 535-547,

ICOMAA 2023

High Order Numerical Approach for Solving the Modified Equal Width Equation

Emre Kırılı

Graduate School of Natural and Applied Sciences, Eskisehir Osmangazi University,
emre.26@windowslive.com

Abstract

The Modified Equal Width (MEW) equation

$$u_t + 3u^2u_x - \mu u_{xxt} = 0$$

was explored by Morrison et al [1] to model the nonlinear dispersive waves. Since the analytical solutions of the MEW equation are obtained only for restricted solution set of initial and boundary conditions, it has gained great importance to obtain the numerical solutions of the MEW equation over the last years. In the present work, a high order computational approach is introduced to find the approximate solutions of the MEW equation. This approach is formed by using quintic trigonometric B-spline collocation procedure in the spatial discretization and fourth-order two step scheme in the temporal discretization. As a test problem, the motion of the single solitary wave is examined. To see accuracy of the proposed approach and to show the improvement in the results, the L_∞ error norm is computed and compared with the existing techniques in the literature. The obtained results verify that the proposed approach has marked advantage over some existing studies in terms of accuracy. Also, the temporal order of convergence and three invariant constants are computed numerically and found to agree with their theoretical.

Keywords: Collocation method, Trigonometric B-splines, Modified equal width equation, Solitary waves.

References:

1. Morrison P.J., Meiss J.D., & Carey J.R. (1984). Scattering of Regularized-Long-Wave Solitary Waves. *Physica D: Nonlinear Phenomena*, 11(3), 324-336.
2. Saka B. (2007). Algorithms for numerical solution of the modified equal width wave equation using collocation method. *Mathematical and Computer Modelling*, 45(9), 1096-1117.
3. Celikkaya I. (2019). Operator splitting method for numerical solution of modified equal width equation. *Euro-Tbilisi Mathematical Journal*, 12(3), 51-67.
4. Karakoc S.B.G., Zeybek H. (2016). A cubic B-spline Galerkin approach for the numerical simulation of the GEW equation. *Statistics, Optimization & Information Computing*, 4(1), 30-41.

Effect of Caputo-Fabrizio Derivative on Scabies Model

Esmehan Uçar¹

¹*Department of Mathematics, Balıkesir University,
esucarr@gmail.com*

Abstract

A tiny burrowing mite known as *Sarcoptes scabiei* causes the itching skin condition known as scabies. In a family, daycare center, classroom, or nursing home, close person-to-person contact can quickly spread scabies, which really is contagious, fairly quickly. In this study, Caputo-Fabrizio (CF) derivative is originally used to remodel a scabies disease model and what's more we closely analyse numerical aspects of this fractional model and generate some graphical outcomes in order to forecast difficult and complicated occurrences. With a greater understanding of what is vital to public health research and decision-making processes, such analyses will be beneficial to subsequent research efforts.

Keywords: Caputo-Fabrizio derivative, scabies model

References:

1. M. Caputo and M. Fabrizio, A new definition of fractional derivative without singular kernel, *Progress in Fractional Differentiation and Applications*, 1 (2015), 73-85.
2. Y. T. Toh, C. Phang and J.R. Loh, New predictor-corrector scheme for solving nonlinear differential equations with Caputo-Fabrizio operator, *Mathematical Methods in the Applied Sciences*, 42(2019), 175-185.
3. S. Rashid, B. Kanwal, F. Jarad and S. K. Elagan, A peculiar application of the fractal–fractional derivative in the dynamics of a nonlinear scabies model, *Results in Physics*, 38 (2022), 105634.
4. E. Uçar, S. Uçar, F. Evirgen and N. Özdemir, A Fractional SAIDR Model in the Frame of Atangana–Baleanu Derivative, *Fractal and Fractional* 5(2) (2021), 18 pages.
5. Z. Hammouch, M. Yavuz and N. Özdemir, Numerical solutions and synchronization of a variable-order fractional chaotic system. *Mathematical Modelling and Numerical Simulation with Applications*, 1(1) 2021, 11-23.

ICOMAA 2023

Alternative Solution Method on Matrix Games with Fuzzy Goals

Adem Cengiz Cevikel¹, Esra Soslu²

^{1,2}Department of Mathematics, Yildiz Technical University,
acevikel@yildiz.edu.tr
esrasoslu@hotmail.com

Abstract

Two-person zero-sum games have been widely studied in game theory. In these games, each player is mainly interested in maximizing her/his payoff by choosing the best strategy. In the crisp scenario, this is done by maximizing an objective function under the linear constraints. However, the player cannot always be in the crisp scenario and can have to play under the fuzzy environment. One of the situation that fuzziness can occur in two-person zero-sum games is that player's ambiguous preferences. This kind of games is expressed as the matrix games with fuzzy goals. Nishizaki and Sakawa (2001) considered fuzzy goals as player's degree of satisfaction for a payoff. They suggested a solution method for solving these games by constructing linear programming problems for both players. The method consists of a membership function that is aimed to be optimized for each fuzzy goal. In this work, it is studied whether the membership function in the optimization problem can also be directly applied to the elements of payoff matrix. In this way, it was questioned whether a more direct solution method could be obtained. Multi-objective games with fuzzy goals are examined as well as single objective games.

Keywords: Two-person zero-sum games, fuzzy goal, membership function, linear programming.

References:

1. Barron, E.N., (2013). *Game Theory: An Introduction* (2nd ed.). Wiley.
2. Bector, C.R., Chandra, S., (2005). *Fuzzy Mathematical Programming and Fuzzy Matrix Games*. Springer Berlin Heidelberg, New York.
3. Klir, G.J., Yuan, B. (1995). *Fuzzy Sets and Fuzzy Logic: Theory and Applications*. Prentice Hall.
4. Nishizaki I., Sakawa M. (2001). *Fuzzy and Multiobjective Games for Conflict Resolution*. Physica-verlag, Heidelberg.
5. Zadeh, L.A. (1965). Fuzzy Sets. *Information and Control*, 8(3), 338-353. [https://doi.org/10.1016/S0019-9958\(65\)90241-X](https://doi.org/10.1016/S0019-9958(65)90241-X)

ICOMAA 2023

A Note on Singular Integral Operators Associated with Laplace-Bessel Differential Operator in Variable Lorentz Space

Esra Kayal¹, Canay Aykol²

A Local Taylor Series Method for Non-linear Singular Boundary Value Problems

Esra Celik¹, Huseyin Tunc², Murat Sari³

¹Department of Mathematics, Yildiz Technical University,

²Department of Biostatistics and Medical Informatics, Bahcesehir University

³Department of Mathematics Engineering, Istanbul Technical University

esra.celik@std.yildiz.edu.tr

huseyin.tunc@bau.edu.tr

muratsari@itu.edu.tr

Abstract

In this paper, we propose the utility of the implicit-explicit local differential transformation method (IELDTM) based on the local Taylor series approximation for solving two-point singular boundary value problems. These problems arise in many fields of science such as gas dynamics, nuclear physics, atomic structures, and electro hydrodynamics, etc. First, we give some basic information to explain the mathematical structure of the method, then consider the results produced by the method. We show that IELDTM eliminates the disadvantages of existing differential transform-based methods. It has been shown that the IELDTM yields more accurate solutions with fewer CPU times than the rival numerical methods.

Keywords: implicit-explicit local differential transform method, Taylor series, differential transform method, non-linear singular boundary value problems.

References:

1. T. Huseyin and M. Sari, An implicit-explicit local differential transformation method for parabolic partial differential equations, *Engineering Computations*. 39.3 (2022) 1020-1037.
2. P. Roul and V. M. K. Goura, A fast numerical scheme for solving singular boundary value problems arising in various physical models, *Journal of Mathematical Chemistry*. (2022) 1-28.
3. P. Roul and T. Kumari, A quartic trigonometric B-spline collocation method for a general class of nonlinear singular boundary value problems, *Journal of Mathematical Chemistry*. 60(1)(2022) 128-144.
4. Y. Lin, H. W. Tang and C. O. K. Chen, Modified differential transform method for two singular boundary value problems, *Journal of Applied Mathematics*. (2014).

Mechanistic Modelling of Contact Tracing and Isolation Policies

Esra Ozge Asan¹, Murat Sari², Enes Seyfullah Kotil³, Huseyin Tunc⁴

¹Department of Mathematics, Yildiz Technical University,

²Department of Mathematics Engineering, Istanbul Technical University,

³Department of Molecular Biology and Genetics, Boğaziçi University,

⁴Department of Biostatistics and Medical Informatics, Bahcesehir University

ozgeasan@std.yildiz.edu.tr

muratsari@itu.edu.tr

enesseyfullah.kotil@boun.edu.tr

huseyin.tunc@med.bau.edu.tr

Abstract

The COVID-19 pandemic has become a global crisis since its emergence, affecting all aspects of life, including social, economic, physical, and mental health. Various intervention policies have been implemented since the early days of the disease, such as the use of masks, social distancing, contact tracing and isolation, lockdowns, and other restrictions. In this study, we derive the S(Susceptible)- E(Exposed)- (P)Presymptomatic- A(Asymptomatic)- D (Symptomatic)- R(Reported) model including contact tracing and isolation policies. The SEPADR model is represented by the system of integro-differential equations.

Keywords: COVID-19, Compartmental model, SEIR model, Integro-differential equations

References:

1. Chen, Y., Huang, H. (2022). Modeling the impacts of contact tracing on an epidemic with asymptomatic infection. *Applied Mathematics and Computation*, 416.
2. Ge, Y., Chen, Z., Handel, A., Martinez, L., Xiao, Q., Li, C., Chen, E., Pan, J., Li, Y., Ling, F., & Shen, Y. (2021). The impact of social distancing, contact tracing, and case isolation interventions to suppress the COVID-19 epidemic: A modeling study. *Epidemics*, 36.
3. Wu, P., Liu, F., Chang, Z., Lin, Y., Ren, M., Zheng, C., Li, Y., Peng, Z., Qin, Y., Yu, J., Geng, M., Yang, X., Zhao, H., Li, Z., Zhou, S., Ran, L., Cowling, B. J., Lai, S., Chen, Q., ... Li, Z. (2021). Assessing Asymptomatic, Presymptomatic, and Symptomatic Transmission Risk of Severe Acute Respiratory Syndrome Coronavirus 2. *Clinical Infectious Diseases*, 73(6), E1314–E1320.
4. Giordano, G., Blanchini, F., Bruno, R., Colaneri, P., di Filippo, A., di Matteo, A., & Colaneri, M. (2020). Modelling the COVID-19 epidemic and implementation of population-wide interventions in Italy. *Nature Medicine*, 26(6), 855–860.

Thermoelastic laminated Timoshenko beam with distributed Delay: global existence and stability

Fares Yazid¹, Fatima Siham Djaradi² and Nor-El-Houda Beghersa³

¹Department of Mathematics, Amar Teledji Laghouat University, Algeria
f.yazid@lagh-univ.dz¹

²Department of Mathematics, Amar Teledji Laghouat University, Algeria
fs.djeradi@lagh-univ.dz²

³Department of Mathematics, University of Sciences and Technology of Oran,
Mohamed Boudiaf (USTO-MB)
norelhouda.beghersa@univ-usto.dz³

Abstract

In this work, we consider a linear thermoelastic laminated Timoshenko beam with distributed delay, where Cattaneo as law gives the heat conduction. We establish the well posedness of the system. For stability results, we prove exponential and polynomial stabilities of the system for the cases of equal and nonequal speeds of wave propagation. Our work is a natural extension of Feng's work in [4].

Keywords: Well-Posedness, Exponential Decay, Polynomial Decay, Thermoelastic Laminated, Distributed delay term.

References:

1. M. Afilal and S. Messaoudi, Stabilization of coupled hyperbolic equations with a heat equation of second sound, Mediterranean Journal of Mathematics, vol. 14, no. 2 (2017) P.39.
2. M. S. Alves and R. N. Monteiro, Exponential stability of laminated Timoshenko beams with boundary & internal controls, Journal of Mathematical Analysis and Applications. vol. 482 (2020).
3. Chen Z., Liu W. and Chen D. General decay rates for a laminated beam with memory. Taiwanese Journal of Mathematics. Vol. 23, No. 5 (2019) (25 pages).
4. B. Feng, Well-posedness and exponential decay for laminated Timoshenko beams with time delays and boundary feedbacks, Mathematical Methods in the Applied Sciences, vol. 41, no3, (2018), 1162-1174.

On Some Beta Type Operators

Faruk Özger

Department of Engineering Sciences, İzmir Katip Çelebi University, İzmir, Türkiye
farukozger@gmail.com

Abstract

Benrtein type operators have been widely studied by many researchers. In this presentation, we focus on some beta type operators and their approximation properties such as a direct approximation theorem by means of modulus of continuity and approximation rate for functions having derivatives of bounded variation. We support the theoretical parts by computer graphics.

Keywords: Bernstein-type operators, bounded variation, beta type operators, rate of approximation, approximation properties.

References:

1. Kajla, A., Özger, F., & Yadav, J. (2022). Bézier-Baskakov-Beta Type Operators. *Filomat*, 36(19), 6735-6750.
2. Adell, J., Badia, F. G., & de la Cal, J. (1993). Beta-type operators preserve shape properties. *Stochastic processes and their applications*, 48(1), 1-8.
3. Alotaibi, A., Özger, F., Mohiuddine, S. A., & Alghamdi, M. A. (2021). Approximation of functions by a class of Durrmeyer–Stancu type operators which includes Euler’s beta function. *Advances in Difference Equations*, 2021(1), 1-14.
4. Qasim, M., Khan, A., Abbas, Z., Özger, F., & Raina, P. (2022). Approximation by generalized Baskakov-beta operators and its convergence properties. *The Journal of Analysis*, 1-17.

ICOMAA 2023

On Some Beta Type Operators

Faruk Özger

Department of Engineering Sciences, İzmir Katip Çelebi University, İzmir, Türkiye
farukozger@gmail.com

Abstract

Benrtein type operators have been widely studied by many researchers. In this presentation, we focus on some beta type operators and their approximation properties such as a direct approximation theorem by means of modulus of continuity and approximation rate for functions having derivatives of bounded variation. We support the theoretical parts by computer graphics.

Keywords: Bernstein-type operators, bounded variation, beta type operators, rate of approximation, approximation properties.

References:

1. Kajla, A., Özger, F., & Yadav, J. (2022). Bézier-Baskakov-Beta Type Operators. *Filomat*, 36(19), 6735-6750.
2. Adell, J., Badia, F. G., & de la Cal, J. (1993). Beta-type operators preserve shape properties. *Stochastic processes and their applications*, 48(1), 1-8.
3. Alotaibi, A., Özger, F., Mohiuddine, S. A., & Alghamdi, M. A. (2021). Approximation of functions by a class of Durrmeyer–Stancu type operators which includes Euler’s beta function. *Advances in Difference Equations*, 2021(1), 1-14.
4. Qasim, M., Khan, A., Abbas, Z., Özger, F., & Raina, P. (2022). Approximation by generalized Baskakov-beta operators and its convergence properties. *The Journal of Analysis*, 1-17.

ICOMAA 2023

Lie Group Property of Generalized Linear Deformation Transformation

Fatih Tuğrul¹, Şenay Baydaş² and Bülent Karakaş³

¹Department of Mathematics, Van Yuzuncu Yil University,

fatihtugrul@yyu.edu.tr

²Van Yüzüncü Yil University

senaybaydas@gmail.com

³Bartın University

bulentkarakas@gmail.com

Abstract

In this study, using the definition of linear anisotropic transformation in 3-dimensional Euclidean space E^3 , we have defined a linear generalized deformation transformation. Unlike homothetic rotation matrices, we have shown that this transformation consists of the product of different expansion (or shrinkage) and different infinitesimal orthogonal matrices for each $x \in \mathbb{R}^3$ and $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}^+$ scalars. We have shown that matrix B is a regular matrix and that this set of matrices has a group structure. We examined the Lie group structure of this group.

Keywords: Anisotropic transformation, Linear anisotropic transformation, Generalized linear deformation transformation, Lie group

References:

1. Brickell, F., Clark, R. S., 1970. **Differentiable Manifolds**. Van Nostrand Reinhold Company Ltd., London.
2. Courant, R., Hilbert, D., 1953. **Methods of Mathematical Physics Vol. I**. Jhon Wiley & Sons., New York. (Translated and revised from the German original, into English)
3. Mortenson, M., M., 2009. **Geometric Transformation for 3D Modeling**. Industrial Press Inc., New York.

ICOMAA 2023

A Note About Pseudospectrum Of The Tensor Product Operator

*Fatih YILMAZ*¹

¹ Graduate School of Natural and Applied Sciences, Karadeniz Technical University,
fatih.yilmaz@ktu.edu.tr

Abstract

The connection between the ε -pseudospectrum of the tensor product operator $A_1 \otimes A_2$ and the ε -pseudospectrums of A_1 and A_2 has been investigated and some results are given about the this connection under certain conditions.

Keywords: Tensor product, Pseudospectrum.

References:

1. A. Brown, C. Percy, Spectra of tensor products of operators, Proc. Amer. Math. Soc., 17 (1966) 162-166.
2. C.S. Kubrusly, A concise introduction to tensor product, Far East J. Math. Sci., 22 (2006), 137-174.
3. L.N. Trefethen, Pseudospectra of linear operators, SIAM Rev. 39(3) (1997), 383-406.
4. J. Cui, C. Li, Y. Poon, Pseudospectra of special operators and pseudospectrum preservers, Journal of Mathematical Analysis and Applications, 419 (2) (2014), 1261-1273.
5. G.K. Kumar, S.H. Kulkarni, Linear maps preserving pseudospectrum and conditionspectrum, Banach Journal of Mathematical Analysis, 6(1) (2012), 45-60.



ICOMAA 2023

On basicity of the system of eigenfunctions for a discontinuous spectral problem in variable exponent Lebesgue space

Fatih Sirin¹

¹*Department of Mathematics, Halic University,
fatihSirin@halic.edu.tr*

Abstract

In this study, the basicity properties of the eigenfunctions system of a second order differential operator containing spectral parameters under boundary conditions in variable exponent Lebesgue space are investigated. For this purpose, we apply the theory of close bases. In this method we use the classical systems such as exponential and trigonometric systems to obtain the basicity properties of the system.

Keywords: variable exponent space, theory of close base, basicity, second-order discontinuous spectral problem.

References:

1. Zeren, Yusuf, MIQDAD ISMAILOV, and FATİH ŞİRİN. "On basicity of the system of eigenfunctions of one discontinuous spectral problem for second order differential equation for grand-Lebesgue space." *Turkish Journal of Mathematics* 44.5 (2020): 1595-1611. doi:10.3906/mat-2003-20.
2. Zeren, Yusuf, Migdad Ismailov, and Fatih Sirin. "On the basis properties of a system of eigenfunctions of a spectral problem for a second-order discontinuous differential operator in the weighted grand-Lebesgue space with a general weight." *Filomat* 36.17 (2022): 6037-6050.
3. Bilalov, Bilal T., and Zafar G. Guseynov. "Basicity of a system of exponents with a piecewise linear phase in variable spaces." *Mediterranean journal of mathematics* 9 (2012): 487-498..
4. Bilalov, Bilal T., Ali A. Huseynli, and Miran I. Aleskerov. "On the basicity of unitary system of exponents in the variable exponent Lebesgue spaces." (2017)..

ICOMAA 2023

Constructing Lyapunov Functions for Delay ReactionDiffusion Systems and Application

Fatiha Najm¹, Radouane Yafia¹, M. A. Aziz Alaoui²

¹Department of Mathematics, Ibn Tofail University, Kénitra Morocco

fatiha.najm@uit.ac.ma

radouane.yafia@uit.ac.ma

²Normandie Univ, France; ULH, LMAH, F-76600 Le Havre

aziz.alaoui@univ-lehavre.fr

Abstract

Motivated by some biological and ecological problems given by reaction diffusion system with delays and boundary conditions of Neumann type; knowing their associated Lyapunov functions for delay ordinary differential equations, we review a method for determining their Lyapunov functions to establish the local/global stability. The method is essentially based on adding integral terms to the corresponding Lyapunov function for ordinary differential equations. The new approach is not general but it is applicable in a wide variety of delays reaction diffusion models with one discrete delay or more, distributed delay and combination of both of them. To illustrate our result, we give an application to a reaction diffusion epidemiological model with time delay (latency period) and indirect transmission effect.

Keywords: Reaction-diffusion system with delay, Lyapunov function, epidemiological model, latency period, basic reproduction number R_0 .

References:

1. McCluskey C. C. Using Lyapunov Functions to Construct Lyapunov Functionals for Delay Differential Equations. *SIAM J. Appl. Dyn. Syst.* 14(1), 1–24, (2015).
2. Huang G., Ma W, Takeuchi Y. Global properties for virus dynamics model with Beddington–DeAngelis functional response. *Appl. Math. Lett.* 22, 1690–1693, (2009)
3. Hsu S. B. Constructing Lyapunov Functions for Mathematical Models in Population Biology. *Taiwanese Journal of Math.* 9, 151-173, (2005).
4. Korobeinikov A. Lyapunov functions and global properties for SEIR and SEIS epidemic models. *Math. Med. Biol.* 21,2175-83, (2004).

Solution Of Composite Beam With Periodical Curvature By Finite Difference Method Under Uniform Loadings

Fatma Saydam Akman¹, Zafer Kütüğü², Onur Çınar³, Özlem Vardar⁴ and Yousaf Anwar⁵

^{1,3,4}Graduate School of Science and Engineering, Yıldız Technical University

fatmaa.saydam@gmail.com

onurcinar98@gmail.com

ozlemvardar@gmail.com

^{2,5}Civil Engineering Faculty, Yıldız Technical University

kutug@yildiz.edu.tr

yousafanwarkhan@yahoo.com

Abstract

In this study, deflection equations of the simply and/or fixed supported, composite plate-strip (beam) having periodical curved material in structure has been obtained in the framework of Akbarov and Guz' continuum theory by using Kirchhoff-Love plate theory. Computer codes prepared in the programming language Matlab, used to get the numerical results. Deflection equation of the above plate-strip at a point x_i has been defined for different curved values. Finite-difference method has been used in the solution of the deflection equations of the plate-strip, subjected to different distributed load. In numerical analysis, finite-difference methods (FDM) area class of numerical techniques for solving differential equations by approximating derivatives with finite differences. The effects on the deflection value have been studied, by changing the value of the elasticity modulus ratio of the plate-strip laminates and the amplitude, wavelength and phase angle of the periodic curvature.

Keywords: composite, plate-strip, Finite difference method, Kirchhoff-Love plate theory

References:

1. Kütüğü, Z., Natural Frequency and Stability Of Beam and Plate Of Which Composite Material Having Local and Periodical Curvature (1998).
2. Akbarov, S.D., On the determination of normalized nonlinear mechanical properties of composite materials with periodically curved layers, Int. J. Solids and Structures. (1995).
3. Christensen, R.M., Mechanics of composite materials, John Wiley and Sons, Inc. (1979).

ICOMAA 2023

Machine Learning Models in the Diagnosis of Alzheimer Disease

Fatma Nur Budak¹, Murat Sari², Huseyin Tunc³

¹Department of Mathematics, Yildiz Technical University,

²Department of Mathematics Engineering, Istanbul Technical University

³Department of Biostatistics and Medical Informatics, Bahcesehir University

nur.budak@std.yildiz.edu.tr

muratsari@itu.edu.tr

huseyin.tunc@med.bau.edu.tr

Abstract

Dementia has become a major global problem in the 21st century, with Alzheimer's disease (AD) the most common type of dementia. Due to the increasing elderly population, there are difficulties in meeting the demand for diagnosis due to the lack of specialists, especially in many middle and low-income countries. New methods are needed in the diagnosis of individuals with cognitive impairment due to various reasons. In this study, we present some robust machine learning models for diagnosing normal cognitive function, non-AD cognitive disorders and individuals with AD. We use different classification algorithms such as LGBM, Random Forest, MLP, CatBoost, and evaluate the performance of our framework using performance metrics. We show that our machine learning models provide reliable predictions for AD diagnosis.

Keywords: Alzheimer disease, Machine learning, Demantia, Computational neuroscience, Artificial intelligence.

References:

1. An, N., Ding, H., Yang, J., Au, R., & Ang, T. F. (2020). Deep ensemble learning for Alzheimer's disease classification. *Journal of biomedical informatics*, 105, 103411.
2. Ballard, C., Gauthier, S., Corbett, A., Brayne, C., Aarsland, D., & Jones, E. (2011). Alzheimer's disease. *the Lancet*, 377(9770), 1019-1031.
3. Qiu, S., Miller, M. I., Joshi, P. S., Lee, J. C., Xue, C., Ni, Y., ... & Kolachalama, V. B. (2022). Multimodal deep learning for Alzheimer's disease dementia assessment. *Nature communications*, 13(1), 3404.
4. Qiu, S., Joshi, P. S., Miller, M. I., Xue, C., Zhou, X., Karjadi, C., ... & Kolachalama, V. B. (2020). Development and validation of an interpretable deep learning framework for Alzheimer's disease classification. *Brain*, 143(6), 1920-1933.

ICOMAA 2023

A New Function Type in Grill Minimal Spaces and Its Properties

Ferit Yalaz

Department of Mathematics, Selcuk University
ferit.yalaz@selcuk.edu.tr

Abstract

In this work, a new function is defined on the power set of a non-empty set by using concepts such as minimal structure, grill and minimal semi-closure. According to the relation of being a subset, this new function is compared with the function shown with $(\cdot)^{*M}$ in the literature. Moreover, the properties of this function are examined.

Keywords: Minimal structure, Minimal space, Minimal semi-closure, Grill, Grill minimal space.

References:

1. Choquet, G., Theorie des ensembles-sur les notions de filtre et de grille, Comptes rendus hebdomadaires des seances de l'academie des sciences, 224(3) (1947), 171-173.
2. Maki H. Every topological space is pre- $T_{1/2}$. Mem. Fac. Sci. Kochi Univ. Ser. A Math. 17 (1996) 33-42.
3. MinW. K., m-Semiopen sets and m-semicontinuous functions on spaces with minimal structures, Honam Mathematical J. 31(2) (2009) 239-245.
4. Modak, S., Minimal spaces with a mathematical structure, Journal of the Association of Arab Universities for Basic and Applied Sciences 22 (2017), 98-101.
5. Popa V, Noiri T., On the definitions of some generalized forms of continuity under minimal conditions. Mem. Fac. Sci. Kochi Univ. Ser. A Math. (22) (2001) 9-18.



ICOMAA 2023

Approximative Properties of Perturbed Exponential System in Orlicz Spaces

Bilal Bilalov^{1,2}, Yusuf Zeren¹, Fidan Alizadeh², Feyza Elif Dal¹

¹Department of Mathematics, Yildiz Technical University,

²Institute of Mathematics and Mechanics of NAS of Azerbaijan, Baku, Azerbaijan

bilal.bilalov@yildiz.edu.tr

yzeren@yildiz.edu.tr

fidanalizade95@mail.ru

feyzadal@hotmail.com

Abstract

The non-homogeneous Riemann boundary value problem with piecewise Hölderian coefficient is considered in Hardy-Orlicz classes. Based on N-function we introduce new characteristic of Orlicz space and establish its relationship between Boyd indices of considered Orlicz space.

Keywords: Orlicz space, Hardy-Orlicz classes, approximative properties.

Acknowledgements: This work is supported by the Azerbaijan Science Foundation-Grant No: AEF-MQM-QA-1-2021-4(41)-8/02/1-M-02. The research of Feyza Elif Dal is supported by TÜBİTAK 2211-E Domestic Direct Doctorate Scholarship Program.

References:

1. Y. Zeren, F.A. Alizadeh and F.E. Dal On solvability of homogeneous Riemann boundary value problems in Hardy-Orlicz classes, Turk. J. of Math. (2023) (Accepted).
2. B.T. Bilalov, T.B. Gasymov, A.A. Guliyeva, On solvability of Riemann boundary value problem in Morrey-Hardy classes, Turk. J. of Math. 40(50) (2016), 1085-1101.
3. B.T. Bilalov, The basis property of a perturbed system of exponentials in Morrey-type spaces, Sib. Math. Journ. 60(2)(2019), 323-350.
4. B.T. Bilalov, F.A. Alizade, M.F. Rasulov, On bases of trigonometric systems in Hardy-Orlicz spaces and Riesz theorem, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. Mathematics, 39 (4), (2019), 1-1

ICOMAA 2023

Almost Shortest Paths: To Alternate Routes for Security Reasons in Graphs

Gökçe Çaylak Kayaturan

³Department of Mathematics, Ordu University,
gkccaylak@gmail.com

Abstract

This study discusses how form of a path in a graph representing a computer network affects the message delivery during distribution. In order to reduce the delivery time and other costs, shortest paths are widely used for message transfer by the computer network users. However, this sometimes can cause traffic in the network. Additionally, the users also consider other routing issues such as delivery time, security etc. In this case, the users search for alternative routes that do not waste the sources as well as the shortest ones in the network. In this study, we present a secure and fast way to choose the appropriate routes among alternatives. The computer network is represented by a mathematical graph and we label the paths with binary strings by considering structural properties of the graph. We support the path-coding method with mathematical proofs and theorems.

Keywords: Graph, computer network, path, message delivery, routing.

References:

1. Bloom, B. H. (1970). Space/time trade-offs in hash coding with allowable errors. *Communications of the ACM*, 13(7), 422-426.
2. L. Carrea, A. Vernitski, and M. Reed, "Optimized hash for network path encoding with minimized false positives," *Computer networks*, vol. 58, pp. 180–191, 2014.
3. Kayaturan, G. Ç., Özçelik, G., & Gökçe, A. (2023). Decision-aided evaluation of paths for routing on multi-attribute computer network consisting of encoded paths under uncertainty. *Expert Systems with Applications*, 119881.

ICOMAA 2023

Upper bounds for blow up of solutions for a fourth-order parabolic equation with viscoelastic term

Gülistan Butakin¹, Erhan Pişkin²

¹*Department of Mathematics, Dicle University*

²*Institute of Natural and Applied Sciences, Dicle University*

gulistanbutakin@gmail.com

episkin@dicle.edu.tr

Abstract

In this paper, we study a fourth-order parabolic type equation with viscoelastic term. This type problem occurs in many mathematical models in engineering and physical sciences, such as nuclear science, chemical reactions, heat transfer, population dynamics. We prove the upper bounds for blow up of solutions.

Keywords: Blow up, parabolic equation, viscoelastic term.

References:

1. E. Pişkin, B. Okutmuşur, An Introduction to Sobolev Spaces, Bentham Science, 2021.
2. E. Pişkin, G. Butakin, Existence and Decay of solutions for a parabolic-type Kirchhoff equation with variable exponents, Journal of Mathematical Sciences and Modelling, (in press).
3. R.Abita, Bounds for Blow-up Solutions of a Semilinear Pseudo-Parabolic Equation with a Memory Term and Logarithmic Nonlinearity in Variable Space, Mathematica Scandinavica, (in press).

ICOMAA 2023

Blow up solutions for a fourth-order parabolic equation with variable exponents

Gülstan Butakın¹, Erhan Pişkin²

¹*Department of Mathematics, Dicle University*

²*Institute of Natural and Applied Sciences, Dicle University*

gulistanbutakin@gmail.com

episkin@dicle.edu.tr

Abstract

In this presentation, we consider a fourth-order parabolic type equation with a variable exponents. This type problem occurs in many mathematical models of applied science, such as electrorheological fluids, population dynamics, heat transfer, chemical reactions. We prove the blow up of solutions.

Keywords: Blow up, parabolic equation, variable exponents.

References:

1. E. Pişkin, B. Okutmuşur, An Introduction to Sobolev Spaces, Bentham Science, 2021.
2. E. Pişkin, G. Butakın, Existence and Decay of solutions for a parabolic-type Kirchhoff equation with variable exponents, Journal of Mathematical Sciences and Modelling, (in press).
3. X. Wu, B. Guo, W. Gao, [Blow-up of solutions for a semilinear parabolic equation involving variable source and positive initial energy](#), Applied Mathematics Letters, 26 (2013) 539-543.

ICOMAA 2023

Investigation of Hadamard Fractional Differential Inclusions

HabibDjourdem¹

¹ *Department of Mathematics, Relizane University, Algeria*
djourdem.habib7@gmail.com

Abstract

This paper investigates the existence of solutions for hybrid fractional differential inclusions involving Hadamard fractional derivative together with hybrid Hadamard nonlocal integral boundary conditions. By using a fixed point theorem due to Dhage, we establish our main results.

Keywords: Hybrid differential inclusions, Fixed point, Multivalued map, Caratheodory.

References:

1. I. Podlubny, Fractional Differential Equations (Academic Press, San Diego, 1999).
2. A.A. Kilbas, H.M. Srivastava and J.J. Trujillo, Theory and Applications of Fractional Differential Equations, North-Holland Mathematics Studies, 204 (Elsevier Science B.V., Amsterdam, 2006).
3. H. Djourdem, S. Benaïcha, Positive solutions for fractional differential equations with non-separated type nonlocal multi-point and multi-term integral boundary conditions, Stud. Univ. Babeş-Bolyai Math. 66(4)(2021), 691-708.
4. B. C. Dhage, V. Lakshmikantham, Basic results on hybrid differential equation, Nonlinear Anal. Hybrid Syst. 4 (2010), 414-424.



ICOMAA 2023

Neutrosophic Statistical Analysis of Temperatures of Cities in the Southeastern Anatolia Region of Turkey

Hacer Şengül Kandemir¹, Nazlım Deniz Aral², Murat Karakaş³, Mikail Et⁴

¹Faculty of Education, Harran University,
hacer.sengul@hotmail.com

^{2,3}Department of Mathematics, Bitlis Eren University,
ndaral@beu.edu.tr
mkarakas@beu.edu.tr

⁴Department of Mathematics, Fırat University,
mikail68@gmail.com

Abstract

Turkey is one of the countries that will be most affected by a possible climate change within its complex climate structure, especially due to global warming. It is naturally surrounded by seas on three sides and has a faulty topography. Different regions of Turkey will be affected by climate change differently and to varying extents due to its orographic characteristics. In recent years, many heat strokes have been recorded, causing many problems in the environment. Animals die because of water due to environmental change. Statistical methods are widely applied for estimation and estimation of temperature.

In this work, neutrosophic statistical analysis of temperature data of different cities in the southeastern Anatolia region of Turkey is given. The neutrosophic mean and neutrosophic coefficient of variation are computed using the temperature data. In addition, the neutrosophic results are compared with results under classical statistics.

Keywords: Neutrosophic sets, Neutrosophic statistics, temperature, indeterminacy.

Acknowledgements: This projected work was partially (not financial) supported by Harran University with the project HUBAP ID: 22261.

References:

1. Afzal, U., Alrweili, H., Ahamd, N., Aslam, M. Neutrosophic statistical analysis of resistance depending on the temperature variance of conducting material. Scientific reports, Vol. 11, No. 1 (2021) 1-6.
2. Aslam, M. A new method to analyze rock joint roughness coefficient based on neutrosophic statistics. Measurement. Vol. 146 (2019) 65-71.
3. Chen, J., Ye, J., Du, S. Scale effect and anisotropy analyzed for neutrosophic numbers of rock joint roughness coefficient based on neutrosophic statistics. Symmetry. Vol. 9, No. 10 (2017) p.208.
4. Shahzadi, I. "Neutrosophic Statistical Analysis of Temperature of Different Cities of Pakistan. Neutrosophic Sets and Systems. 53(1) (2023) 156-164.

Semi Global Domination Number in Product Fuzzy Graphs

Haifa Ahmed¹, Mahioub Shubatah², Mohammed Alsharafi³

¹Department of Mathematics, Aden University, Yemen

²Department of Mathematics, Sheba Region University, Yemen

³Department of Mathematics, Yildiz Technical University, Turkey

haifaahmed010@gmail.com

mahioub70@yahoo.com

alsharafi205010@gmail.com

Abstract

In this paper we introduced the concepts of semi global domination number in product fuzzy graph and is denoted by $\gamma_{sg}(G)$ and semi complimentary product fuzzy graph. We determine the semi global domination number $\gamma_{sg}(G)$ for several classes of product fuzzy graph and obtain Nordhaus-Gaddum type results for this parameter. Further, some bounds of $\gamma_{sg}(G)$ are investigated. Moreover, the relations between $\gamma_{sg}(G)$ and other known parameter in Product fuzzy graphs are investigated.

Keywords: Product fuzzy graphs, semi complimentary product fuzzy graph, semi global domination number in product fuzzy graph.

References:

1. M. Q Shubatah and H. Ahmed. The Global Domination Number in Product Fuzzy Graphs, EPH, International Journal of Mathematics and statistics, (2020), pp. 60-76.
2. Q.M. Mahioub. Domination in product fuzzy graph, ACMA, 1(2), (2012), pp. 119-125.
3. A. Nagoor Gani, S. Yahya Mohammed and R. Jahir Husain. The semi Global Dominating set in Fuzzy Graphs, Intern. J Fuzzy Mathematical Archive, 4(2), (2014), pp. 104-111.
4. Ramaswmy. Product fuzzy graph, Int. Jon. of Com. Sci. and Net.Sec, 9(1), (2009), pp. 114-118.
5. E. Sampathkumar. The global Domination Number of A Graph, Jour. Math. Phy. Sci. 23(5), (1989), pp. 377-385.

ICOMAA 2023

Trace formula for differential equation with coefficient containing first degree polynomial depending on spectral parameter

Hajar Movsumova

Institute of Mathematics and Mechanics of NAS of Azerbaijan. Baku, Azerbaijan.

movsumovahecer@gmail.com

Abstract

In the Hilbert space $L_2(H, (0,1))$ (where H is separable Hilbert space) we consider the boundary value problem for differential equation with unbounded operator coefficient containing first degree polynomial depending on spectral parameter:

$$l[y] \equiv -y''(t) + Ay(t) + q(t)y(t) = \lambda y(t), \quad y'(0) = 0, \quad y(1)(1 + \lambda) = y'(1)(1 + h + \lambda), \quad h \in \mathbb{R}.$$

A is a self-adjoint positive-definite operator in H and $A^{-1} \in \sigma_\infty$. $\gamma_1 \leq \gamma_2 \leq \dots$ and $\varphi_1, \varphi_2, \dots$, are eigenvalues and eigenvectors of A . $q(t)$ is a self-adjoint operator-valued function, suppose that $q(t)$ is weakly measurable and the following conditions are satisfied: $\|q(t)\| \leq C, t \in [0,1]$;

$$q'(0) = q'(1) = 0; \quad \text{for each } t \quad \sum_{k=1}^{\infty} |(q(t)\varphi_k, \varphi_k)| < \text{const}; \quad \int_0^1 (q(t)\varphi_k, \varphi_k) dt = 0, \quad k = \overline{1, \infty}.$$

Lemma. If eigenvalues of A $\gamma_k \sim gk^\alpha, k \rightarrow \infty, g > 0, \alpha > 0$, then for the eigenvalues of L_0 we can write $\lambda_n(L_0) \sim dn^{\frac{2\alpha}{\alpha+2}}, d = \text{const}$

Theorem. If the operator function $q(t)$ satisfy above conditions. Then under the conditions of lemma, we get the following regularized trace formula

$$\lim_{m \rightarrow \infty} \sum_{n=1}^m (\lambda_n - \mu_n) = \frac{\text{tr } q(0) + \text{tr } q(1)}{4}.$$

Keywords: Hilbert space, eigenvalue parameter, the first regularized trace.

References:

1. Gorbachuk V.I., Gorbachuk M.L. On some class of one boundary value problems for Sturm-Liouville equation with operator coefficient. Ukrainian Journal of Math., 24 (3) (1972) 291-306.
2. Maksudov F.G., Bayramogly M., Adigezalov A.A. On regularized trace of Sturm-Liouville operator on finite segment with unbounded operator coefficient. DAN SSSR. 277(4) (1984) 795-799
3. Fulton, ChT: Two-point boundary value problems with eigenvalue parameter contained in the boundary condition. Proc R Soc Edinburgh., 77A (1977) 293-308.
4. Nigar M., Aslanova, M., Bayramoglu, Khalig M., Aslanov. On one class eigenvalue problem with eigenvalue parameter in boundary condition at one end point. Filomat 32(19) (2018) 6667-6674.

Some Improvement of Berezin Radius inequalities via Specht's Ratio

Hamdullah Başaran¹, Mehmet Gürdal²
^{1,2}Department of Mathematics, Süleyman University,
07hamdullahbasaran@gmail.com
gurdalmehmet@sdu.edu.tr

Abstract

The Berezin symbol \tilde{X} and the Berezin number of an operator X on the reproducing kernel Hilbert space $\mathcal{H}(\Lambda)$ over some set Λ with the reproducing kernel are defined, respectively, by

$$\tilde{X}(\varrho) = \langle X \frac{k_\varrho}{\|k_\varrho\|}, \frac{k_\varrho}{\|k_\varrho\|} \rangle, \varrho \in \Lambda \text{ and } \text{ber}(X) = \sup_{\varrho \in \Lambda} |\tilde{X}(\varrho)|$$

By using this bounded function \tilde{X} , we discover a few inequalities pertaining to Berezin number inequalities of functional Hilbert space operators. There are also some conclusions drawn using Hermite-Hadamard inequality. We strengthen and broaden a few inequalities in relation to Specht's ratio. With these enhancements, we also demonstrate a number of new inequalities for the Berezin norm and Berezin radius of operators.

Keywords: Berezin number, Hermite Hadamard inequality, Specht's ratio, positive operator.

References:

1. F.A. Berezin, Covariant and contravariant symbols for operators, Math. USSR-Izvestiya, 6 (1972), 1117-1151.
2. V. Gürdal, H. Başaran, Berezin radius inequalities via geometric convexity, Palestine J. Math., 12(1) (2023), 240-253.
3. M. Gürdal, H. Başaran, Berezin number inequalities in terms of Specht's ratio, El-Cezeri, 9(4) (2022), 1201-1214.
4. M.T. Karaev, Berezin symbol and invertibility of operators on the functional Hilbert spaces, J.Funct. Anal., 238 (2006), 181-192.
5. W. Specht, Zur theorie der elementaren Mittel, Math. Z., 74 (1960), 91-98.

ICOMAA 2023

Exponential, Logarithmic and Trigonometric Functions on Split Quaternions and Their Characterizations

Hamide Feyza Aykut¹, Salim Yüce²

^{1,2}Department of Mathematics, Yıldız Technical University,
hamidefeyzaaykut@gmail.com
sayuce@yildiz.edu.tr

Abstract

In the literature, there are two definitions of the set of split quaternions that is defined by the metrics $(-, -, +, +)$ or $(+, +, -, -)$. Polar forms and exponential functions have already been defined over the set of split quaternions with the metric $(-, -, +, +)$.

In this study, we considered the set of split quaternions with the metric $(+, +, -, -)$. It is clear that the set of the vector part of them is isomorphic to \mathbb{R}_2^3 with the metric $(+, +, -)$. We discussed the properties and characterizations of the exponential function on split quaternions considering definition of the exponential function via this approach. Additionally, we defined logarithmic and trigonometric functions on split quaternions and discussed their properties and characterizations.

Keywords: Split quaternions, exponential function, logarithmic function, trigonometric function.

References:

1. Özdemir, M., Ergin, A. (2005). Some geometric applications of timelike quaternions. 10.13140/2.1.2870.5609.
2. Kula, L., Yaylı, Y. (2007). Split quaternions and rotations in semi Euclidean space E^4_2 . Journal of The Korean Mathematical Society - J Korean Math Soc. 44. 1313-1327. 10.4134/JKMS.2007.44.6.1313.
3. Alagöz, Y., Oral, K., Yüce, S. (2012). Split quaternion matrices. Miskolc Mathematical Notes. 2. 10.18514/MMN.2012.364.
4. Erdoğan, M., Özdemir, M., (2017). On exponential of split quaternionic matrices. Applied Mathematics and Computation, vol.315, 468-476.
5. Öztürk, İ., Özdemir, M. (2022). On geometric interpretations of split quaternions. Mathematical Methods in the Applied Sciences. 46. 10.1002/mma.8518.

Some Combinatorial Results for the Hyperoctahedral Group

Hasan Arslan

Department of Mathematics, Faculty of Science, Erciyes University,
hasanarslan@erciyes.edu.tr

Abstract

In this work, we will present that the flag major index $fmaj$ defined in [1] on the hyperoctahedral group is equidistributed over the group with the inversion statistic introduced in [3] with respect to the root system of this real reflection group. In addition, we will combinatorially show that $Dmaj$ permutation statistic defined in [2] on the group of even-signed permutations and the inversion statistic we constructed depending on the root system of the group are equidistributed over this group.

Keywords: Hyperoctahedral group, permutation statistic, root system.

References:

1. R.M. Adin, F. Brenti, Y. Roichman, Descent numbers and major indices for the hyperoctahedral group, *Adv. Appl. Math.* 27 (2001) 210–224.
2. R. Biagioli, Signed Mahonian polynomials for classical Weyl groups, *European J. Combin.* 27 (2006), 207–217.
3. I. V. Raharinirina, On hyperoctahedral enumeration system, application to signed permutations, *Asian Research Journal of Mathematics*, 16(8) (2020) 40–49.



ICOMAA 2023

A Relation between Bosbach State Operator and Very True Operator

Ibrahim Senturk

Department of Mathematics, Ege University,
ibrahim.senturk@ege.edu.tr


Abstract

In this study, the concepts of Bosbach state operator and Very True operator in many-valued algebraic structures corresponding to many-valued logic are discussed and the relations between these concepts are revealed. To support mentioned concepts, we give place to some interesting examples.

Keywords: Bosbach state operator, Very True operator, many-valued algebra, many-valued logic.

References:

1. T. Oner and I. Senturk, The Sheffer stroke reducts of basic algebras, *Open Mathematics* 15(1) (2017) 926 – 935.
2. I. Senturk, A view on state operators in Sheffer stroke basic algebras, *Soft Comput.* 25 11471–11484.
3. I. Senturk, A bridge Construction from Sheffer stroke basic algebras to MTL-algebras, *BAUN Fen. Bil. Enst. Dergisi* 22(1) (2020)193–203.
4. I. Senturk, Riečan and Bosbach state operators on Sheffer stroke MTL-algebras, *Bull. Int. Math. Virtual Inst.* 12(1) (2022) 181–193.
5. L. C. Ciungu, Internal states on equality algebras, *Soft Comput.* 19(4) (2015) 939–953.



ICOMAA 2023

Some Existence Theorems for Solutions of Set-Valued Optimization Problems with Respect to Various Set Order Relations

İlknur Atasever Güvenç¹

^{1,2,3}Department of Mathematics, Eskişehir Technical University,
iatasever@eskisehir.edu.tr

Abstract

In this work, relations among existence of solutions of a set-valued optimization problem with respect to vector and set optimization criteria are given. For set optimization criterion set order relations $\leq_c, \leq_{mc}, \leq_{mn}, \leq_m$ given by Jahn and Ha [1], \leq_s given by Nishnianidze [2] and Young [3] are considered. Also, under some cone compactness assumptions existence theorems for solutions of a set optimization problem with respect to these order relations are presented.

Keywords: set-valued optimization, set optimization criterion, vector criterion, set order

References:

1. J. Jahn, T.X.D. Ha, New order relations in set optimization, *J. Optimiz. Theory App.* 148 (2003) 209–236.
2. Z.G. Nishnianidze, Fixed points of monotonic multiple-valued operators, *Bull Georgian Acad. Sci.* 114 (1984) 489–491.
3. R.C. Young, The algebra of many-valued quantities, *Math. Ann.* 104 (1931) 260–290.
4. E. Hernández, L. Rodríguez-Marín and M. Sama, On solutions of set-valued optimization problems, *Computers and Mathematics with Applications*, 60 (2010) 1401-1408.
5. D. Kuroiwa, Existence theorems of set optimization with set-valued maps, *J. Inf. Optim.* 24 (2003) 73–84.

ICOMAA 2023

Revenue Maximization For Polynomial Demand Function Using Interval-Valued Trapezoidal Fuzzy Numbers

Ismail TEMBELO¹, Salih AYTAR²

^{1,2}Department of Mathematics, Suleyman Demirel University,
salihaytar@sdu.edu.tr,
ismailtembelo20@gmail.com

Abstract

In this talk, we first consider the linear demand function

$$p = a - bx$$

where $a, b > 0$ and p is the unit price with respect to the demand quantity x . Then we use the parabolic demand function

$$p = a - bx - cx^2,$$

where $a, b, c > 0$. After the coefficients a, b and c are fuzzified by interval-valued fuzzy numbers, we calculate maximum revenue.

KeyWords: Demand function; Revenue function; Trapezoidal fuzzy number; Interval-valued trapezoidal fuzzy number; Graded mean defuzzification method.

References:

1. Chang S.-C., Yao J.-S., Optimal revenue for demand function in fuzzy sense, *Fuzzy Sets and Systems*, 111(2000) 419-431.
2. Chang S.-C., Optimal revenue for fuzzy demand quantity, *Fuzzy Sets and Systems* 111(2000) 399-417.
3. Chiao, K. P. 2011 Multiple criteria group decision making with triangular interval type-2 fuzzy sets, *In 2011 IEEE International Conference on Fuzzy Systems*, 2575-2582.

ICOMAA 2023

Minkowski difference and sum of half planes

Jalolxon Nuritdinov

*Department of digital technologies and mathematics, Kokand University,
nuritdinovjt@gmail.com*

Abstract

A straight line given in a plane divides it into two convex areas. We call one of these spheres a half-plane. One can think about the Minkowski difference and sum of such half-planes. Works [1], [3] provide information about the Minkowski sum and Minkowski difference of straight lines. Using these, we will consider the problem of finding the Minkowski sum and difference of half-planes.

In this article, the conditions for the Minkowski difference and sum of two half-planes to be non-empty are derived and proved in the form of a theorem. Also, the method of calculating the given two half-planes in the presence of the Minkowski difference is described. In writing this article, we widely used works [2], [4], [5]. Continuing the results presented in this paper, these results can be used to calculate the Minkowski difference of other unbounded sets.

Keywords: Minkowski difference, Minkowski sum, half plane, a straight line, unbounded set.

References:

1. D. Velichova, Notes on properties and applications of Minkowski point set operations, South Bohemia Mathematical Letters. 2016. Volume 24. □1. pp.57-71.
2. M.Mamatov, J.Nuritdinov, Some properties of the sum and geometric differences of Minkowski, Journal of Applied Mathematics and Physics, 2020, 8, 2241-2255.
3. J.Nuritdinov, On the Minkowski difference of lines and planes, Modern problems of applied mathematics and information technologies. Al-Khwarizmi 15-17 November, 2021, Fergana, Uzbekistan. pp 252.
4. J.Nuritdinov, About the Minkowski difference of squares on a plane, Differential Geometry - Dynamical Systems, Vol.24, 2022, pp. 151-156. © Balkan Society of Geometers, Geometry Balkan Press 2022.
5. J. T. Nuritdinov, On Minkowski difference of triangles, Bull. Inst. Math., 2021, Vol.4, №6, pp. 50-57.

ICOMAA 2023

On the completeness and minimality in the space $L_p(0, \pi)$ of a weighted system of sines with a power-law weight with an excluded arbitrary element

Yusuf ZEREN¹, Migdad I. ISMAILOV², Kader ŞİMŞİR ACAR³

¹Department of Mathematics, Yıldız Technical University,
yzeren@yildiz.edu.tr

²Department of Mathematics, Baku State University
migdad-ismailov@rambler.ru

³Graduate School of Natural and Applied Sciences, İstanbul Commerce University
ksacar@ticaret.edu.tr

Abstract

The paper considers the issue of completeness and minimality of a weighted exponential system in the grand Lebesgue space. It follows from the results of [1] that a weighted exponential system does not form a basis in the grand Lebesgue space. Also in [1], sufficient conditions for the completeness and minimality of the weighted exponential system were found. In this paper, we give criteria for the completeness and minimality of the weighted exponential system in grand Lebesgue space. Note that in the case of a weight function of a general form, the criteria for completeness and minimality of weight systems of exponents and systems of cosines with an excluded arbitrary element were studied in [2, 3].

Keywords: completeness, minimality, Lebesgue space, weighted system, sines

References:

1. B. T. Bilalov and F. Guliyeva, "On the frame properties of degenerate system of sines", J. Funct. Space Appl., 2012, Art. ID 184186, pp. 1-12.
2. A. Sh. Shukurov, "Basic properties of weighted exponential systems with excess", Bulletin of Samara University. Vestn. SamGU. Natural Science Ser., 2018, Issue 24(1), 14-19.
3. A. Sh. Shukurov, "On basicity of the degenerate trigonometric system with excess", Acta Comment. Univ. Tartu. Math. 21 (2017), no.2, 249–257.

ICOMAA 2023

A Novel Alternative Algorithm to Find All Multiple Solutions of General Integer Linear Programs

Kadriye Şimşek Alan

*Department of Mathematical Engineering, Yıldız Technical University,
ksimsek@yildiz.edu.tr*

Abstract

Integer linear programming (ILP) is often used to model and solve real-life problems. In practice, alternative solutions are very useful as they significantly increase flexibility for the decision maker. In this study, an alternative method based on parameterization obtained from the Diophantine equation is developed to find all alternative solutions of ILP problems and an easy to implement, efficient and reliable algorithm is presented. The proposed method was used without being affected by the number of variables and constraints in the problem. Numerical examples are presented to demonstrate the usefulness of the proposed method. In addition, these examples are coded in the MAPLE programming language according to the proposed algorithm.

Keywords: Linear integer programming, Linear Diophantine equations, Optimal hyperplane, Pure integer programming problems.

References:

1. Schrijver, A. Theory of Linear and Integer Programming, John Wiley & Sons Ltd. (1986).
2. Joseph, A. Parametric formulation of the general integer linear programming problem. Computers & operations research, 22(9) (1995) 883-892.
3. Pandian, P., Jayalakshmi, M. A. New Approach for solving a Class of Pure Integer Linear Programming Problems. Journal of Advanced Engineering Technology, 3(2012) 248-251.
4. Tsai, J. F., Lin, M. H., Hu, Y. C. Finding multiple solutions to general integer linear programs. European Journal of Operational Research, 184(2), (2008) 802-809.

ICOMAA 2023

A Novel Alternative Algorithm for Solving Integer Linear Programming Problems

Kadriye Şimşek Alan

*Department of Mathematical Engineering, Yıldız Technical University,
ksimsek@yildiz.edu.tr*

Abstract

This study proposes an integer linear programming (ILP) an iterative method for solving integer programs with linear objectives and linear constraints. The proposed method is based on the parameterization obtained from the Diophantine equation, and using this parameterization the original problem is reformulated as another ILP problem that can be more efficiently solved using simple mathematical programming. The method is used without being affected by the number of constraints of the problem and also offers all alternative solutions to the decision maker. It is demonstrated by some examples that this method provides an efficient algorithm.

Keywords: Linear integer programming, Linear Diophantine equations, Optimal hyperplane, Pure integer programming problems.

References:

1. Schrijver, A. Theory of Linear and Integer Programming, John Wiley & Sons Ltd. (1986).
2. Genova, K., & Guliashki, V., 2011. Linear integer programming methods and approaches—a survey. Journal of Cybernetics and Information Technologies, 11(1)(2011), 1-23.
3. Joseph, A. Parametric formulation of the general integer linear programming problem. Computers & operations research, 22(9) (1995) 883-892.
4. Pandian, P., Jayalakshmi, M. A. New Approach for solving a Class of Pure Integer Linear Programming Problems. Journal of Advanced Engineering Technology, 3(2012) 248-251.

ICOMAA 2023

Effects of Musics Composed Using Mathematical Methods and DNA On the EEG Frequency Bands of Healthy Individuals

Cemil Karaçam¹, Halil Yakıt², Kayra Ege Altun³, Şerif Efe Dartar⁴

¹Koycegiz Science and Art Center,
Koycegiz, Mugla, Türkiye

cemil-karacam@hotmail.com

^{2,3,4}Kabatas Erkek High School,

Beşiktaş, İstanbul, Türkiye

halilyakit@gmail.com

altunkayra07@gmail.com

dserif.efe@gmail.com

Abstract

Researches about the mathematical techniques that famous musicians used are still being debated by musicians and mathematicians. In this paper, composing new musics using new mathematical techniques or DNA was aimed to remove the lack of literature on this topic. Totally five musics were composed for this project by using DNA, Golden Ratio, and Fibonacci numbers. After the composing process, the EEG tests of the healthy volunteers were taken while they were listening to composed musics. After that, the results were evaluated (by comparing them), and found new techniques for music therapy.

Keywords: Mathematics, Golden Ratio, Fibonacci numbers, Music, DNA, EEG

References:

1. Gena, P., & Strom, C. (1995). Musical synthesis of DNA sequences. XI Colloquio di Informatica Musicale, 203-204.
2. Mutver, S. (2007). Müzikte altın oran. (Yüksek Lisans Tezi). İstanbul Üniversitesi, İstanbul.
3. Yılmaz, B., & Can, Ü. K. (2019). Türkiye'de müzik terapi uygulamalarında kullanılan müzikler. OPUS International Journal of Society Researches, 13(19), 592-620.

New Triangular q -Fibonacci Matrix and Its Applications

Koray İbrahim Atabey¹, Muhammed Çınar², Mikail Et³

¹Muş Nizamülmülk Girl Anatolia İmam Hatip High School Muş,

²Department of Mathematics, Muş Alparslan University,

³Department of Mathematics, Fırat University,

korayatabey7@gmail.com

muhammedcinar23@gmail.com

mikailet68@gmail.com

Abstract

In this study, we construct a new triangular q -analogue of the q -Fibonacci matrix $\tilde{F}_q = (F_{nk}(q))$ defined by

$$F_{nk}(q) = \begin{cases} \frac{q^k F_k(q)}{F_{n+2}(q) - 1}, & 1 \leq k \leq n \\ 0, & \text{otherwise} \end{cases}.$$

Later, we define the sequence spaces $c(\tilde{F}_q)$, $c_0(\tilde{F}_q)$, $\ell_\infty(\tilde{F}_q)$, $\ell_p(\tilde{F}_q)$ ($1 \leq p < \infty$) with this analogue and study some topological properties, and give some inclusion relations for these spaces. In addition, we build a bases for the space $\ell_p(\tilde{F}_q)$, compute α -, β -, γ - duals of the same space, characterize some matrix classes and examine some geometric properties.

Keywords: q -Fibonacci numbers, q -analogue, dual spaces, matrix transform, Banach-Saks property.

References:

1. Aktuğlu, H., Bekar, Ş.: On q -Cesàro matrix and q -statistical convergence. J. Comput. Appl. Math. 235(16), 4717–4723 (2011).
2. Andrews, G.E.: Fibonacci numbers and the Rogers-Ramanujan identities. Fibonacci Quart. 42(1), 3–19 (2004).
3. Atabey, K. İ., Çınar, M. and Et, M.: q -Fibonacci sequence spaces and related matrix transformations. Journal of Applied Mathematics and Computing, (2022) 1-20
4. Aytaç, P.: Some arithmetic properties of q -Fibonacci numbers. Master thesis, Akdeniz University (2018).

ICOMAA 2023

A new approach to the generalization of Hardy-Littlewood integral inequalities on time scale

Lütfi Akın¹, Yusuf Zeren²

¹Department of Business Administration, Mardin Artuklu University, Mardin, Turkey

²Department of Mathematics, Yildiz Technical University, Istanbul, Turkey

lutfiakin@artuklu.edu.tr

yzeren@yildiz.edu.tr

Abstract

Hardy-Littlewood type integral inequalities have an important place in harmonic analysis and time scale. In this study, we will present a new approach to generalize multivariate fractional Hardy-Littlewood type inequalities on time scale calculus.

Keywords: Morrey-type classes, harmonic functions, Statistical Convergence, λ - statistical convergence, Intuitionistic fuzzy 2-normed space.

References:

1. Agarwal, R.P., Bohner, M., O'Regan, D., Saker, S.H. (2011). Some Wirtinger-type inequalities on time scales and their applications, *Pacific J. Math.* **252**, 1–26 (2011)
2. Hilger, S. (1990). Analysis on measure chains-a unified approach to continuous and discrete calculus, *Results Maths.* **18**(1-2) 18–56.
3. Akın L.(2021). On innovations of n-dimensional integral-type inequality on time scales. *Adv. Differ. Equ.* 2021:**148**.
4. Bohner, M., Petereson, A. (2001). *Dynamic Equations on Time Scales. An Introduction with Applications*, Birkhauser, Boston
5. Akın L. (2023). On generalized weighted dynamic inequalities for diamond- α integral on time scales calculus. *Indian J Pure Appl Math.* <https://doi.org/10.1007/s13226-023-00366-6>.

ICOMAA 2023

Several characterizations of the Bessel functions and their applications

Mahvish Ali

Department of Mathematics

Faculty of Science

Shree Guru Gobind Singh Tricentenary University

Gurugram, Haryana, India

mahvishali37@gmail.com

Abstract

The complex form of certain special polynomials are investigated by many authors, see for example [1-3]. Recently, Srivastava and co-authors [3] show that the real and imaginary parts of a general set of complex Appell polynomials can be represented in terms of the Chebyshev polynomials of the first and second kind. After that, Kim [2] introduced the degenerate form of the complex Appell polynomials. The main motive of this paper is to show that the generating function can be employed efficiently to obtain certain results for the special functions. The complex form of the Bessel functions is introduced by means of the generating function. Certain enthralling properties for the complex Bessel functions are investigated in view of generating function method. By considering the real and imaginary parts of the complex Bessel functions separately, we get the cosine-Bessel functions and the sine-Bessel functions. Several novel identities and Jacobi-Anger expansions for the cosine-Bessel functions and sine-Bessel functions are established. Moreover, the generating function of the degenerate Bessel functions is investigated and certain novel identities for these functions are obtained. A hybrid form of these degenerate Bessel functions, namely degenerate Fubini-Bessel functions is constructed using the replacement technique. Finally, the explicit forms of the real and imaginary parts of the complex Bessel functions are established by using hypergeometric approach.

Keywords: Bessel functions; 2-variable Fubini polynomials; complex Bessel functions; degenerate Bessel functions, hypergeometric functions

References:

1. M.E.H. Ismail, Analytic properties of complex Hermite polynomials, *Trans. Am. Math. Soc.* 368 (2016) 1189-1210.
2. D. Kim, A Note on the Degenerate Type of Complex Appell polynomials, *Symmetry*, 11(2019) 1339.
3. H.M. Srivastava, P.E. Ricci, P. Natalini, A family of complex Appell polynomial sets, *Rev. Real Acad. Cienc. Exact. Fis. Nat. Ser. A Mat.*, 113(2019) 2359-2371.

ICOMAA 2023

Pseudo-implication and fuzzy ideals in BL-algebras

Manal A. Alqahtany^{1,2}, Dr Tanisha Baghwan Dale³

1. Department of Mathematics, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad, India.

2. Department of Mathematics, Taiz University, Taiz, Yemen.

3. Department of Mathematics, Rajashri Shahu Arts, Comm. Science College, Aurangabad, India.

manal.abdullah@taiz.edu.ye

daletanu@gmail.com

Abstract

In this work, the concept of pseudo-implication was presented by Yongwei Yang and Xiaolong Xin. They defined this new operation as a binary operation in BL-algebras that is non-abelian and gave another description of ideals in BL-algebras. Our aim is to employ this operation to define the concept of a fuzzy ideal, which serves as a natural extension of the idea of a fuzzy ideal in MV-algebras. By utilizing pseudo-implication, we obtain another simplified description of fuzzy ideals in BL-algebras and provide several characterizations.

Keywords: BL-algebra, fuzzy ideal, fuzzy filter.

References:

1. B.L. Meng, X.L. Xin, On fuzzy ideals of BL-algebras, The Scientific World Journal, 2014, Article ID 757382, 12 pages.
2. C. Lele and J.B. Nganou, Pseudo-addition and fuzzy ideals in BL-algebras. Annals of Fuzzy Mathematics and Informatics. Volume 8, No.2, (August 2014), pp. 193–207.
3. C. Lele and J.B. Nganou, MV-algebras derived from ideals in BL-algebras, Fuzzy Sets, and Systems, 218 (2013), 103–113.
4. Yang, Yongwei & Xin, X., On characterizations of BL-Algebras via implicative ideals. Italian Journal of Pure and Applied Mathematics., 37 (2017), 493-506.
5. Paad Akbar, Folding Theory of Implicative and Obstinate Ideals in Algebras, Discussiones Mathematicae -BL- General Algebra and Applications, vol.38, no.2, 2018, pp.255-271.

ICOMAA 2023

Existence of Solutions to Fuzzy Stochastic Differential Equations

Marek T. Malinowski

Department of Applied Mathematics, Cracow University of Technology, Cracow, Poland
malinowskimarek@poczta.fm

Abstract

Stochastic differential equations are a good tool for modeling dynamic phenomena subjected to uncertainty resulting from the action of random forces. To formulate and apply them, concepts of stochastic integrals are needed. In practice, there is a need to take into account uncertainties that are not of the stochastic type. For example, some human judgments about the parameters of the considered dynamical system may be vague, ambiguous or fuzzy. To model the dynamics of systems operating in random and unclear/fuzzy environments, we propose to use fuzzy stochastic differential equations and their particular version, i.e. set-valued stochastic differential equations. In this communication, we will define how fuzzy stochastic differential equations should be understood and focus on the question of the existence of solutions to such equations.

Keywords: Fuzzy stochastic differential equation, Fuzzy stochastic process, Fuzzy integrals.

References:

1. M.T. Malinowski (2022), Symmetric functional set-valued integral equations and Bihari–LaSalle inequality, *Symmetry* 14(11), 2246.
2. M.T. Malinowski (2021), Bilateral multivalued stochastic integral equations with weakening of the Lipschitz assumption”, *Bulletin des Sciences Mathématiques* 172, 103041.
3. M.T. Malinowski (2020), Symmetric fuzzy stochastic differential equations with generalized global Lipschitz condition, *Symmetry* 12, 819.
4. M.T. Malinowski (2016), Bipartite fuzzy stochastic differential equations with global Lipschitz condition, *Math. Probl. Eng.* vol. 2016, 13 pages.
5. M.T. Malinowski (2015), Stochastic fuzzy differential equations of a nonincreasing type”, *Commun. Nonlinear Sci. Numer. Simulat.* 33, 99-117.

ICOMAA 2023

A unified Feng–Liu type result in relational metric spaces with an application

Md Hasanuzzaman¹ and Mohammad Imdad²

¹*School of Mathematic, Thapar Institute of Engineering and Technology, Patiala Punjab-147004, India*

²*Department of Mathematics, Aligarh Muslim University, Aligarh-202002, India*

md.hasanuzzaman1@gmail.com

mhimdad@gmail.com

Abstract

The purpose of this paper is to obtain an existence Feng–Liu type fixed point result for multi-valued mappings in relational metric spaces which generalizes, extends and unifies several relevant fixedpoint results of the existing literature. Suitable illustrative examples are adopted to substantiate the genuineness of our newly proved result. Besides this, we present a result in a Banach space and utilize the same to show the convergence of successive approximation for the modified nonlinear Bernstein operators in $C[0, 1]$.

Keywords: Fixed point, multi-valued mapping, binary relations, relational metric space, Bernstein operators.

References:

1. Mizoguchi, N., Takahashi, W.: Fixed point theorems for multivalued mappings on complete metric spaces. *J. Math. Anal. Appl.* 141(1), 177–188 (1989).
2. Feng, Y., Liu, S.: Fixed point theorems for multi-valued contractive mappings and multi-valued Caristi type mappings. *J. Math. Anal. Appl.* 317(1), 103–112.
3. Alam, A., Imdad, M.: Relation-theoretic contraction principle. *J. Fixed Point Theory Appl.* 17(4), 693–702 (2015).
4. Kelisky, R., Rivlin, T.: Iterates of Bernstein polynomials. *Pac. J. Math.* 21(3), 511–520 (1967).

ICOMAA 2023

Analytical approximate solutions of some nonlinear evolution equations of fractional order via a novel fractional variational iteration method

Md. Asaduzzaman¹ and Faruk Özger²

¹*Department of Mathematics, Islamic University, Kushtia-7003, Bangladesh,*
asad@math.iu.ac.bd

²*Department of Engineering Sciences, Izmir Katip Celebi University, 35620, Izmir, Türkiye,*
farukozger@gmail.com

Abstract

In this paper, first we formulate and analyze a novel fractional analytical approximation method named as fractional alternative variational iteration method (FAVIM). After formulation of the FAVIM, we consider the initial value problems (IVPs) corresponding to nonlinear time-fractional Fornberg-Whitham equation (NLTFWE), modified nonlinear time-fractional Fornberg-Whitham equation (mNLTFWE), time fractional modified Camassa-Holm equation (TFmCHE), time fractional modified Degasperis-Procesi equation (TFmDPE). Then, we apply our formulated FAVIM for solving the considered IVPs. Here, we use the symbolic software package “Maple” for finding the solution of the considered IVPs and for plotting the 3D and contour figures of the obtained analytical approximate solutions. Finally, we check the accuracy of the obtained solutions by making a numerical comparison with the exact solutions and other existing analogous approximate solutions and which verify that the effectiveness, suitability and time consumed ability of the proposed FAVIM is better.

Keywords: Fractional alternative variational iteration method, Nonlinear evolution equations of fractional order, Caputo’s fractional derivative, Maple software package.

References:

1. Sakar, M.G., Ergoren, H. (2015). Alternative variational iteration method for solving the time-fractional Fornberg–Whitham equation. *Appl. Math. Model.*, 39, 3972-3979.
2. Ibis, B., Bayram, M. (2014). Analytical approximate solution of time-fractional Fornberg–Whitham equation by the fractional variational iteration method. *Alexandria Engineering Journal*, 53, 911-915.
3. Fornberg, B. Whitham, G.B. (1978). A Numerical and Theoretical Study of Certain Nonlinear Wave Phenomena, *Philosophical Transactions of the Royal Society of London. Series A* 289, 373-404.
4. He, B., Meng, Q., Li, S. (2010), Explicit Peakon and Solitary Wave Solutions for the Modified Fornberg-Whitham Equation. *Applied Mathematics and Computation* 217, 1976-1982.
5. Wazwaz, A.M. (2006). Solitary Wave Solutions for Modified Forms of Degasperis-Procesi and Camassa-Holm Equations. *Physics Letter A* 352, 500-504.

Stancu-Type Generalized q -Bernstein–Kantorovich Operators Involving Bezier Bases

Md. Nasiruzzaman

Department of Mathematics, Faculty of Science, University of Tabuk, PO Box-4279, Tabuk-71491, Saudi Arabia,

nasir3489@gmail.com

Abstract

We construct the Stancu-type generalization of q -Bernstein operators involving the idea of Bézier bases depending on the shape parameter $1 \leq \lambda \leq 1$ and obtain auxiliary lemmas. We discuss the local approximation results in term of a Lipschitz-type function based on two parameters and a Lipschitz-type maximal function, as well as other related results for our newly constructed operators. Moreover, we determine the rate of convergence of our operators by means of Peetre's K -functional and corresponding modulus of continuity.

Keywords:

(λ, q) -Bernstein operators; bezier bases; uniform convergence; Lipschitz-type functions; rate of convergence

References:

1. S.N. Bernstein, Demonstration du theoreme de Weierstrass fondee sur le calcul des probabilites, Commun. Kharkov Math. Soc. 13 (1912/13), 1-2
2. Mohiuddine, S.A.; Ozger, F. Approximation of functions by Stancu variant of Bernstein-Kantorovich operators based on shapeparameter a . Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat. RACSAM 2020, 114, 70.
3. Ozger, F. On new Be_zier bases with Schurer polynomials and corresponding results in approximation theory. Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat. 2020, 69, 376–393

ICOMAA 2023

The Marchenko Method for the Derivative NLS Equation

Mehmet Unlu¹, Ramazan Ercan²

¹*Department of Mathematics, Recep Tayyip Erdogan University,*

²*Department of Mathematics and Statistics, Saint Louis University,*

mehmet.unlu@erdogan.edu.tr

ramazan.ercan@slu.edu

Abstract

We present the Marchenko method to solve the derivative NLS (nonlinear Schrödinger) equation. This is done by deriving a linear integral equation, which is our Marchenko integral equation, using as input the time-evolved reflection coefficient and the time-evolved bound-state data expressed in terms of a matrix triplet. In the reflectionless case, we present a solution formula, and we illustrate our method with some explicit examples.

Keywords: Derivative NLS equation, inverse scattering, Marchenko method

References:

1. T. Aktosun, R. Ercan and M. Unlu (2022), The Marchenko method to solve the general system of derivative nonlinear Schrödinger equations, preprint, arXiv:2209.08426.
2. T. Aktosun and R. Ercan (2022), The generalized Marchenko method in the inverse scattering problem for a first-order linear system, preprint, arXiv:2203.02663.
3. H. H. Chen, Y. C. Lee, and C. S. Liu (1979), Integrability of nonlinear Hamiltonian systems by inverse scattering method, Phys. Scr. 20, 490-492.
4. D. J. Kaup and A. C. Newell (1978), An exact solution for a derivative nonlinear Schrödinger equation, J. Math. Phys. 19, 798-801.
5. T. Tsuchida and M. Wadati (1999), New integrable systems of derivative nonlinear Schrödinger equations with multiple components, Phys. Lett. A 257, 53-64.

ICOMAA 2023

A New Note on Summability of Infinite Series and Fourier Series

Hikmet Seyhan Özarslan¹ and Mehmet Öner Şakar²

^{1,2}*Department of Mathematics, Erciyes University,*

seyhan@erciyes.edu.tr

mehmethaydaroner@hotmail.com

Abstract

In this paper, we have generalized two main theorems dealing with absolute Riesz summability factors of infinite series and Fourier series to the $|A, p_n, \beta; \delta|_k$ summability method. Also, some results related to the new theorem are obtained.

Keywords: Absolute matrix summability, Infinite series, Summability factors, Fourier series.

References:

1. H. Bor, On two summability methods, *Math. Proc. Cambridge Philos Soc.* 97(1) (1985) 147-149.
2. H. S. Özarslan and A. Karakaş, A new study on absolute summability factors of infinite series, *Maejo Int. J. Sci. Tech.* 13 (2019) 257-265.
3. H. Bor, A new note on factored infinite series and trigonometric Fourier series, *C. R. Math. Acad Sci.* 359 (2021) 323-328.
4. H. S. Özarslan, A new factor theorem for absolute matrix summability, *Quaest. Math.* 42 (2019) 803-809.
5. H. Bor, On the absolute Riesz summability factors, *Rocky Mt. J. Math.* 24(4) (1994) 1263-1271.

ICOMAA 2023

A NOTE ON CONSTRAINED MULTIVARIATE LINEAR MODEL AND ITS REDUCED MODELS

Melek Eriş Büyükkaya¹, Nesrin Güler²

¹*Department of Statistics and Computer Sciences, Karadeniz Technical University,*

²*Department of Econometrics, Sakarya University,*

melekeris@ktu.edu.tr

nesring@sakarya.edu.tr

Abstract

The fundamental prediction problems relating to a constrained multivariate linear model (CMLM) and its reduced models are covered in this work. We obtain an unconstrained multivariate linear model (UMLM) from the CMLM by using a reparameterization approach. Thus, we present the derivation of analytical formulas for calculating the best linear unbiased predictors (BLUPs) of all unknown parameter matrices by using some quadratic matrix optimization methods after converting the CMLM to the UMLM. We may also refer to [1-5], for the corresponding studies in which similar approaches were conducted.

Keywords: BLUP, constrained multivariate linear model, reparameterization, reduced models.

References:

1. B. Jiang, Y. Tian, On equivalence of predictors/estimators under a multivariate general linear model with augmentation, *J. Korean Stat. Soc.* 46 (4) (2017a) 551–561.
2. B. Jiang, Y. Tian, On additive decompositions of estimators under a multivariate general linear model and its two submodels, *J. Korean Stat. Soc.* 162 (2017b) 193–214.
3. B. Jiang, Y. Tian, Equivalence Analysis of Statistical Inference Results under True and Misspecified Multivariate Linear Models, *Mathematics* 11(1) (2023) 182.
4. Y. Sun, H. Jiang, Y. Tian, A prediction analysis in a constrained multivariate general linear model with future observations, *Commun. Stat. - Theory Methods* 50(2) (2021) 345–357.
5. Y. Tian, C. Wang, On simultaneous prediction in a multivariate general linear model with future observations, *Stat. Probab. Lett.* 128 (2017) 52–59.

ICOMAA 2023

Soliton Solutions of the (3+1)-dimensional Generalized Kadomtsev-Petviashvili Equation

Melih Cinar¹

¹Department of Mathematical Engineering, Yildiz Technical University,
mcinar@yildiz.edu.tr

Abstract

This study investigates soliton solutions of the generalized Kadomtsev-Petviashvili equation in (3+1)-dimension that was recently introduced. The KP equation is used to model shallow water waves and the topic is a vital aspect of fluid dynamics, enabling researchers to gain valuable insights into the behavior of fluids in diverse scenarios. First, the considered method was converted into an ordinary differential equation (ODE) using an appropriate wave transformation and the solution of the ODE was supposed to be the form suggested by the used method. The trial solution includes some unknowns to be found. The trial solution and its necessary derivatives were substituted into the ODE and equating the terms with the same coefficients to zero, a system of algebraic equations was derived. Solving the system, the unknowns in the trial solution and so diverse soliton solutions of the KP equation were extracted. It is validated that the obtained solutions in analytical form satisfy the generalized KP equation via a computer algebraic system, and some plots of the solutions are demonstrated. The obtained results might be significant in terms of gaining a better understanding of the dynamic features of the generalized KP equation in fluid mechanics.

Keywords: Generalized (3+1)-dimensional Kadomtsev-Petviashvili equation, soliton, shallow water.

References:

1. Wazwaz, A.-M., Painlevé integrability and lump solutions for two extended (3+1)- and (2+1)-dimensional Kadomtsev-petviashvili equations, *Nonlinear Dynamics*, vol. 111, no. 4, pp. 3623–3632, 2022. N. Samko, Weight Hardy and singular operators in Morrey spaces, *J. Math. Anal. Appl.* 35(1) (2009) 183–188.
2. Wazwaz, A.-M. and El-Tantawy, S. A., A new (3+1)-dimensional generalized Kadomtsev-Petviashvili equation, *Nonlinear Dynamics*, vol. 84, no. 2, pp. 1107–1112, 2015.
3. Shen, Y., Tian, B., Cheng, C.D. and Zhou, T.Y., N-soliton, M th-order breather, H th-order lump, and hybrid solutions of an extended (3+ 1)-dimensional Kadomtsev-Petviashvili equation. *Nonlinear Dynamics*, pp.1-18.
4. Alquran, M., Sulaiman, T.A., Yusuf, A., Alshomrani, A.S. and Baleanu, D., Nonautonomous lump-periodic and analytical solutions to the (3+1)-dimensional generalized Kadomtsev-Petviashvili equation. *Nonlinear Dynamics*, pp.1-8.
5. Stuhlmeier, R., Stiassnie, M., Nonlinear dispersion for ocean surface waves. *J. Fluid Mech.* 859, 49 (2019).

ICOMAA 2023

Modular Grad-Div Stabilization for Linearly Extrapolated Blended Three-Step Differentiation Scheme for the Incompressible Natural Convection

Merve Ak¹, Mine Akbaş²

^{1,2}Department of Mathematics, Duzce University,
merveak67@hotmail.com
mineakbas@duzce.edu.tr

Abstract

This paper proposes a modular grad-div stabilization method for solving multiphysics flow problems. The method proposed here adds a modular grad div step for the velocity into a numerical scheme for the incompressible natural convection flow problem. The proposed numerical scheme uses an extrapolated blended three-step Backward-differentiation (BDF) in time and finite element discretization in space. We obtain the scheme is unconditionally stable over finite time interval and has optimal convergence rates both in time and in space. We also provide some numerical tests in order to support our theoretical finding and show the efficiency of the scheme.

Keywords: Modular grad-div, finite element, incompressible natural convection.

References:

1. Ravindran, S. S., An analysis of the belended three-step backward-differentiation Formula time-stepping schem for the Navier-Stokes-type system related to soret convection, Numerical Functional Analysis and Optimization. 36 (2015)658–686.
2. Cıbık, A., Eroglu, F. G., Kaya, S., Long Time Stability of a Linearly Extrapolated Blended BDF Scheme for Multiphysics Flows, Int. J. Numer. Anal. Mod. 17 (2020) 24-41.
3. Fiordilino, J. A. , Layton, W., Rong Y. An efficient modular grad-div stabilization, Computer Methods in Applied Mathematics and Engineering, 335(2018)327-346.
4. Akbas, M., Rebholz, L. G., Modular grad-div stabilization for the incompressible non-isothermal fluid flows, Applied Mathematics and Computation, 393(2021) 125748.

ICOMAA 2023

Scattering analysis of Sturm-Liouville differential equation with sign-changing weight

Merve Görgülü

Department of Mathematics, Karamanoğlu Mehmetbey University,
mervegorgulu@kmu.edu.tr

Abstract

The scattering analysis of Sturm-Liouville type singular operator with an impulsive condition and turning point will be presented in this study. Different from the classical literature [1-3] considering the positive values of the weight function, we focus on the case including an impulsive condition at the turning point. We obtain the scattering function, resolvent operator, and discrete spectrum of the operator using the hyperbolic representations of the fundamental solutions and give an example illustrating the main results [4].

Keywords: Sturm-Liouville equations, scattering theory, impulsive differential equations, turning point, discontinuous weight function.

References:

1. Gasyimov, M. G., & El-Raheem, Z. F. (1993). On the theory of inverse Sturm-Liouville problems with discontinuous sign-alternating weight, Dokl. Akad. Nauk Azerb., 48(50), 1-12.
2. El-Raheem, Z. F., & Nasser, A. H. (2014). On the spectral investigation of the scattering problem for some version of one-dimensional Schrödinger equation with turning point, Bound. Value Probl., 1, 1-12.
3. Öznur, G. B., & Bairamov, E. (2022). Scattering theory of the quadratic eigenparameter depending impulsive Sturm-Liouville equations, Turkish J. Math., 46(2), 406-415.
4. Çoşkun, N., & Görgülü, M. (forthcoming). Scattering properties of Sturm-Liouville equations with sign-alternating weight and transmission condition at turning point, Open Math. <https://doi.org/10.1515/math-2022-0566>

ICOMAA 2023

A New Transform Method and Its Application

Merve Yücel¹, Oktay Sh. Mukhtarov², Kadriye Aydemir³

¹Hitit University, Faculty of Science, Department of Mathematics, ²Tokat Gaziosmanpasa University, Faculty of Science, Department of Mathematics, ³Amasya University, Faculty of Science, Department of Mathematics

merve.yucel@outlook.com.tr

omukhtarov@yahoo.com

kadriyeaydemr@gmail.com

Abstract

Recently, different semi-analytical or numerical methods, such as the finite difference method, the Adomian decomposition method, the homotopy perturbation method, the differential transform method are of increasing interest due to the complexity of searching for analytical solutions to many initial or boundary value problems. Differential transform method (DTM) is one of the numerical methods which enables to find an approximate solution in case of linear and nonlinear systems of differential equations. The concept of the DTM was first proposed by Zhou, who solved linear and nonlinear initial value problems in electric circuit analysis. In this study, we propose a new generalization of DTM, which we call the α -parameterized differential transform method for the numerical solution of various type boundary value problems. In this work, we applied the proposed new method to solve the following illustrative differential equation

$$u''(x) + xu'(x) + \left(\frac{1}{4}x^2 + \frac{1}{2}\right)u(x) = 0, \quad x \in [0,1]$$

together with the boundary conditions

$$u(0) = 0, \quad u(1) = 1.$$

We have also, drew a graph of the approximate solution to demonstrate the reliability and effectiveness of our own method. The obtained results showed that the proposed new method can be alternative way to solve various type boundary value problems.

Keywords: Boundary value problems, differential transform method, numerical solution

References:

1. Zhou JK. Differential transformation and its application for electrical circuits. Wuuhahn, China: Huarjung University Press; 1986 [in Chinese]
2. Mukhtarov, O. S., Yücel, M., & Aydemir, K. (2020). Treatment a new approximation method and its justification for Sturm–Liouville problems. Complexity, 2020, 1-8.
3. Hassan, I. A. H. (2002). On solving some eigenvalue problems by using a differential transformation. Applied Mathematics and Computation, 127(1), 1-22.

A Finite Element Method for Solving 2D Contact Problems with Tresca Friction law

DALAH Mohamed¹, FADLIA Besma², DERBAZI Ammar³

^{1,2} Department of Mathematics, FMC-Constantine 1,

³ Department of Mathematics, University BBA

mdalah.ufmc@gmail.com

aderbazi@yahoo.com

Abstract

In this work we give the mathematical model of our phenomena. Second, we describe the classical formulation for the antiplane problem and we give the corresponding variational formulation which is given by a method coupling an evolutionary variational quality for the displacement field and a time-dependent variational equation for the electric potential field. Then we prove the existence of a unique weak solution to the model.

Keywords: Antiplane problem, electro-viscoelastic material, contact problem, friction, weak solution.

Mathematics Subject Classification: 74M10, 74F15, 49J40.

References:

1. Rademacher, Andreas. Space time finite element methods for dynamic frictional contact problems. Techn. Univ., Fak. für Mathematik, 2011.
2. G. Gilardi, I. Sharf. Literature survey of contact dynamics modeling. Mechanism and Machine Theory 37 (2002) 1213–1239.
3. R. Weyler, J. Oliver, T. Sain, J.C. Cante. On the contact domain method: A comparison of penalty and Lagrange multiplier implementations. Comput. Methods Appl. Mech. Engrg. 205–208 (2012) 68–82.
4. S. Hübner, B. Wohlmuth. Equilibration techniques for solving contact problems with Coulomb friction. Comput. Methods Appl. Mech. Engrg. 205–208 (2012) 29–45.

Bernstein type operators on elliptic domain and their interpolation properties

Mohammad Ilyas¹, Asif Khan¹ and Mohammad Mursaleen²

¹Department of Mathematics, Aligarh Muslim University, Aligarh, India

ilyas2695@gmail.com, asifjnu07@gmail.com

² Department of Medical Research, China Medical University Hospital,

China Medical University (Taiwan), Taichung, Taiwan

mursaleenm@gmail.com

Abstract

Motive of this paper is to construct univariate Bernstein-type operators $B_x^m G(x, z)$ and $B_z^n G(x, z)$, their Products $P_{mn} G(x, z)$, $Q_{nm} G(x, z)$ their Boolean sums $S_{mn} G(x, z)$, $T_{nm} G(x, z)$ on elliptic region which interpolate the given real valued function G defined on elliptic region on its boundary. The bound of the remainders of each approximation formula of corresponding operators are computed with the help of Peano's theorem and modulus of continuity, and rate of convergence for functions of Lipschitz class is computed.

Keywords: Bernstein type operators, elliptic domain, interpolation, product operators, Boolean sum operators, modulus of continuity, Peano's theorem.

References:

1. D. D. Stancu, The remainder of certain linear approximation formulas in two vari-ables, SIAM Numer. Anal. Ser. B, 1(1964), 137-163.
2. P. Blaga and G. Coman, Bernstein-type operators on triangles, Rev. Anal. Num\{e}r. Th\{e}or. Approx., 38(2009), 1, 11-23.
3. T. Catinas, Extension of some particular interpolation operators to a triangle with one curved side}, Appl. Math. Comput., 315(2017), 286-297.
4. R. E. Barnhill, G. Birkhoff and W. J. Gordon, Smooth interpolation in triangle, J. Approx. Theory, 8(1973), 2, 114-128.

ICOMAA 2023

FIXED POINT THEOREMS UNDER LOCALLY T -TRANSITIVE BINARY RELATIONS EMPLOYING MATKOWSKI CONTRACTIONS

MOHAMMAD ARIF¹, MOHAMMAD IMDAD², AFTAB ALAM³
^{1,2,3}Department of Mathematics, Aligarh Muslim University, Aligarh-202002, India,
mohdarif154c@gmail.com
mhimdad@gmail.com
aafu.amu@gmail.com

Abstract

In this paper, we extend the relation-theoretic contraction principle due to Alam and Imdad (J. Fixed Point Theory Appl. 17 (2015) 693-702) for Matkowski contractions employing a locally T -transitive binary relation. Our results improve and enrich several fixed-point theorems of the existing literature.

Keywords: Locally T -transitive binary relation, comparison function, S -connected sets.

References:

1. A. Alam and M. Imdad, "Relation-theoretic contraction principle," *J. Fixed Point Theory Appl.*, vol. 17, no. 4, pp. 693–702, 2015.
2. S. Banach, "Sur les operations dans les ensembles abstraits et leur application aux equations intgerales," *Fund. Math.*, vol. 3, no. 1, pp. 133–181, 1922.
3. F. E. Browder, "On the convergence of successive approximations for nonlinear functional equations," *Proc. K. Ned. Akad. Wet. Ser. A, Indag. Math.*, vol. 30, no. 27, p. 25, 1968.

ICOMAA 2023

Self Adaptive inertial Yosida approximation iterative algorithms for split variational inclusion and fixed point problems

Mohammad Dilshad

Department of Mathematics, University of Tabuk, Tabuk-71491, Saudi Arabia

mdilshaad@gmail.com

Abstract

We present two self-adaptive inertial iterative algorithms involving Yosida approximation to investigate a split variational inclusion problem (SVIP) and common solutions of a fixed point problem (FPP) and SVIP in Hilbert spaces. We analyze the weak convergence of the proposed iterative algorithm to explore the approximate solution of the SVIP and strong convergence to estimate the common solution of the SVIP and FPP under some mild suppositions. A numerical example is demonstrated to validate the theoretical findings, and comparison of our iterative methods with some known schemes is outlined.

Keywords: split variational inclusion; fixed point problem; Yosida approximation; algorithms; weak convergence; strong convergence

References:

1. M. Akram, M. Dilshad, A. K. Rajpoot, F. Babu, R. Ahmad, J. C. Yao, Modifies iterative schemes for fixed point problems and split variational inclusion problems, *Mathematics*, 10 (2022), 2098. <https://doi.org/10.3390/math10122098>
2. C. Byrne, Y. Censor, A. Gibali, S. Reich, Weak and strong convergence of algorithms for split common null point problem, *J. Nonlinear Convex Anal.* 13 (2012), 759--775.
3. K. R. Kazmi, S. H. Rizvi, An iterative method for split variational inclusion problem and fixed point problem for a nonexpansive mapping, *Optimization Letters*, 8 (2014), 1113--1124.
4. F. Alvarez, H. Attouch, An inertial proximal method for maximal monotone operators via discretization with damping, *Set-Valued Anal.* 9 (2001), 3—11.
5. M. Dilshad, A.F. Aljohani, M. Akram, Iterative scheme for split variational inclusion and a fixed-point problem of a finite collection of nonexpansive mappings, *J. Funct. Spaces*, 2020 (2020), 3567648, <https://doi.org/10.1155/2020/3567648>

ICOMAA 2023

On two sequential hybrid fractional q-differential inclusions

Mohammad Esmael Samei¹

¹ Department of Mathematics, Bu-Ali Sina University, Hamedan, Iran

mesamei@basu.ac.ir

mesamei@gmail.com

Abstract

In this study, we investigate a new class of the sequential fractional hybrid q-differential inclusion in Caputo sense under three-point q-integro-derivative boundary conditions. To this aim, we employ various novel analytical techniques based on α - ψ -contractive mappings, endpoints, and the fixed points of the product operators to obtain the main results. Finally, we provide some examples to illustrate our main results

Keywords: Endpoint, α - ψ -contractive, Sequential hybrid q -inclusion problem, Nonlinear analysis

References:

1. M.H. Annaby and Z.S. Mansour, *q*-Fractional Calculus and Equations, Springer Heidelberg, Cambridge, 2012.
2. R.A.C. Ferreira, Nontrivial solutions for fractional q -difference boundary value problems, *Elect. J. Qualit. Theory Diff. Eq.* 70 (2010) 1–101.
3. M.E. Samei and S. Rezapour, On a system of fractional q -differential inclusions via sum of two multi-term functions on a time scale, *Boundary Value Problems*, (2020) 2020:135.
4. M. Aydogan, D. Baleanu, J.F. Gomez Aguilar, S. Rezapour and M.E. Samei, Approximate endpoint solutions for a class of fractional q -differential inclusions, *Fractals*, 28(8) (2020), Article ID 2040029, 18 pages.
5. M. Houas, F. Martinez, M.E. Samei and M.K.A. Kaabar, Uniqueness and Ulam-Hyers-Rassias stability results for sequential fractional pantograph q -differential equations, *Journal of Inequalities and Applications*, (2022) 2022:93.

Solving A Simple Harmonic Oscillator Equation By Adomian Decomposition Method

Mohammed Mohswn¹, Yahya Qaid²

^{1,2}Department of Mathematics, Saba Region University, YEMEN

alhassnymohammed31@gmail.com

yahya217@yahoo.com

Abstract

In this paper, we will display Adomian decomposition method (ADM) for solving a simple harmonic oscillator equation. It is shown that the Adomian decomposition method (ADM) efficiency, simple, easy to use in solving physical equation. The proposed method can be applied to linear problem. Some examples were presented to show the ability of the method for linear ordinary differential physical equations.

Keywords: Adomian decomposition method, harmonic oscillator, Schrodinger equation, physical equation.

References:

1. G. Adomian. (1994). Solving frontier problem of physics, The Decomposition Method, Kluwer Academic press.
2. A.H. AL-Karawi and Inaam R. AL-Saiq (2020). Application modified Adomian Decomposition method for solving second order ordinary differential Equations, Journal of physics, 9(1): 1-18.
3. N.M. Dabwan and Y. Q. Hasan (2020). solving second order ordinary differential equation using a new modified Adomian method, Advances in mathematics: Scientific Journal 9(3): 937-943.
4. D. Andrew Lewis (2017). Introduction to differential equation, Kingston, Ontario, Canada.
5. Y. Q. Hasan and L. M. Zhu (2009). Solving second order differential equation with constant coefficients by Adomian decomposition method, Journal of concrete and Applicable Mathematics (JCAAM), 370-378.

ICOMAA 2023

Adomian decomposition method to Solve the second order ordinary differential equations with constant coefficient

Yahya Qaid¹, Mohammed Mohswn²
^{1,2}Department of Mathematics, Saba Region University, YEMEN
yahya217@yahoo.com
alhassnymohammed31@gmail.com

Abstract

In this paper, we will display Adomian decomposition method for solving second-order ordinary differential equations with constant coefficient. The Adomian decomposition method (ADM) is a creative and effective method for exact solution. It is important to note that a lot of research work has been devoted to the application of the Adomian decomposition method to a wide class of linear and non-linear problems. Some examples were presented to show the ability of method for linear and non-linear ordinary differential equations.

Keywords: Adomian decomposition method, second-order ordinary differential equations.

References:

1. G. Adomian, (1997). A review of the decomposition method and some results for nonlinear equation, Math. Model, 13(7): 17-43.
2. Y.Q. Hasan (2009). Solving second order ordinary differential equations with constant coefficients by Adomian decomposition method, JCAAM., 7(4):70-378.
3. S.Gh. Othman and Y.Q. Hasan (2020). New development of Adomian Decomposition Method for solving second order ordinary differential Equations, EPH-International Journal of Mathematics and Statistics, 6(2): 28-48.
4. S.S. Salim and Y.Q. Hasan (2022). By Adomian Decomposition Method solving the second order ordinary differential equations with singular points, International Journal of Recent Scientific Research, 13(5): 1247-1250.
5. A.M.Wazaz, Analytical approximations and pade approximations for volterra populations model, Apple, Mth. compu.102(1999) 13.

ICOMAA 2023

Computational Analysis of Degree-Based Topological Descriptors of Triphenylene Benzenoid System

Mohammed Alsharafi¹, Yusuf Zeren², Mutlu Akar³, Abdu Alameri⁴

^{1,2,3}Department of Mathematics, Yildiz Technical University, Turkey

⁴Department of Biomedical Engineering, University of Science and Technology, Yemen

alsharafi205010@gmail.com

yzeren@yildiz.edu.tr

makar@yildiz.edu.tr

a.alameri2222@gmail.com

Abstract

Topological descriptors are used to model various chemical and physical properties of molecules, such as boiling points, solubility, reactivity, biological activity, etc. By analyzing the relationship between the values of these indices and the properties of interest, QSPR and QSAR models can be developed to predict the properties of new molecules. Triphenylene is an aromatic hydrocarbon with the molecular formula $C_{18}H_{12}$. It is a planar molecule consisting of three fused benzene rings. The benzenoid triphenylene series consists of compounds derived from triphenylene by replacing one or more of its benzene rings with other aromatic rings. In this study, we computed some topological descriptors of the benzenoid triphenylene series S_r .

Keywords: Topological Descriptors, Molecular Graph, benzenoid triphenylene series.

References:

1. P. Ranjini, V. Loksha, A. Usha. Relation between phenylene and hexagonal squeeze using harmonic index, International Journal of Graph Theory. Vol. 1, no.4, (2013); pp. 116-121.
2. Y. Zeren, A. Alameri, M. Alsharafi, A. Ayache, M. Kamran Jamil. Degree-Based Molecular Descriptors of Chain Biphenylene, Biointerface Research in Applied Chemistry, Vol. 13, no. 5, (2023); 496.
3. M. Alsharafi, M. Akar, Y. Zeren, and A. Alameri. Degree-Based Topological Descriptors of Hexaphenylbenzene Molecule Graphs. Polycyclic Aromatic Compounds, (2023); (20 pages).
4. R. Pradeep Kumar, D. Soner Nandappa, and M. R. Rajesh Kanna, "Redefined Zagreb, Randic, Harmonic, GA Indices of Graphene," International Journal of Mathematical Analysis, Vol. 11, no. 10 (2017): 493–502.

ICOMAA 2023

Blending type Approximations of Sz'asz-Mirakjan operators depending on parameter λ

Mohd Raiz¹, Vishnu Narayan Mishra², Nadeem Rao³

^{1,2}Department of Mathematics, Indira Gandhi National Tribal University, Lalpur, Amarkantak, Anuppur-484887, Madhya Pradesh, India.

mohdraizp@gmail.com
vishnunarayanmishra@gmail.com
nadeemrao1990@gmail.com

Abstract

In the present manuscript, we study the approximation properties of modified Sz'asz operators with a new modification of Blending type polynomials which depends on parameters, λ and $\rho > 0$. Further, we prove a Korovkin type approximation theorem and obtain the rate of convergence of these operators. Next numerically and graphically, error analysis and convergence of the operator for the different function are presented. Moreover, local and global approximation properties are studied via the first and second-order modulus of smoothness, also Peetre's K-functional and weight functions for these sequences in different spaces of functions. Lastly, A-Statistical approximation results are investigated.

Keywords: Blending type operators, Local and global approximation, Peetre's K-functional, Rate of convergence, Korovkin theorem.

References:

1. R. Aslan, A. 'Izgi, Approximation by one and two variables of Bernstein-Schurer type operators and associated GBS operators on symmetrical mobile interval, J. Fun. Spac. 2021, 2021, 9979286.
2. 'A. Acu, M. N. Manav and M.N. Sofonea, Approximation properties of λ -Kantorovich operators. J. Inequal. Appl. 2018, 2018, 202.
3. R. Aslan, M. Mursaleen, Approximation by bivariate Chlodowsky-Sz'asz-Durrmeyer type operators and associated GBS operators on weighted spaces. J. Ineq. Appl. 2022, 2022, 26.
4. R. Aslan, M. Mursaleen, Some approximation results on a class of new type λ -Bernstein polynomials. J. Math. Ineq. 2022, 16(2), 445-462.
5. R. Aslan, Some approximation results on λ -Sz'asz-Mirakjan-Kantorovich operators. FUJMA, textbf2021, 4, 150-158.

ICOMAA 2023

A Novel Numerical Manner for Non-Linear Coupled Variable Order Reaction-Diffusion Equation

Mohd Kashif

Department of Mathematical Sciences IIT (BHU), Varanasi,
mohd.kashif.rs.mat18@itbhu.ac.in

Abstract

In this work, an efficient variable order Bernstein collocation technique, which is based on Bernstein polynomials, is applied to a non-linear coupled system of variable order reaction-diffusion equations with given initial and boundary conditions. The operational matrix of Bernstein polynomials is derived for variable order derivatives w.r.t. time and space. The Bernstein operational matrix and collocation technique are applied to the concerned non-linear physical model to achieve a system of non-linear algebraic equations, which are further solved by using Newton method. A few examples are presented to demonstrate the accuracy and stability of the scheme by comparing L_2 and L_∞ norm errors between the obtained numerical solutions and existing solutions. The important feature of this article is the graphical exhibitions of the effects of variable order derivatives on the solutions of the considered non-linear coupled reaction-diffusion equation for different particular cases.

Keywords: Variable order derivatives, diffusion equation, Bernstein Polynomials, convergence analysis, error bounds.

References:

1. Mirzaee, F., Samadyar, N., Numerical Solution Based on 2-D Orthonormal Bernstein Polynomials for Solving Some Classes of 2-D Non-linear Integral Equations of Fractional Order, *Applied Mathematics and Computation*, 344 (2019), Mar., pp. 191-203.
2. Das, S., et al., Solution of the Non-Linear Fractional Diffusion Equation with Absorbent Term and External Force, *Applied Mathematical Modelling*, 35 (2011), 8, pp. 3970-3979
3. Mirzaee, F., Hoseini, S. F., Hybrid Functions of Bernstein Polynomials and Blockpulse Functions for Solving Optimal Control of the Non-linear Volterra Integral Equations, *Indagationes Mathematicae*, 27 (2016), 3, pp. 835-849.
4. Ortigueira, M. D., et al., Variable Order Fractional Systems, *Communications in Non-linear Science and Numerical Simulation*, 71 (2019), June, pp. 231-243.
5. Yousefi, S. A., Behroozifar, M., Operational Matrices of Bernstein Polynomials and Their Applications, *International Journal of Systems Science*, 41 (2010), 6, pp. 709-716

ICOMAA 2023

Approximation by nonlinear Lupas type Bernstein operator of Max-Product kind

Mohd Shanawaz Mansoori¹, Asif Khan²

^{1,2}Department of Mathematics, Aligarh Muslim University, India

shanawaz110592@gmail.com

asifjnu07@gmail.com

Abstract

We construct a new analogue of nonlinear Lupas type Bernstein operator using max-product algebra and q -integers, which possess the endpoint interpolation property. We study the Shape-preserving property and the graphs have also been added to check the theory.

Keywords: Nonlinear max-product operators, Lupas type operators, Degree of approximation, Shape-preserving properties.

References:

1. B. Bede, L. Coroianu, S.G. Gal, Approximation and shape preserving properties of the Bernstein operator of max-product kind, *Int. J. Math. Math. Sci.*, Art. ID 590589, 26pp. (2009).
2. B. Bede, S.G. Gal, Approximation by nonlinear Bernstein and Favard-Szasz-Mirakjan operators of max-product kind, *J. Concr. Appl. Math.*, 8, 193–207 (2010).
3. L. Coroianu, S.G. Gal, Approximation by nonlinear Lagrange interpolation operators of maxproductkind on Chebyshev knots of second kind, *J. Comput. Anal. Appl.*, 13, 211–224 (2011).
4. O. Duman, Statistical convergence of max-product approximating operators, *Turkish J. Math.*34, 501–514 (2010).
5. O. Duman, Nonlinear Approximation: q -Bernstein Operators of Max-Product Kind. In: Anastassiou G., Duman O. (eds) *Intelligent Mathematics II: Applied Mathematics and Approximation Theory. Advances in Intelligent Systems and Computing*, vol 441. Springer, Cham., 2016. https://doi.org/10.1007/978-3-319-30322-2_3

ICOMAA 2023

New Auxiliary Filled Function without Parameter for Solving Optimization Problems

Muhammad Wasim Awan¹, Ahmet ŞAHINER², Temel ERMIŞ³,

^{1, 2} Department of Mathematics, Suleyman Demirel University, Isparta, 32260, Turkey

³ Department of Mathematics and Computer Sciences, Eskisehir Osmangazi University, 26480, Eskisehir

Wasimawan2012@gmail.com

ahmetsahiner@sdu.edu.tr

termis@ogu.edu.tr

Abstract

This study offers a new filled function method to solve unconstrained global optimization problems. First, we analytically show that our filled function holds all conditions to be a filled function. Then an algorithm constructed for this new filled function and apply constructed algorithm to an unconstrained global optimization problem. At the end we present the numerical results and compare with results of other competing algorithms in literature to show the efficiency of our filled function method.

Keywords: Global optimization, Filled function, Global minimizer, Algorithm, Local minimizer.

References

1. Ge, R.: The theory of filled function method for finding global minimizers of nonlinearly constrained minimization problems. *J. Comput. Math.* **5**, (1987) 1–9.
2. El-Gindy, T.M., Salim, M.S., Ahmed, A.I.: A new filled function method applied to unconstrained global optimization. *Appl. Math. Comput.* **273**, (2016) 1246–1256.
3. A. I. Ahmed: A new parameter free filled function for solving unconstrained global optimization problems, *International Journal of Computer Mathematics*, 98:1, (2021) 106-119.
4. Ridwan Pandiya, A Novel Parameter-Free Filled Function Applied to Global Optimization Volume 29, Issue 1: *Engineering Letters*, 29:1, EL-29(2021) 1-22.
5. Ahmet Sahiner, Nurullah Yilmaz and Gulden Kapusuz, a novel modeling and smoothing technique in global optimization, *journal of industrial and management optimization*, Vol.15, Num.1, (2019) 113-130.

ICOMAA 2023

Risk Assessment of Autonomous Vehicles in terms of Public Health

Murat Kirişci¹

¹*Department of Biostatistics and Medical Informatics, Istanbul University-Cerrahpaşa,
Fatih, Istanbul, Türkiye
mkirisci@hotmail.com*

Abstract

Risk assessment is required to manage the negative consequences of Autonomous Vehicle (AV) implementation and maximize its benefits. In this study, which will be conducted on the beneficial and negative effects of AVs on public health, risk factors related to transportation that affect public health will be given first. When uncertainties are encountered in hazard assessment and risk assessment for transport-related exposures and risk factors that may affect public health, a new method will be developed to overcome these problems. Examining the health effects of AVs, identifying risks, and trying to reduce these risks will help produce strategies that are beneficial to overall public health.

Keywords: Autonomuou vehicles, public health, risk assessment, AHP, TOPSIS, MABAC

References:

1. S. Sohrabi, H. Khreis, & D. Lord. Impacts of autonomous vehicles on public health: a conceptual model and policy recommendations. *Sustainable cities and society*, 2020, 63: 102457.
2. D. Rojas-Rueda, M. J. Nieuwenhuijsen, H. Kheris, H. Frumkin. Autonomous Vehicles and Public Helath, *Annu. Rev. Public Health* 2020. 41:18.1–18.17
3. İ. Kaya, A. Karaşan, B. Özkan & M. Çolak. An integrated decision-making methodology based on Pythagorean fuzzy sets for social robot evaluation. *Soft Computing*, 2022; 26:9831-9858.
4. B. Kelley. Public Helath, Autonomous Automobiles, and the Rush to Market. *Journal of Public Health Policy*, 2017; 38: 167-184.

ICOMAA 2023

Childhood Cancer Risk Analysis with Integrated Decision-Making Method

Murat Kirişci¹

¹Department of Biostatistics and Medical Informatics, Istanbul University-Cerrahpaşa,
mkirisci@hotmail.com

Abstract

Childhood cancer is the leading cause of death in children, especially in low and middle-income countries. Their likelihood of survival heavily rests on the country where they live. The chance of curing childhood cancer in high-income countries is above eighty percent, whereas in low and in middle-income countries, it is a mere forty-five percent. The most productive way to reduce the effects of childhood cancer is through effective and evidence-based therapy with appropriate nurturing care.

To address the risk assessment process at the moment, a new risk evaluation technique will be offered. The suggested technique uses MCDM, which has a hybrid structure made up of the Neutrosophic AHP and Fermatean Fuzzy AHP methods, to enable dealing with ambiguity in the risk assessment process for children cancer.

Keywords: Childhood cancer, risk analysis, fermatean fuzzy AHP, Neutrosophic AHP.

References:

1. S. Habib, M. Akram, M.M. Ali Al-Shamri. Comparative analysis of pythagorean MCDM methods for the risk assessment of childhood cancer. *Computer Modeling in Engineering & Sciences*, 135 (3), 2023, 2585-2615.
2. Johnston, W. T., et al. "Childhood cancer: Estimating regional and global incidence." *Cancer epidemiology* 2021; 71:101662.
3. M. Kirisci, N. Şimşek. Decision making method related to Pythagorean Fuzzy Soft Sets with infectious diseases application. *Journal of King Saud University-Computer and Information Sciences*, 2022, 34.8: 5968-5978.
4. L.G. Spector, P. Nathan, L. M. Erin. Genetic and nongenetic risk factors for childhood cancer. *Pediatric Clinics* 2015; 62: 11-25.

ICOMAA 2023

Some applications of Krasnosel'skii's theorem for multi-valued operators

Cesim Temel¹, Müberra Selah²

^{1,2}Department of Mathematics, Faculty of Sciences, Van Yuzuncu Yil University, 65080, Van/Turkey

cesimtemel@yyu.edu.tr
muberra.selah@gmail.com

Abstract

In this work, we discovered some results about the existence of a solution of the sum of a multi-valued operator and a single-valued operator in the weak topology. The results are specifically presented using the weak non-compactness technique. Some previously known single-valued Krasnosel'skii-type theorems are extended into multivalued versions.

Keywords: Multivalued operator, Krasnosel'skii's theorem, Measure of weak noncompactness.

References:

1. R. Agarwal and D. O'Regan. (2002). Fixed-point theory for weakly sequentially upper semicontinuous maps with applications to differential inclusions. *Nonlinear Oscillations*, 5, 277-286.
2. A. Ben Amar and A. Sikorska-Nowak. (2011). On some fixed point theorems for 1-set weakly contractive multi-valued mappings with weakly sequentially closed graph. *Adv. Pure Math.* 1, 163-169.
3. M.A. Taoudi. (2010). Krasnosel'skii type fixed point theorems under weak topology features, *Nonlinear Anal.* (1)72, 478-482.
4. C. Temel. (2019). Multivalued types of Krasnosel'skii's fixed point theorem for weak topology. *U.P.B. Sci. Bull., Series A*, 81 (2), 139-148.

ICOMAA 2023

Basicity of double system of exponents with linear phase in the weighted Lebesgue space

Natavan Nasibova¹, Nazir Ismailov²

¹Azerbaijan State Oil and Industry University, Baku, Azerbaijan,

²Ganja State University, Ganja, Azerbaijan,

natavan2008@gmail.com

department2011@mail.ru

Abstract

This paper considers double system of exponentials with linear phase in the weighted space $L_{p,\rho}$ with power weight $\rho(\cdot)$ on the segment $[-\pi, \pi]$. Under certain conditions on the weight function $\rho(\cdot)$ and on the perturbation parameters, the completeness, minimality and basicity of this system in $L_{p,\rho}$ is proved. An explicit expression for the biorthogonal system in the case of minimality is derived and its integral representation is obtained. The obtained results generalize all previously known results in this direction.

Keywords: exponential system, basicity, weighted space.

References:

1. A.Sh. Shukurov, N.A. Ismailov, On the Minimality of Double Exponential System in Weighted Lebesgue Space, Azerbaijan Journal of Mathematics, V. 7, No 1, 2017, pp. 125-134.
2. B.T. Bilalov, Z.G. Guseinov, Basicity of a system of exponents with a piece-wise linear phase in variable spaces. Mediterr. J. Math., 9(3), 2012, 487-498
3. B.T. Bilalov, Bases of a system of exponentials in L_p , Dokl. Akad. Nauk, 392(5), 2003, 583-585

ICOMAA 2023

On Harmonic Summability of Difference Sequences of Fractional Order

Nazlım Deniz Aral¹, Hacer Şengül Kandemir², Mikail Et³

¹Department of Mathematics, Bitlis Eren University,
ndaral@beu.edu.tr

²Faculty of Education, Harran University,
hacer.sengul@hotmail.com

³Department of Mathematics, Fırat University,
mikaillet68@gmail.com

Abstract

In this work, we generalized the concept of lacunary $(H, 1)$ summability from real sequences to difference sequences of fractional order. The notation of the lacunary harmonic summability, lacunary strongly harmonically summability and lacunary statistically logarithmic convergence of difference sequences of fractional order is introduced. Also we have established theorems presenting a connection between these notions.

Keywords: Harmonic convergence, lacunary sequence, difference sequence.

References:

1. Fridy, J., Orhan, C. Lacunary statistical convergence, Pacific J. Math. Vol.160, No. 1 (1993) 43-51.
2. Nuray, F. Lacunary Statistical Harmonic Summability, Journal of Applied Analysis and Computation. Vol. 12, No.1 (2022).
3. Şengül Kandemir, H., Et, M. On Harmonic Summability of Order α . doi: <https://doi.org/10.21203/rs.3.rs-1064137/v1> (2022).

ICOMAA 2023

Existence and blow up of solutions for a viscoelastic Petrovsky equation with variable exponents

Nebi Yılmaz¹, Erhan Pişkin²

¹*Department of Mathematics, Dicle University*

²*Institute of Natural and Applied Sciences, Dicle University*

nebiyilmaz1981@gmail.com

episkin@dicle.edu.tr

Abstract

In this presentation, we consider a viscoelastic Petrovsky equation with variable exponents. This type problem occurs in many mathematical models of applied science, such as electrorheological fluids, population dynamics, heat transfer, chemical reactions. We prove the existence and blow up of solutions.

Keywords: Blow up, existence, Petrovsky equation, variable exponents.

References:

1. Stanislav N. Antontsev, Jorge Ferreira, Erhan Pişkin, Sebastiao Martins Siqueira Cordeiro, Existence and non-existence of solutions for Timoshenko-type equations with variable exponents, *Nonlinear Analysis: Real World Applications*, 61 (2021) 103341: 1-13.
2. E. Pişkin, B. Okutmuşur, *An Introduction to Sobolev Spaces*, Bentham Science, 2021.
3. Erhan Pişkin, Nebi Yılmaz, Blow up of solutions for a system of strongly damped Petrovsky equations with variable exponents, *Acta Universitatis Apulensis*, 71 (2022) 87-99.

ICOMAA 2023

Stability and stabilization of the implicit differential equations in Hilbertspaces

Nor-El-Houda Beghersa¹ and Mehdi Benabdallah²

¹Department of Mathematics, University of Sciences and Technology Mohamed Boudiaf of Oran
USTO-MB, El Mnaouar, BP, 1505, Bir El Djir 31000, Oran, ALGERIA.

norelhouda.beghersa@univ-usto.dz

²University of Sciences and Technology Mohamed Boudiaf of Oran USTO-MB, El Mnaouar, BP, 1505,
Bir El Djir 31000, Oran, ALGERIA.

mehdi.benabdallah@univ-usto.dz

Abstract

We apply the generalization of Lyapunov theorem for some stationary differential equations of the form:
 $A'x(t)=Bx(t)$, for all $t \geq 0$;

Where A and B are bounded linear operators in Hilbert spaces, in order to obtain a necessary and sufficient conditions about the stability of those equations. The achieved results can be used also to study the stability of another type of differential equations called quasi-linear degenerate differential equations.

Keywords: Exponential stability, Operator theory, Implicit equations, Hilbert space.

References:

1. R. Bellman and K.L. Meskhi, Differential-Difference Equations, vol 6, Academic Press. London, (1963).
2. M. Benabdallah, A.G. Rutkas and A.A Soloviev, On the stability of degenerate difference systems in Banach spaces, J.Sov. Math, 57(1991) 3435–3439.
3. Doleckii, J. L. Krein, M. G. (1975). Stability of solutions of differential equations in Banach space. Amer. Math. Soci. Providence. RI.
4. A. Favini, A. Yagi, Degenerate differential equations in Banach spaces, Marcel Dekker-Inc. New York- Basel-Hong Kong, (1999).
5. A.G. Rutkas, Spectral methods for studying degenerate differential-operator equations, J. Math.Sci, 144(2007) 4246–4263.

ICOMAA 2023

On the exponential stability of the stationary and perturbed implicit equations

Nor-El-Houda Beghersa¹ and Mehdi Benabdallah²

¹Department of Mathematics, Faculty of Mathematics and Computer Sciences, University of Sciences and Technology Mohamed Boudiaf of Oran USTO-MB, El Mnaouar, BP, 1505, Bir El Djir 31000, Oran, ALGERIA

norelhouda.beghersa@univ-usto.dz

²Department of Mathematics, Faculty of Mathematics and Computer Sciences, University of Sciences and Technology Mohamed Boudiaf of Oran USTO-MB, El Mnaouar, BP, 1505, Bir El Djir 31000, Oran, ALGERIA

mehdi.benabdallah@univ-usto.dz

Abstract

In this work, we study the exponential stability of the stationary differential equations of the form: $Ax'(t) - Bx(t) = 0$ for all $t \geq t_0$, where A and B are linear bounded operators in Hilbert spaces. The obtained results on the one hand are the generalization of the Liapounov theorem for the spectrum of the operator pencil $\lambda A - B$. On the other hand, the establishment of the exponential stability conditions for the stationary and perturbed equation described by: $Ax'(t) - (B + B(t))x(t) = 0$.

Keywords: Exponential stability, Operator pencil, Implicit stationary equation, Perturbed equation.

References:

1. M. Benabdallah and A.G. Rutkas, A.A. Soloviev, On the stability of degenerate difference systems in Banach spaces, J.Sov.Math, 57(1991) 3435–3439.
2. Y. Doletski, M.G. Krein, Stability of solutions of differential equations in Banach spaces, Transactions of math monographs 43. American Math Society Providence. (1974).
3. Gantmacher F.R. The theory of matrices. Chelsea Publishing Company: New York. Vol. 1, 2, (1960).
4. Rutkas, A. G., (2007). Spectral methods for studying degenerate differential-operator equations. Springer Science, 144: 4246-4263.

ICOMAA 2023

Finite volume method for Seawater intrusion problem in Porous Media

Noura Djedaidi¹, Souad Gasmi² and Fatma Zohra Nouri³
*^{1,2,3}Mathematical Modeling and Numerical Simulation Research Laboratory,
Badji Mokhtar University, Annaba, Algeria*

souad_gasmi@yahoo.com

tassili.nan09@gmail.com

¹University Algiers 1, Algiers, Algeria

noura.djedaidi@gmail.com

Abstract

In this paper, we are interested in studying the dynamics of seawater intrusion problem in heterogeneous porous media. This problem concerns immiscible fluids, where these fluids are separated by an abrupt interface. Here we propose a numerical approach to solve our problem by using finite volume method and then present numerical results.

Keywords: Seawater intrusion, Porous media, Finite volume method, Stability.

References:

1. Bear, J. (1972). Dynamics of Fluids in Porous Media, Elsevier, New York.
2. Djedaidi, N. & Nouri, F.Z. (2016). A study of sea water flow interactions in heterogeneous porous media', International Journal of Ecology and Development, Vol. 31, No. 2, pp.70–77.
3. Eymard, R., Gallouet, T. & Herbin, R. (2000). Finite volume methods, Handbook of numerical analysis, North Holland, P.G. Ciarlet, J.L. Lions eds, vol 7, pp 713-1020.
4. Gasmi, S. & Nouri, F.Z. (2015). Numerical simulation for two phase flow in a porous medium, Boundary Value Problems, Vol. 7, pp.1–7.

ICOMAA 2023

Existence and nonexistence results of a fifth-order BVP

Nourredine Houari¹, Faouzi Haddouchi^{1,2}

¹Laboratory of Fundamental and Applied Mathematics of Oran (LMFAO), Department of Mathematics, University of Oran 1, Oran, Algeria,

²Department of Physics, University of Sciences and Technology of Oran-MB, Oran, Algeria ,

nouredlinehouari.mi@gmail.com

fhaddouchi@gmail.com

Abstract

The aim of this talk, is to discuss the existence and nonexistence of solutions for a class of nonlinear fifth-order boundary value problems. The main tool of our analysis is the classical norm type compression-expansion fixed point theorem of Krasnoselskii.

Keywords: Existence, nonexistence, boundary value problem, Krasnoselskii fixed point theorem.

References:

1. F. Haddouchi, N. Houari, Monotone positive solutions of fourth order boundary value problem with mixed integral and multi-point boundary conditions, J. Appl. Math. Comput. 66 (2021), 87–109.
2. N. Houari, F. Haddouchi, Existence and nonexistence results for fifth-order multi-point boundary value problems involving integral boundary condition, Filomat, 37-19 (2023), 6463–6486.
3. S. N. Odda, Existence solution for 5th order differential equations under some conditions, Appl. Math. 1 (2010), 279-282.
4. M. A. Krasnosel'skii, Positive Solutions of Operator Equations, P. Noordhoff, Groningen, The Netherlands, 1964.

ICOMAA 2023

Multiple Pursuers and One Evader in A Pursuit Differential Game with Gronwall-Type Constraints

Gafurjan Ibragimov¹, Omongul Egamberganova², Risman Mat Hasim³

¹Department of General and Exact Subjects, Tashkent State University of Economics, 100006, Tashkent, Uzbekistan,

²Department of Mathematics, University Putra Malaysia, Serdang 43400, Malaysia and Department of Higher and Applied Mathematics, Tashkent Institute of Finance, Tashkent 100000, Uzbekistan,

³Department of Mathematics, University Putra Malaysia, Serdang 43400, Malaysia;

⁴ Department of Mathematics, University Putra Malaysia, Serdang 43400, Malaysia

gofurjon.ibragimov@tsue.uz

intelegentniy21@gmail.com and gs64479@student.upm.edu.my

risman@upm.edu.my

Abstract

We study a differential game with many pursuers and one evader in \mathbf{R}^n . The control functions of the players are constrained by Gronwall type constraints. If the state of a pursuer coincides with the state of the evader, then pursuit is considered completed. Pursuers try to complete the game and the aim of the evader is opposite. We obtain the conditions of completion of pursuit and possibility of evasion in terms of the convex hull of the initial positions of pursuers.

Keywords: Differential game, Gronwall constraints, pursuer, evader, strategy.

References:

1. Garcia E., Casbeer D.W., Von Moll A., Pachter M. (2021). Multiple Pursuer Multiple Evader Differential Games. *IEEE Transactions on Automatic Control*, 66(5): 2345--2350, May 2021, doi: 10.1109/TAC.2020.3003840.
2. Ibragimov G.I., Ferrara M., Kuchkarov A.Sh., Pansera B.A. (2018). Simple motion evasion differential game of many pursuers and evaders with integral constraints. *Dynamic Games and Applications*. 8: 352--378. <https://doi.org/10.1007/s13235-017-0226-6>.
3. Makkapati V.R., Tsiotras P. (2019). Optimal Evading Strategies and Task Allocation in Multi-player Pursuit-Evasion Problems. *Dyn Games Appl* 9, 1168--1187. <https://doi.org/10.1007/s13235-019-00319-x>
4. Samatov B.T., Ibragimov G.I., Khodjibayeva I.V. (2020) Pursuit-evasion differential games with Gronwall type constraints on controls. *Ural Mathematical Journal*, 6(2): 95--107, DOI: 10.15826/umj.2020.2.010.

Gauss Decomposition of \mathbb{Z}_3 -Graded Quantum Group $\widetilde{GL}_q(2)$

Salih Çelik¹, Öztürk Savaş²

^{1,2}Department of Mathematics, Yıldız Technical University,

sacelik@yildiz.edu.tr

ozturksavas1992@gmail.com

Abstract

The general linear group obtained by the matrix representations of the linear transformations is denoted by $GL(n)$. This group is deformed in two ways. In the classical and deformed cases, a matrix T belonging to the group $GL(n)$ can be written as $T = T_L T_D T_U$, where T_L , T_D and T_U are a lower triangular, a diagonal, and an upper triangular matrices, respectively. In the literature, it has been shown that this case is also valid for a matrix belonging to the quantum group $GL_q(n)$ and the quantum supergroup $GL_q(m|n)$.

In this study, we show that the Gauss decomposition is valid for the \mathbb{Z}_3 -graded quantum group $\widetilde{GL}_q(2)$ as well. When the Gauss decomposition of a matrix in the quantum group $\widetilde{GL}_q(2)$ is performed, two new \mathbb{Z}_3 -graded quantum subgroups will emerge and the properties of these quantum subgroups will be examined: If $T \in \widetilde{GL}_q(2)$, we can write $T = T_L T_D T_U$. Then, it can be seen that the product matrices $T_L T_D$ and $T_D T_U$ form both \mathbb{Z}_3 -graded quantum groups. The coordinate algebras of both quantum subgroups have a \mathbb{Z}_3 -graded Hopf algebra structure. Finally, it has been seen that the product of three matrices in the Gauss decomposition of a matrix belonging to the quantum group $\widetilde{GL}_q(2)$ also admits a Hopf algebra structure.

Keywords:

\mathbb{Z}_3 -graded Hopf algebra, \mathbb{Z}_3 -graded quantum group, Gauss decomposition.

References:

1. S. Çelik, "Differential geometry of the \mathbb{Z}_3 -graded quantum superplane", J. Phys. A: Math. Gen., vol. 35, pp. 4257–4268, 2002.
2. S. Çelik, "A differential calculus on \mathbb{Z}_3 -graded quantum superspace $\mathbb{R}_q(2|1)$ ", Algebras Representation Theory, vol. 19, pp. 713–730, 2016.
3. S. Çelik, "A new \mathbb{Z}_3 -graded quantum group", J. Lie Theory 27, pp. 545–554, 2017.
4. S. Çelik and S. A. Celik, "Differential calculi on \mathbb{Z}_3 -graded Grassmann plane", Adv. Appl. Clifford Algebras, vol. 27, pp. 2407–2427, 2017.
5. S. Celik, "Left-covariant differential calculi on $\widetilde{GL}_q(2)$ ", J. Math Phys., vol. 62, 073504:1-16, 2021.

Inverse Boundary Problem for Construction of the Right Side in a Third Order Two-Dimensional Pseudohyperbolic Equation

Yashar Mehraliyev¹, Rena Sadikhzada²

^{1,2}Department of Differential and Integral Equations, Baku State University,

yashar_aze@mail.ru

rena7097@gmail.com

Abstract

In mathematics and mathematical modeling, tasks that require the determination of the coefficient (coefficients) of the equation itself or the right-hand side of equation in addition to solving a definite differential equation are called inverse problems.

In this paper, we study the inverse boundary problem for a third order two-dimensional pseudohyperbolic equation. When solving inverse boundary problem, a transition from an initial boundary problem to some auxiliary problem is performed.

After applying the formal Fourier scheme to the auxiliary problem, the auxiliary problem is reduced to a decomposable system of integral equations. Then the existence and uniqueness of the integral equation will be proved. It will also be shown that the solution of the integral equation is the solution of the auxiliary problem.

Keywords: inverse boundary problem, third-order pseudohyperbolic equation, Fourier method.

References:

1. Kozhanov A.I., Safiullova R.R., On a class of pseudohyperbolic equations with an unknown coefficient, Chelyab. Phys-Math. journal, 2022, volume 7, issue 2, 164180
2. Mehraliev Ya. T. On the solvability of an inverse boundary value problem for a fourth order pseudohyperbolic equation with an additional integral condition // Izv. universities. Povolzh. region. Ser. Phys.-Math. Sciences. 2013. No. 1. S. 1933.
3. Romanov VG Inverse problems of mathematical physics. M. : Nauka, 1989.

ICOMAA 2023

A new family of symmetric functions of binary products of bivariate Mersenne Lucas polynomials with (p,q) -numbers

Rokiya Sahali¹ and Ali Boussayoud²

¹LMAM Laboratory and Department of Mathematics,
Mohamed Seddik Ben Yahia University, Jijel, Algeria
rokasali18@gmail.com

²LMAM Laboratory and Department of Mathematics,
Mohamed Seddik Ben Yahia University, Jijel, Algeria
alboussayoud@gmail.com

Abstract

In this work, we study the generalized (p,q) -numbers and bivariate Mersenne Lucas polynomials defined in [4, 6]. We first give some new properties and results on the generalized (p,q) -numbers. Moreover, by using the symmetric functions we obtain the new generating functions for the products of bivariate Mersenne and bivariate Mersenne Lucas polynomials with (p,q) -numbers at positive and negative indices.

Keywords: (p,q) numbers; generating functions; explicit formula; Bivariate Mersenne Lucas polynomials.

References:

1. A. Boussayoud, M. Kerada, Symmetric and generating functions, Int. Electron. J. Pure Appl. Math., 7, 195-203, 2014.
2. A. Suvarnamani, M. Tatong, Some properties of (p,q) -Fibonacci numbers, Science and Technology RMUTT Journal, 5, 17-21, 2015.
3. F.R.V. Alves, Bivariate Mersenne polynomials and matrices, Notes Number Theory Discrete Math., 26, 83-95, 2020.
4. N. Saba, A. Boussayoud, A. Abderrezzak, Symmetric and generating functions of generalized (p,q) -numbers, Kuwait J. Sci., 48, 1-15, 2021.
5. N. Saba, A. Boussayoud, K.V.V. Kanuri, Mersenne Lucas numbers and complete homogeneous symmetric functions, J. Math. Computer Sci., 24, 127-139, 2022.

Quasilinear parabolic equations with partialy VMO coefficients

Rosamaria Rescigno, Lubomira Softova
Department of Mathematics, University of salerno,
rorescigno@unisa.it


Abstract

We study strong solvability of the Cauchy-Dirichlet problem for parabolic quasilinear equations with discontinuous principal coefficients. The dependence on x is of VMO type while with respect to t , the coefficients are only measurable. We obtain existence and uniqueness of the solution, under suitable structural conditions, and by the imbedding theorems also Holder regularity of the solution.

Keywords: Quasilinear parabolic equations, Fixed point theorem, strong solutions, VMO x coefficients.

References:

1. Krylov N., Parabolic and elliptic coefficients with VMO coefficients, Commun. Partial Differ. Equ., 32 (3), (2007), 453-475.
2. Rescigno R., Softova L., Quasilinear parabolic equations with partialy VMO coefficients (submitted)
3. Softova L., Quasilinear parabolic operators with discontinuous ingredients, Nonlinear Anal., Theory Methods Appl., Ser A, 52 (4), (2003), 1079-1093.



ICOMAA 2023

Impact of large claims on the stability bound of a bivariate risk model

Safia Hocine¹, Djamil Aïssani², Aïcha Bareche³

^{1,2,3}Research Unit LaMOS, Faculty of Exact Sciences, University of Bejaia, Algeria.

safia.hocine@univ-bejaia.dz

djamil.aissani@univ-bejaia.dz

aicha.bareche@univ-bejaia.dz

Abstract

In ruin theory, stochastic processes are used to model the surplus of an insurance company and to evaluate its ruin probability. This characteristic is a much studied risk measure in the literature. In general, this measure is very difficult or even impossible to evaluate explicitly. Thus, different approximation methods have been proposed to estimate this characteristic [2]. However, its assessment is not evident in several cases since it cannot be found explicitly in closed forms. Besides, it is not easy to determine the parameters governing these models since they are often unknown or partially known. For all these reasons, the stability analysis becomes crucial for studying such models. Hence, it is necessary to obtain explicit stability bounds. The strong stability method, which was developed by Aïssani and Kartashov (1983) [1], makes it possible to clarify the conditions for which the ruin probability of the complex risk model (real model) can be approximated by the corresponding ruin probability in the simple risk model (ideal model). In this work, we consider the problem of stability of the bivariate classical risk model, in particular at the stability bounds of the classical risk model established by Benouaret and Aïssani (2010) [4] using the approach based on the strong stability method developed for general Markov chains [1]. Based on numerical and graphical results, we study the stability bounds of the ruin probability considering different heavy-tailed claim distributions [3], and we show the impact of large claims on the stability bound of a bivariate classical risk model.

Keywords: Risk model, Ruin probability, Approximation, Large claims, Strong stability, Markov chain.

References:

1. D. Aïssani, and N. V. Kartashov, Ergodicity and stability of Markov chains with respect to operator topology in the space of transition kernels, Doklady Akademii Nauk Ukrainkoï S.S.R., Series A (11) : 3-5, 1983.
2. S. Asmussen, and H. Albrecher, Ruin Probabilities, Advanced Series on Statistical Science & Applied Probability, World Scientific Publishing, Singapore, Vol. 14, 2010.
3. T. Buch-Larsen, J.P. Nielsen, M. Guillen and C. Bolancé. Kernel density estimation for heavy-tailed distribution using the Champernowne transformation, Statistics, (6) : 503-518, 2005.
4. Z. Benouaret, and D. Aïssani, Strong stability in a two dimensional classical risk model with independent claims, Scandinavian Actuarial Journal, 2010(2) : 83-92, 2010.

Global solvability of the Laplace equation in weighted Sobolev spaces

B. Bilalov, N.Nasibova, L.Softova, S. Tramontano

¹*Department of Mathematics, University of Salerno*
lsoftova@unisa.it

Abstract

We consider a non-local boundary value problem for the Laplace equation in unbounded domain. Our interest is pointed on the weak solvability of that problem in the framework of the weighted Sobolev space with Muckenhoupt weight. We have proved that any weak solution heaving second generalized derivatives is also a strong solution and satisfies the corresponding boundary conditions, extending in such a way some classical results.

Keywords:Laplace equation, weak solutions, anisotropic Lebesgue spaces, Morrey spaces.

References:

1. **Bilalov B., Nasibova N., Softova L, Tramontano S.,** Global solvability of the Laplace equation in weighted Sobolev spaces (submitted).
2. **Moiseev E.I.,** The solution of a nonlocal boundary value problem by the spectral method, Differ. Equations, 35 (8, 1105--1112 (1999), transl. from Differ. Uravn. 35, (8), 1094-1100 (1999).
3. Zygmund A., Trigonometric Series, Vol. I and II combined, 3rd ed., Cambridge: Cambridge University Press, pp. 364 (2002).



ICOMAA 2023

On Evolution Equations and Operator Semigroups in Banach Spaces

Erdal Gül¹, Samet Karasu²

^{1,2}*Department of Mathematics, Yildiz Technical University,*

gul@yildiz.edu.tr

sametkrs58@gmail.com

Abstract

The theory of operator semigroups is a field of study widely used in many areas of applied mathematics, including analysis, probability theory, partial differential equations, dynamical systems, and quantum theory. In this work, the qualitative properties of solutions of evolution equations corresponding to different types of initial value problems considered for applications will be analyzed through one-parameter semigroups on Banach spaces.

Keywords: Linear operator, Banach space, Semigroup, Evolution equation

"This work was supported by Research Fund of the Yildiz Technical University. Project Number: FYL-2023-5606"

References:

1. E.B. Davies, One-Parameter Semigroups, Academic Press, London (1980).
2. L. C. Evans, Partial Differential Equations, Graduate Studies in Mathematics, AMS, Providence, Rhode Island, Vol. 19, 1998 (Reprinted 2002).
3. J. A. Goldstein, Abstract evolution equations, Trans. Amer. Math. Soc. 141 (1969), 159-185.
4. S. Kesavan, Topics in Functional Analysis and Applications, New Age International(P) Limited, New Delhi, 1989 (Reprint 2003).
5. A. Pazy, Semigroups of Linear Operators and Applications to Partial Differential Equations, Springer-Verlag, Applied Math. Sciences, Vol. 44, 1983.

ICOMAA 2023

Data Dependence Result for A New Iterative Algorithm

Samet Maldar¹ and YunusAtalan²

^{1,2}*Department of Mathematics, Aksaray University,*

mmaldar@aksaray.edu.tr

yunusatalan@aksaray.edu.tr


Abstract

In this work, we defined a different version of an iteration method known in the literature. We examined the convergence of this new iteration method under certain conditions. We also proved that this iteration method is data-dependent. Finally, we presented some examples for the obtained results.

Keywords: Iteration method, Convergence, Data dependence.

References:

1. Noor, M.A.; Noor, K.I. (2013). Some parallel algorithms for a new system of quasi variational inequalities, Applied Mathematics & Information Sciences, 2493-2498.
2. Noor, M.A.; Noor, K.I.; Khan, A.G. (2014). Parallel schemes for solving a system of extended general quasi variational inequalities, Applied Mathematics and Computation, 245, 566–574.
3. Maldar, S. (2022). New parallel fixed-point algorithms and their application to a system of variational inequalities. Symmetry, 14(5), 1025.



ICOMAA 2023

An explicit Milstein-type scheme for interacting particle systems and McKean--Vlasov SDEs with common noise and non-differentiable drift coefficients

Sani Biswas¹, Chaman Kumar¹, Neelima², Gonçalo Dos Reis³, Christoph Reisinger⁴

¹Indian Institute of Technology Roorkee, ²University of Delhi, ³Edinburgh University, ⁴Oxford University
sbiswas2@ma.iitr.ac.in

Abstract

We propose an explicit drift-randomised Milstein scheme for both McKean–Vlasov stochastic differential equations and associated high-dimensional interacting particle systems with common noise. By using a drift-randomisation step in space and measure, we establish the scheme's strong convergence rate of 1 under reduced regularity assumptions on the drift coefficient: no classical (Euclidean) derivatives in space or measure derivatives (e.g., Lions/ Fréchet) are required. The main result is established by enriching the concepts of bistability and consistency of numerical schemes used previously for standard SDE. We introduce certain Spijker-type norms (and associated Banach spaces) to deal with the interaction of particles present in the stochastic systems being analysed. A discussion of the scheme's complexity is provided.

Keywords: stochastic interacting particle systems, McKean-Vlasov equations, common noise, Milstein scheme, non-differentiable drift, drift randomisation, bistability.

References:

1. C. Kumar, Neelima, C. Reisinger and W. Stockinger. Well-posedness and tamed schemes for McKean–Vlasov equations with common noise, *Ann. Appl. Probab.*, 32 (5) (2021), 3283-3330.
2. P. Cardaliaguet (2013). Notes on Mean-field games, notes from P. L. Lions lectures at Collège de France.
3. R. Kruse and Y. Wu. A randomised Milstein method for stochastic differential equations with non-differentiable drift coefficients, *Discrete Contin. Dyn. Syst. Ser. B*, 24(8)(2019), 3475-3502.
4. S. Biswas, C. Kumar, Neelima, G. dos Reis and C. Reisinger. An explicit Milstein-type scheme for interacting particle systems and McKean–Vlasov SDEs with common noise and non-differentiable drift coefficients (2022).

ICOMAA 2023

Numerical Investigation of Free Rotor Aerodynamics for Ship-Rotor Interaction

Sarih Sari¹, Ali Dogrul², Seyfettin Bayraktar³

¹Department of Naval Arch. & Mar. Eng., Yildiz Technical University,

²Department of Naval Arch. & Mar. Eng., Turkish Naval Academy, National Defense University,

³Department of Mar. Eng. Op., Yildiz Technical University,

sarih@yildiz.edu.tr

adogrul@dho.edu.tr

sbay@yildiz.edu.tr

Abstract

Rotor aerodynamics is important for rotary-wing aircrafts especially located in naval surface combatants. The aerodynamic interaction between the helicopter, its rotor and the ship structure should be investigated precisely. This study focuses on the aerodynamic performance of 4-blade rotor geometry to be used in the further studies. Sikorsky S-76 main rotor geometry was analyzed numerically in free condition in an unbounded flow domain. RANS equations were solved by discretizing the computational domain with finite volume elements and using appropriate turbulence model ($k-\omega$ SST). The numerical results were obtained in terms of rotor thrust and rotor torque. The results were discussed for different wind-over-deck (WOD) angles. The results of the present study will provide information about the interaction between ship and rotor, which is to be carried out in further studies.

Keywords: Aerodynamics, Free rotor, RANS, Ship airwake, Ship-rotor interaction.

References:

1. Bardera, R. and J. C. Matías. "A comparison of helicopter recovery maneuvers on frigates by means of PIV measurements." *Ocean Engineering* 219 (2021): 108393.
2. Bardera, R., J. C. Matias-Garcia, and A. Garcia-Magariño. "PIV helicopter rotor-ground and rotor-frigate interaction study." *AIAA Aviation 2020 Forum*. 2020.
3. Crozon, C., Rene S. and G. N. Barakos. "Numerical study of helicopter rotors in a ship airwake." *Journal of Aircraft* 51.6 (2014): 1813-1832.

ICOMAA 2023

Some Results on b_3 -subbalancing Numbers

Selin Sari¹, Gül Karadeniz Gözeri²
^{1,2}Department of Mathematics, Istanbul University,
gulkaradeniz@istanbul.edu.tr
selin.sari@ogr.iu.edu.tr

Abstract

Subbalancing numbers were defined in [4] as the values of n satisfying the Diophantine equations

$$1 + 2 + \dots + (n - 1) + D = (n + 1) + \dots + (n + r)$$

for some positive integer r , where D is a fixed positive integer. In this study, we consider the b_3 -subbalancing numbers obtained by taking the D positive integer as the 3th cobalancing number. We give the recurrence relation, Binet formula and some algebraic identities of b_3 -subbalancing numbers. Furthermore, we investigate algebraic relations between the sequence of b_3 -subbalancing numbers and other integer sequences.

Keywords: Balancing numbers, Cobalancing numbers, Subbalancing numbers, Binet formula.

References:

1. A. Behera, G.K. Panda, On the square roots of triangular numbers, Fibonacci Quarterly 37(2) (1999) 98-105.
2. G.K. Panda, P.K. Ray, Some links of balancing and cobalancing numbers with Pell and associated Pell numbers, Bulletin of the Institute of Mathematics Academia Sinica(New Series) 6(1) (2011) 41-72.
3. G.K. Gozeri, A. Ozkoc, A. Tekcan, Some algebraic relations on balancing numbers, Util. Math. 103 (2017) 217-236.
4. R.K. Davala, G.K. Panda, Subbalancing numbers, Matematika 34(1) (2018) 163-172
5. S. Sari, G.K Gözeri, b_3 -Subbalancing and b_3 -Lucas subbalancing numbers, Filomat, In press.

ICOMAA 2023

Split Quaternion Sequences

Sena Nur Aktaş¹, Salim Yüce²

^{1,2}*Department of Mathematics, Yıldız Technical University,*

akts.sna@gmail.com

sayuce@yildiz.edu.tr


Abstract

In this study, we used the set of split quaternions with the metric $(+, +, -, -)$ and we worked on split quaternion sequences, concepts of their convergence and divergence. Also, we tested the convergence of split quaternion sequences. Additionally, the properties of split quaternion sequences are discussed.

Keywords: Real quaternion sequence, split quaternions, split quaternion sequences, split quaternion series.

References:

1. Couto, I. T., Lymberopoulos, A. (2021). Introduction to Lorentz geometry: curves and surfaces. CRC Press.
2. Kantor, I. L., Solodovnikov, A. S., Shenitzer, A. (1989). Hypercomplex numbers: an elementary introduction to algebras (Vol. 302). New York: Springer-Verlag.
3. Alagöz, Y., Oral, K. H., Yüce, S., Split Quaternion Matrices. Miskolc Mathematical Notes, 13 (2). pp. 223-232, 1787-2413, (2012).



ICOMAA 2023

A Bound Set of Chaotic Satellite System and Its Application in Chaos Synchronization

Ayub Khan¹, Shadab Ali², Arshad Khan³

^{1,2,3}Department of Mathematics, Jamia Millia Islamia, New Delhi, India

akhan12@jmi.ac.in

shadabbsr1@gmail.com

akhan2@jmi.ac.in

Abstract

In this manuscript, we have estimated the bound set and achieved synchronization of the chaotic satellite system. To obtain the ultimate bound set of chaotic satellite system, we have employed Lagrange optimization method. The idea of the ultimate bound has been used to the chaos control and synchronization. By Lyapunov stability theory, synchronization has been achieved between two chaotic systems via a globally exponential scheme. We use the linear feedback control and applied inequality techniques to achieve the globally exponential synchronization of two identical chaotic satellite systems. To establish the efficiency of the developed method, we have performed the numerical simulations through MATLAB.

Keywords: Ultimate bound set, Lagrange optimization, Globally exponential synchronization, Lyapunov stability theory.

MSC 2020 Classifications: 34D06, 34H10, 34D20.

References:

1. Thomas L Carroll and Louis M Pecora. Synchronizing chaotic circuits. In *Nonlinear Dynamics in Circuits*, pages 215–248. World Scientific, 1995.
2. Ayub Khan and Sanjay Kumar. Study of chaos in chaotic satellite systems. *Pramana*, 90(1):1–9, 2018.
3. Pei Wang, Damei Li, Xiaoqun Wu, Jinhua Lu, and Xinghuo Yu. Ultimate bound estimation of a class of high dimensional quadratic autonomous dynamical systems. *International Journal of Bifurcation and Chaos*, 21(09):2679–2694, 2011.
4. Guo-Ping Jiang, Guanrong Chen and Wallace Kit-Sang Tang. A new criterion for chaos synchronization using linear state feedback control. *International Journal of Bifurcation and Chaos*, 13(8):2343–2351, 2003.
5. Damei Li, Junan Lu, Xiaoqun Wu, and Guanrong Chen. Estimating the bounds for the Lorenz family of chaotic systems. *Chaos, Solitons & Fractals*, 23(2):529–534, 2005.

Optimal points in b -metric spaces endowed with graph

Shagun Sharma and Sumit Chandok

^{1,2}School of Mathematics, Thapar Institute of Engineering and Technology, Patiala- 147004, India.

shagunsharmapandit8115@gmail.com

sumit.chandok@thapar.edu

Abstract

In this paper, we focus on the best proximity point theorems to prove their uniqueness on a b -metric space endowed with a graph. We also furnish some numerical examples to support our claims. As an application of our main result, we find the solution of a nonlinear integral equation.

Keywords: best proximity point, fixed point, weak PS - property, graphical b -metric space.

References

1. A. Abkar and M. Gabeleh, The existence of best proximity points for multivalued non-self-mappings, Acad. Cienc. Exactas Fs. Nat. Ser. A Math. RACSAM 107(2013), 319 - 325.
2. M. U. Ali, T. Kamram and N. Shahzad, Best proximity point for α - ψ proximal contractive multimaps, Abstr. Appl. Anal. 2014(2014), 6 pages.
3. S. Czerwik, Nonlinear set valued contraction mappings in b -metric spaces, Atti Sem. Math Fis. Univ. Modena 46(1998), 263-276.
4. D. Singh, V. Chauhan, I. Altun, Common fixed point of power graphic (F, ψ) contraction pair on partial b -metric spaces with application, Nonlinear Anal. Model Control, 22(2017), 662 - 678.
5. M. Younis, D. Singh, A. Goyal, A novel approach of graphical rectangular b -metric spaces with an application to the vibrations of a vertical heavy hanging cable, J. Fixed point Theory Appl., 21(2019), 33.

Solution of Nonlinear Integral Equations by Generalized Contractive Condition

Shiv Kant Tiwari¹

¹Department of Mathematics,
Lukhdhirji Engineering College, Morbi, Gujarat-363642, India
shivkant.math@gmail.com

Abstract

In this paper, we prove the existence of a solution of some nonlinear integral equations with the help of common fixed-point theorems satisfying generalized contractive condition in complete metric space for two pairs of weakly compatible mappings.

Keywords: Nonlinear integral equation, Fixed point, Compatible mappings, Contractive condition.

References:

1. A. Altun, D. Turkoglu, B.E. Rhoades, Fixed points of weakly compatible maps satisfying a general contractive condition of integral type, Fixed Point Theory Appl., Vol. 2007, Article ID 17301, 9 pages; doi:10/1155/2007/17301.
2. D.P. Shukla, S.K. Tiwari, S.K. Shukla, Fixed point theorems for a pair of compatible mappings in integral type equation, Int. J. of Math. Sci. & Engg. Appls., 7(VI)(2013), 413 – 419.
3. G. Jungck, B.E. Rhoades, Fixed point theorems for occasionally weakly compatible mappings, Fixed Point Theory 7 (2006), 287-296.
4. M. Abbas, B.E. Rhoades, Common fixed point theorems for hybrid pairs of occasionally weakly compatible mappings satisfying generalized contractive condition of integral type, Fixed Point Theory Appl., Vol. 2007, Article ID 54101, doi:10/1155/2007/54101.
5. T. Abdeljawad, Fixed points for generalized weakly contractive mappings in partial metric spaces, Math. Comput. Modelling, 54(2011)2923 –2927.

ICOMAA 2023

Fractional Mathieu-Duffing System with the stability condition and chaos control

Shiva Eshaghi

Department of Basic Science, Kermanshah University of Technology, Kermanshah, Iran
sh.eshaghi@kut.ac.ir

Abstract

In this paper, we introduce the chaotic fractional Mathieu-Duffing system and state a theorem to analyze stability of the system base on the Lyapunov second method. Next, we eliminate the chaotic behaviors of the system by means of feedback controller and presented theorem. We further present numerical simulations and reveal chaotic and asymptotic stability behaviors of the system to verify the theoretical analysis.

Keywords: Chaos, Fractional Mathieu-Duffing System, Stability.

References:

1. R. Garra, R. Gorenflo, F. Polito, and Z. Tomovski, Hilfer-Prabhakar derivatives and some applications, *Applied Mathematics and Computation*, vol. 242 pp. 576-589, 2014.
2. A. A. Kilbas, M. Saigo, and R. K. Saxena, Generalized Mittag-Leffler function and generalized fractional calculus operators, *Integral Transforms and Special Function*, vol. 15 pp. 31-49, 2004.
3. Y. Lei, W. Xu, Y. Xu, and T. Fang, Chaos control by harmonic excitation with proper random phase, *Chaos Solitons and Fractals*, vol. 21, pp. 75-81, 2004.
4. I. Petras, *Fractional-order Nonlinear Systems: Modeling, Analysis and Simulation*, Springer, London, Beijing, 2011.



ICOMAA 2023

DRBEM solutions of singularly perturbed MHD flow in a square duct with variably conducting and no-slip/slipping walls

Sinem Arslan Ölçer¹, Münevver Tezer-Sezgin²

^{1,2} *Department of Mathematics, Middle East Technical University, Ankara, Turkey*

arsinem@metu.edu.tr

munt@metu.edu.tr

Abstract

In this study, the Dual Reciprocity Boundary Element Method (DRBEM) solutions of singularly perturbed magnetohydrodynamic (MHD) flow equations in a square duct with variably conducting and either slipping or no-slip walls are investigated. The MHD flow equations governed by the velocity $V(x,y)$ of the fluid and the induced magnetic field $B(x,y)$ are coupled and convection-diffusion type including the so-called Hartmann number (Ha) as a coefficient of the convection terms. When Ha is large because of the high intensity of the external magnetic field, then MHD flow equations become convection dominated. That is, the coefficients of the diffusion terms are very small giving the singularly perturbed MHD flow equations whose numerical solutions are difficult to be found especially on the thin boundary layer regions. These singularly perturbed MHD equations are solved using Shishkin mesh which consists of the transition points depending on Ha and the number of nodes taken on each side of the duct. DRBEM numerical results show that the well-known behaviors of $V(x,y)$ and $B(x,y)$ are deduced for large values of Ha such as 500, 700, and 1000. That is, the flattening flow and boundary layers formation are observed. For variably conducting and no-slip walls, for a fixed Ha and increasing wall conductivity c , the velocity $V(x,y)$ decreases in magnitude whereas the profiles of $B(x,y)$ become perpendicular to the duct walls. On the other hand, for variably conducting and slipping walls, the slip length α has such an impact on the velocity $V(x,y)$ that, its magnitude increases with an increase in α for each fixed Ha . Moreover, it is seen that the induced magnetic field $B(x,y)$ profiles is not much effected from the increase in the slip length.

Keywords: MHD flow, DRBEM, Hartmann number, Shishkin mesh, singular perturbation, transition point, slip length.

References:

1. P. W. Partridge, C. A. Brebbia, et al. Dual reciprocity boundary element method, Springer Science & Business Media, 2012.
2. N. Kopteva, E. O'Riordan, Shishkin meshes in the numerical solution of singularly perturbed differential equations, International Journal of Numerical Analysis and Modeling 7(3) (2010) 393-415.

Assessment of Bank Liquidity Risk with Artificial Neural Networks

İbrahim Demir¹, Sinem Kübra Kösen²

^{1,2}Department of Statistics, Yıldız Technical University,

idemir@yildiz.edu.tr
sinemkbr2@gmail.com

Abstract

Liquidity risk of bank is defined as the risk of not converting its short-term liabilities into cash. In this study, an estimation has been made for the current ratio, which is used as a liquidity risk indicator. A real-world case study is presented to demonstrate the applicability and exhibit efficiency, accuracy and flexibility of Artificial Neural Networks when modeling ambiguous occurrences related to bank liquidity risk measurement. To deal with these issues, we propose a model that uses Artificial Neural Networks and Time Series regression model. The implementation of these two systems comprises several algorithms and tests for validating the proposed model. In the current ratio estimation in both systems used, the results are discussed and compared. According to the results obtained time series regression model performed worse prediction than artificial neural networks.

Keywords: Current ratio, Artificial neural networks, Time series, Banking, Liquidity,

References:

1. M.Tavana, A.Reza Abtahi , D. Di Caprio , M. Poortarigh (2017). An Artificial Neural Network and Bayesian Network Model for Liquidity Risk Assessment in Banking
2. Ahmed N., Ahmed Z., Naqvi I.H. (2011). Liquidity Risk and Islamic Banks: Evidence from Pakistan. Interdisciplinary Journal of Research in Business, 1(9), 99-102.
3. Arif, A., Anees, N.A. (2012). Liquidity Risk and Performance in the Banking System. Journal of Financial Regulation and Compliance, 20(2), 182-195.
4. Basel Committee on Banking Supervision (BCBS) (2008). Principles for sound liquidity risk management and supervision. September, No.144. Bank of International Settlements, available at <http://www.bis.org/publ/bcbs144.pdf>
5. Basel Committee on Banking Supervision (BCBS) (2010). Basel III: International Framework for Liquidity Risk Measurement, Standards and Monitoring. December, No.188. Bank for International Settlements, available at <http://www.bis.org/publ/bcbs188.pdf>

ICOMAA 2023

On Modification of Hirota Method Through Transition Between Parameters

Sukri Khareng¹, Ömer Ünsal²

^{1,2}Department of Mathematics and Computer Sciences, Eskisehir Osmangazi University,

sukriyou99@gmail.com

ounsal@ogu.edu.tr

Abstract

In this study, three different classes of solutions to a specific form of a new generalized fourth-order nonlinear partial differential equation are presented. This process appears as result of transition between real and complex parameters which provides an useful modification of well known Hirota method [1,2,3]. What enable us to obtain various solutions are restrictions on phase shifts.

Keywords: Modification of Hirota method, bilinear form, complexiton solutions, soliton solutions, soliton-complexiton solutions.

References:

1. Wazwaz A.M., New solutions for two integrable cases of a generalized fifth-order nonlinear equation. Modern Physics Letters B. Vol. 29, (2015) 1590065 (9 pages).
2. Wazwaz A.M., Zhaqilao. Nonsingular complexiton solutions for two higher-dimensional fifth-order nonlinear integrable equations. Physica Scripta. Vol. 88, (2013) 025001 (6 pages).
3. Ünsal Ö., Complexiton solutions for (3+1) dimensional KdV-type equation. Computers and Mathematics with Applications. Vol. 75, (2018) 2466-2472.

ICOMAA 2023

Statistical Convergence of Sequences of Sets of Order (α, β)

Sunil K. Sharma

*School of Mathematics, Shri Mata Vaishno Devi University, Kakryal,
Katra-182320, Jammu and Kashmir, India*

sunilksharma42@gmail.com

Abstract

In this paper we define Wijsman λ - statistical convergence of order (α, β) and studied some properties of this concept. We also make an effort to define Wijsman λ - statistical convergence of Musielak-Orlicz function of order (α, β) and examine some topological properties.

Keywords: Statistical convergence, λ -statistical convergence, Wijsman convergence, Musielak-Orlicz function.

References:

1. Qing-Bo Cai, S. K. Sharma, and M. A. Mursaleen, A note on lacunary sequence spaces of fractional difference operator of order (α, β) , Journal of Function Spaces, Vol. 2022, Article ID 2779479, 9 pages.
2. M. Et, S. A. Mohiuddine, and A. Alotaibi, On λ -statistical convergence and strongly λ - summable functions of order α , Jour. Ineq. Appl., vol. 2013, no. 469, 8 pages, 2013.
3. B. Hazarika and A. Esi, On λ -asymptotically Wijsman generalized statistical convergence of sequences of sets, Tatra Mountains Mathematical Publications-Number Theory, vol. 56,(2013), 67-77.
4. B. Hazarika and A. Esi, Statistically almost λ -convergence of sequence of sets, European Journal of Pure and Applied Mathematic, vol. 6, no. 2,(2013), 137-146.
5. F. Nuray and B. E. Rhoades, Statistical convergence of sequences of sets, Fasciculi Mathematici, vol. 49, (2012), 1-9.

ICOMAA 2023

Prediction of HIV-1 Nucleoside Reverse Transcriptase Resistance with Artificial Neural Networks

Sumeyye Yilmaz¹, Murat Sari², Huseyin Tunc³

¹Department of Mathematics, Yildiz Technical University,

²Department of Mathematics Engineering, Istanbul Technical University,

³Department of Biostatistics and Medical Informatics, Bahcesehir University

sumeyye.yilmaz1@std.yildiz.edu.tr

muratsari@itu.edu.tr

huseyin.tunc@bau.edu.tr

Abstract

HIV is a serious and chronic disease that affects millions of people worldwide. Despite improvements in life expectancy and quality of life for patients receiving antiretroviral therapy, drug resistance development remains a major challenge for effective treatment. Understanding the mechanism of resistance is crucial, as drug resistance is currently the biggest obstacle to the effective treatment of HIV/AIDS. To overcome these challenges, researchers use genotype and phenotype testing, which can be both time-consuming and expensive. Machine learning models are one of the methods used to overcome these challenges by interpreting the available data appropriately. The purpose of this study is to develop artificial neural network (ANN) models for predicting the resistance profiles of nucleoside reverse transcriptase inhibitors (NRTIs) using the Stanford HIV drug resistance database. The Pearson correlation coefficient and mean square error metrics are presented to measure the predictive performances of various ANN models.

Keywords: Artificial neural networks, HIV-1, Drug resistance, Machine learning.

References:

1. Tunc, H., Dogan, B., Darandeli Kiraz, B. N., Sari, M., Durdagi, S., & Kotil, S. (2023). Prediction of HIV-1 protease resistance using genotypic, phenotypic, and molecular information with artificial neural networks. *PeerJ* 11:e14987.
2. Amamuddy, O. S., Bishop, N. T., & Bishop, O. T. (2017). Improving fold resistance prediction of HIV-1 against protease and reverse transcriptase inhibitors using artificial neural networks. *BMC Bioinform*, 369-376.
3. Steiner, M. C., Gibson, K.M., & Crandall, K. A. (2020). Drug resistance prediction using deep learning techniques on HIV-1 sequence data. *Viruses*, 12(5).
4. Rhee, S.Y., Taylor, J., Wadhera, G., Ben-Hur, A., Brutlag, D. L., Shafer, R.W. (2006) Genotypic predictors of human immunodeficiency virus type 1 drug resistance. *Proc Natl Acad Sci USA*, 103(46):17355–60.

Proposing a New DNA and Music Powered Cryptology Technique

Muharrem Bayrak¹, Halil Yakıt², Kayra Ege Altun³, Şerif Efe Dartar⁴

^{1,2,3,4}Kabatas Erkek High School,

Beşiktaş, İstanbul, Türkiye

muharrembayrak26@gmail.com

halilyakit@gmail.com

altunkayra07@gmail.com

dserif.efe@gmail.com

Abstract

In today's completely digitalized world, cryptology techniques both in cryptography and cryptanalysis are needed. Thus, in this paper, the connection between biology, music, and cryptology was examined and three algorithms that can be used both in encryption and decryption processes were proposed.

The process lying behind these three algorithms is a process that matches every character to a three-digit DNA sequence which consists of 'A', 'T', 'G', and 'C' letters. After that, every organic base is matched to three notes. The first algorithm relies on a key that consists of 9 decimal digits. Plaintext is reflected according to this key. The second algorithm has a circular technique and with this technique order of the notes is changed. The third algorithm is a combination of these two algorithms. With these three algorithms, the cryptanalysis process (especially against brute force attacks) was hardened.

Keywords: Cryptology, Cryptography, DNA, Music

References:

1. Zengin, M. (2021). Genetik Kod Yöntemi ile Kriptoloji Uygulaması (Yüksek Lisans Tezi). Karabük Üniversitesi, Karabük.
2. Dutta, S., Kumar, C., & Chakraborty, S. (2013). A symmetric key algorithm for cryptography using music. *International Journal of Engineering and Technology*, 5(3), 3109-3115.
3. Yunpeng, Z., Yu, Z., Zhong, W., & Sinnott, R. O. (2011, October). Index-based symmetric DNA encryption algorithm. In 2011 4th International Congress on Image and Signal Processing (Vol. 5, pp. 2290-2294). IEEE.

A Study on T-Magnetic Curves

Şerife Nur Bozdağ¹, Şevval Kublay²
^{1,2} Department of Mathematics, Ege University,
serife.nur.yalcin@ege.edu.tr
sevvalkublay@hotmail.com

Abstract

In this study we handled T-magnetic curves in three dimensional Kenmotsu manifolds. We obtained necessary and sufficient conditions of T-magnetic and T-magnetic Legendre curves to be biharmonic, f-harmonic and f-biharmonic. We have also introduced some non-existence theorems.

Keywords: T-magnetic curves, Legendre curves, Kenmotsu manifolds, harmonic maps.

References:

1. Özdemir, Z., Gök, I., Ekmekci, F.N. Yaylı, Y. , Notes on magnetic curves in 3D semi Riemannian manifolds, Turkish Journal of Mathematics, 39, (2015) 412-426.
2. Calin C., Crasmareanu M., Munteanu M.I., Slant curves in three-dimensional f -Kenmotsu manifolds, J. Math. Anal. Appl., 394, (2012) 400-407
3. Perkaş S. Y., Acet B. E., Ouakkas S., On Biharmonic and Biminimal Curves in 3-dimensional f-Kenmotsu Manifolds, Fundamentals of Contemporary Math. Sciences, 1(1), (2020) 14-22.
4. Kenmotsu K., A class of almost contact Riemannian manifolds, Tohoku Mathematical Journal, Second Series 24(1) (1972) 93-103.
5. Bozdağ, Ş. N., Erdoğan, F. E., f-Biharmonic and bi-f-harmonic magnetic curves in three-dimensional normal almost paracontact metric manifolds, International Electronic Journal of Geometry 14.2 (2021) 331-347.

ICOMAA 2023

Solvability in the small and Schauder-type estimates for higher order elliptic equations in Banach-Sobolev spaces

Bilal Bilalov^{1,2}, Yusuf Zeren¹, Sabina Sadigova^{2,3}, Seyma Cetin¹
¹Department of Mathematics, Yildiz Technical University,
²Institute of Mathematics and Mechanics of NAS of Azerbaijan, Baku, Azerbaijan,
³Khazar University, Baku, Azerbaijan
bilal.bilalov@yildiz.edu.tr
yzeren@yildiz.edu.tr
s_sadigova@mail.ru
seyymacetin@gmail.com

Abstract

In this work it is considered an elliptic operator L of m -th order with nonsmooth coefficients in Banach-Sobolev function spaces on a bounded domain. Under weaker restrictions on the coefficients of the operator, we prove the local solvability (in the strong sense), and also establish interior Schauder-type estimates for Banach-Sobolev function spaces.

Keywords: elliptic equations, Banach-Sobolev function spaces, local solvability, Schauder-type estimates.

Acknowledgement: This work is supported by TÜBİTAK 2211-A Domestic General Doctorate Scholarship Program.

References:

1. Bilalov, B.T., Sadigova, S. 2021. "On the Fredholmness of the Dirichlet Problem for a second-order Elliptic Equation in grand-Sobolev Spaces", *Ricerche Di Matematica*, 1-40.
2. Bilalov, B.T., Sadigova, S.R. 2021. "On Solvability in the small of higher order Elliptic Equations in grand-Sobolev Spaces", *Complex Variables and Elliptic Equations*, 66(12), 2117-2130.
3. Bilalov, B.T., Sadigova, S., Zeren, Y., Cetin S. 2022. "On Solvability in the small and Schauder-type Estimates for higher order Elliptic Equations in grand-Sobolev Spaces(nonseparable case)", *Applicable Analysis*, 1-14.
4. Bilalov, B.T., Zeren, Y., Sadigova, S., Cetin S. 2022. "Solvability in the small of m -th order Elliptic Equations in weighted grand-Sobolev Spaces", *Turkish Journal of Math*, 46(6), 2078-2095.

ICOMAA 2023

On Left Derivations of Generalized Matrix Algebras

Seymanur Aydin¹, Nurcan Argac²
^{1,2}Department of Mathematics, Ege University,

seymanuraydiinn@gmail.com
nurcan.argac@ege.edu.tr

Abstract

Let $U = \begin{pmatrix} A & M \\ N & B \end{pmatrix}$ be a generalized matrix algebra defined by the Morita context $(A, B, M, N, \Phi_{MN}, \Psi_{NM})$ ([5]). Let R be a ring and M a left R -module. An additive map $D : R \rightarrow M$ is called a left derivation if $D(xy) = xD(y) + yD(x)$ is fulfilled for all $x, y \in R$. The concept of left derivation was introduced by Brešar and Vukman in [2]. They proved every left derivation on a semiprime ring R is a derivation which maps R into its center. Furthermore, they also show that if A is a Banach algebra, then every continuous left derivation $D : A \rightarrow A$ maps A into its radical. The question under which conditions left derivation is zero on a given Banach algebra have attracted much attention of authors (for instance, see [1],[2],[4]). Recently in [3] Ghosseiri proved that if R is a 2-torsion free ring with identity, then any left derivation on the full matrix ring $M_n(R)$, ($n \geq 2$) is identically zero. In this talk we consider the problem of describing the form of left derivation of U and we show that under the certain conditions any left derivation on generalized matrix algebra is zero. Finally as its applications, we prove that the similar conclusion remains valid on full matrix algebras, unital prime rings with a nontrivial idempotents, unital standard operator algebras, triangular algebras and nest algebras.

References:

1. Ali S. On generalized left derivations in rings and Banach algebras. Aequationes Math. 2011;81:209-226.
2. Brešar M, Vukman J. On left derivations and related mappings. Proc. Amer. Math. Soc. 1990;110(1):7-16.
3. Ghosseiri N. M. On Jordan left derivations and generalized Jordan left derivations of matrix rings. Bull. of Iranian Math. Soc. 2012;38(3):689-698.
4. Jung Y. S. On left derivations and derivations of Banach algebras. Bull. Korean Math. Soc. 1998;35:659-667.
5. Morita K. Duality for modules and its applications to the theory of rings with minimum condition. Sci. Rep. Tokyo Kyoiku Daigaku Sect. 1958;A6:83-142.

ICOMAA 2023

Results on Eigenvalue Problems of NonLinear Conformable Fractional Differential Equations

Şuayip Toprakseven¹

¹Artvin Vocational School, Accounting and Taxation, Artvin Çoruh University, Artvin, 08100, Turkey
topraksp@artvin.edu.tr

Abstract

The basic theory on nonlinear conformable fractional boundary value problems is discussed. Necessary conditions for the existence of eigenfunctions are provided with the help of the maximum principle. Further, we present lower and upper bounds for the eigenvalues. A general existence and uniqueness result is obtained using the method of lower and upper solution. Two monotone sequences of lower and upper solutions that convergence to the exact solution of the problem are derived.

Keywords: Fractional differential equations, Maximum principle, Lower and upper solutions, Conformable fractional derivative.

References:

1. R. P. Agarwal, N. Benchohra, and S. Hamani. Boundary-value problems for Fractional Differential Equations, Georgian Mathematical Journal, 16 (2009), 401-411.
2. M. Al-Refai, M. Hajji. Monotone iterative sequences for nonlinear boundary-value problems of fractional order, Nonlinear Analysis Series A: Theory, Methods and Applications 74 (2011), 3531-3539.
3. A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo. Theory and Applications of Fractional Differential Equations, North-Holland Mathematics Studies, vol. 204, Elsevier, Amsterdam, The Netherlands, 2006.
4. Jiangang Qi and Shaozhu. Eigenvalue problems of the model from nonlocal continuum mechanics, Journal of Mathematical Physics, 52 073516(2011).

ICOMAA 2023

On Solution of Some Differential Equations Using Bicomplex Transformations

Serpil Halıcı¹, Sule Curuk²

^{1,2}Department of Mathematics, Pamukkale University,

shalici@pau.edu.tr

sule9220@gmail.com

Abstract

In this study, motivated by studies on integral transformations in bicomplex space, we derived a formula for bicomplex version of the Hartley transform and investigated some fundamental properties of bicomplex Hartley transform by this formula. Moreover, we gave a solution of the Bessel differential equation using bicomplex Hartley transformation.

Keywords: Integral transformations, Bicomplex functions, Bessel equation.

References:

1. H. O. Beca, An orthogonal set based on bessel functions of the first kind/skup ortogonalnih funkcija zasnovan na beselovim funkcijama prvevrste, Publikacije Elektrotehnickog fakulteta., Serija Matematika i fizika, no. 678/715 (1980)85–90.
2. M. P. Goswami, Bicomplex integral transforms and applications, Ph.D.thesis MNIT Jaipur, (2017).
3. S. Halıcı and S. Curuk, On integral transforms of some special functions, in: Conference Proceedings of Science and Technology, 3(2)(2020) 215-221.
4. R. V. Hartley, A more symmetrical fourier analysis applied to transmissionproblems, Proceedings of the IRE, 30(3) (1942). 144-150.
5. M. E. Luna-Elizarraras, M. Shapiro, D. C. Struppa and A. Vajiac, Bicomplex numbers and their elementary functions, Cubo (Temuco), 14(2) (2012) 61-80.

ICOMAA 2023

Graphs of Bitonic Algebras

Refet Polat¹, Şule Ayar Özbal² and Saadet Eskiizmirliler³

¹Department of Mathematics, Yaşar University,
refet.polat@yasar.edu.tr

²Department of Mathematics, Yaşar University,
sule.ayar@yasar.edu.tr

³Coordinator of Central Student Activities Unit, Yaşar University,
saadet.eskiizmirliler@yasar.edu.tr


Abstract

We introduce the concept of graphs associated with commutative bitonic-algebras, which is a bitonic-graph whose vertices are the elements of commutative bitonic-algebra and whose edges are the association of two vertices, that is two elements from commutative bitonic-algebra.

Keywords: Bitonic algebras, graphs, subgraphs, filters.

References:

1. M. A. Ansari, A. Haidar and A. N.A. Koam, On a Graph Associated to UP-Algebras, Math. Comput. Appl. 2018, 23(4), 61.
2. N. Kırçalı Gürsoy, Graphs of Wajsberg Algebras via Complement Annihilating, Symmetry 2023, 15, 121.
3. D. Al-Kadi Quliyeva, Distinguishing Graphs Associated to KU-Algebras using Graph Invariants, Appl. Math. Inf. Sci. 15, No. 1, 17-22 (2021)



ICOMAA 2023

Inequalities of Hermite-Hadamard-Mercer Type for Twice Differentiable with Applications

Francisco Martínez González¹, Muhammad Zafar Iqbal², Talib Hussain³
¹Department of Applied Mathematics and Statistics, Technical University of Cartagena, Spain,
f.martinez@upct.es
^{2,3}Department of Mathematics and Statistics, University of Agriculture Faisalabad, Pakistan,
mzts2004@hotmail.com
talibnazir76@mail.com

Abstract

In this paper, authors use Riemann-Liouville (R-L) fractional integrals to derive a useful lemma for twice differentiable functions. By using the derived lemma, we establish numerous fractional integral inequalities associated to Hermite-Hadamard (H-H) and Hermite-Hadamard-Mercer (H-H-M) type inequalities for twice differentiable in absolute values are convex. Some of these results also generalize findings from earlier research. Additionally, we can observe the efficiency of our inequalities through some applications on special means.

Keywords: Riemann-Liouville fractional integrals, Hermite-Hadamard inequality, Hermite-Hadamard-Mercer inequality, Jensen-Mercer Inequality, Convex function.

References:

1. Podlubny, I. (1999). An introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications. *Math. Sci. Eng*, 198, 340.
2. Hadamard, J. (1893). Study on the properties of entire functions and in particular of a function considered by Riemann. *Journal of Pure and Applied Mathematics*, 9, 171-215.
3. Ögülmüş, H., & Sarıkay, Z. M. (2021). Hermite-Hadamard-Mercer type inequalities for fractional integrals. *Filomat*, 35(7), 2425-2436.
4. Mercer, A. M. (2003). A variant of Jensen's inequality. *J. Inequal. Pure Appl. Math*, 4(4), 73.

ICOMAA 2023

Applied Mathematical Physics: Solving Real-World Problems

Seyyed Ali Mohammadiyeh¹ and Hamid Purhasani²

¹*Department of Pure Mathematics, Faculty of Mathematical Sciences,
University of Kashan, Kashan, I. R. Iran
alim@kashanu.ac.ir*

²*Department of Computer Engineering, Faculty of Engineering,
University of Yazd, Yazd, I. R. Iran
9424303@yazd.ac.ir*

Abstract

Mathematical physics is a field that has made significant contributions to our understanding of the physical world. The interplay between mathematics and physics has allowed us to model and solve real-world problems in a wide range of areas, from understanding the behavior of particles and systems at the quantum level to predicting the dynamics of astronomical objects and understanding the behavior of complex systems in engineering and other applied fields.

Keywords: Mathematical physics, Mathematics, Physics, Applied mathematics, Real-world problems.

References:

1. J. C. Maxwell, *A Treatise on Electricity and Magnetism* (Oxford University Press, 1873).
2. S. L. Sobolev, *Applications of Mathematics in Natural Sciences* (Springer, 1975).
3. R. Penrose, *The Emperor's New Mind: Concerning Computers, Minds, and The Laws of Physics* (Oxford University Press, 1989).
4. M. L. Boas, *Mathematical Methods in the Physical Sciences* (John Wiley & Sons, 2006).
5. S. K. Donaldson and P. B. Kronheimer, *The Geometry of Four-Manifolds* (Oxford University Press, 1990).

Soft Fractional Ideals and Soft Star-Operations of Integral Domains

Tasaduk Rashid Mir
Department of Mathematics
Jamia Millia Islamia
New Delhi – 110025, India
Email: tasadukmr111@gmail.com

Abstract

The concept of soft modules was introduced by Q. M. Sun et al and carried out several properties of soft modules by utilising Molodtsov's concept of soft set theory and modules. In this paper, we demonstrate the operations from the set of soft fractionary ideals into itself. We also introduce the concept of soft star-operations on an integral domain and show that the set of all soft star-operations on the integral domain forms a complete lattice. We introduce a concept of valuation rings in terms of soft modules of the fraction field and demonstrate the equivalence of certain soft modules of the fraction field, called soft fractional ideals.

Keywords: Soft module, Valuation rings, Soft star-operation, Soft ν -ideal, Invertible soft fractional ideal.

2010 AMS Subject Classification: 13A15, 13G05, 03E75.

References:

1. Pradip Kumar Maji, Ranjit Biswas, and A Ranjan Roy. Soft set theory. *Computers & Mathematics with Applications*, 45(4-5):555–562, 2003.
2. Abdelghani Taouti and Waheed Ahmad Khan. A note on some new soft structures. *J. Math. Comput. Sci.*, 4(3):522–530, 2014.
3. Abdelghani Taouti, Waheed Ahmad Khan, and Seema Karkain. Note on soft fractional ideal of ring. *J. Math. Comput. Sci.*, 8(5):579–583, 2018.
4. D.D. Anderson, D.F. Anderson, Examples of star-operations on integral domains, *Comm. Algebra* 18 (1990) 1621–1643.

Nabla Caputo Type q -fractional difference operators and their properties

Taylan Demir¹, Mohammed Esmael Samei²

¹Department of Mathematics, Çankaya University,

²Department of Mathematics, Bu-Ali Sina University

demir.taylan96@gmail.com

mesamei@gmail.com

Abstract

In this section, initially we will mention that the definitions of fractional calculus. To give a detail as, Grunwald-Letnikov, Riemann-Liouville & Caputo's definitions will be given because, these definitions are connected with the difference operators. So, after these definitions are introduced, difference equations and linear type of difference equations will be given in this part. Later, definitions of fractional calculus that are our mentioning will be adapted in q -version as Riemann-Liouville and Caputo type on difference operators.

Keywords: Difference operators, q -Mittag Leffler functions, fractional operators, singular and non-singular kernel

References:

1. Sun, Z. Z., & Gao, G. H. (2020). Fractional differential equations. In Fractional Differential equations. De Gruyter.
2. Diethelm, K., & Ford, N. J. (2002). Analysis of fractional differential equations. *Journal of Mathematical Analysis and Applications*, 265(2), 229-248.
3. Annaby, M. H., & Mansour, Z. S. (2012). q -Fractional calculus and equations (Vol.2056). Springer.
4. Abdeljawad, T., B., & Baleanu, D. (2012, January). A generalized q -Mittag Leffler function by q -Caputo fractional linear equations. In *Abstract and Applied Analysis* (Vol. 2012). Hindawi.
5. Samei, M. E., Zanganeh, H. & Aydogan, S. M. (2021). Investigation of a Class of the Singular Fractional Integro-differential Quantum Equations with Multi-Step Methods. *Journal of Mathematical Extension* (Vol. 15), 1-54. <https://doi.org/10.30495/JME.SI.2021.2070>

ICOMAA 2023

Applications of q-Mittag Leffler function on q-fractional differential operators

Taylan Demir¹, Mohammad Esmael Samei²

¹Department of Mathematics, Çankaya University,

²Department of Mathematics, Bu-Ali Sina University

demir.taylan96@gmail.com

mesamei@gmail.com

Abstract

In this part, we will introduce definitions of special functions. Gamma, beta and Mittag-Leffler functions will be given. Then, derivatives and integration of Mittag-Leffler function will be defined before definitions of fractional operators will have been mentioned. Since, this topic are basic to introduce q-Mittag Leffler function and their applications on q-fractional differential operators. Finally, q-gamma, q-beta and q-Mittag Leffler functions will be mentioned before q-fractional differential operators.

Keywords: q-gamma function, q-beta function, Mittag-Leffler kernel, singular kernel, q-fractional operators, beta function, gamma function, fractional operators

References:

1. Milici, C., Drăgănescu, G., & Machado, J. T. (2018). *Introduction to fractional differential equations* (Vol. 25). Springer.
2. Annaby, M. H., & Mansour, Z. S. (2012). *q-Fractional calculus and equations* (Vol. 2056). Springer.
3. Abdeljawad, T., Benli, B., & Baleanu, D. (2012, January). A generalized q-Mittag-Leffler function by q-Caputo fractional linear equations. In *Abstract and Applied Analysis* (Vol. 2012). Hindawi.
4. Hajiseyedazizi, S. N., Samei, M. E., Alzabut, J. & Chu, Y. (2021). On multi-step methods for singular fractional q -integro-differential equations. *Open Mathematics*, *19*, 1378–1405. <https://doi.org/10.1515/math-2021-0093>
5. Samei, M. E., Baleanu, D. & Rezapour, S. (2020). An increasing variables singular system of fractional q -differential equations via numerical calculations. *Advances in Difference Equations*, *2020*, 452. <https://doi.org/10.1186/s13662-020-02913-5>

ICOMAA 2023

Applications of q-Mittag Leffler function on q-fractional differential operators

Taylan Demir¹, Mohammad Esmael Samei²

¹Department of Mathematics, Çankaya University,

²Department of Mathematics, Bu-Ali Sina University

demir.taylan96@gmail.com

mesamei@gmail.com

Abstract

In this part, we will introduce definitions of special functions. Gamma, beta and Mittag-Leffler functions will be given. Then, derivatives and integration of Mittag-Leffler function will be defined before definitions of fractional operators will have been mentioned. Since, this topic are basic to introduce q-Mittag Leffler function and their applications on q-fractional differential operators. Finally, q-gamma, q-beta and q-Mittag Leffler functions will be mentioned before q-fractional differential operators.

Keywords: q-gamma function, q-beta function, Mittag-Leffler kernel, singular kernel, q-fractional operators, beta function, gamma function, fractional operators

References:

1. Milici, C., Drăgănescu, G., & Machado, J. T. (2018). *Introduction to fractional differential equations* (Vol. 25). Springer.
2. Annaby, M. H., & Mansour, Z. S. (2012). *q-Fractional calculus and equations* (Vol. 2056). Springer.
3. Abdeljawad, T., Benli, B., & Baleanu, D. (2012, January). A generalized q-Mittag-Leffler function by q-Caputo fractional linear equations. In *Abstract and Applied Analysis* (Vol. 2012). Hindawi.
4. Hajiseyedazizi, S. N., Samei, M. E., Alzabut, J. & Chu, Y. (2021). On multi-step methods for singular fractional q -integro-differential equations. *Open Mathematics*, *19*, 1378–1405. <https://doi.org/10.1515/math-2021-0093>
5. Samei, M. E., Baleanu, D. & Rezapour, S. (2020). An increasing variables singular system of fractional q -differential equations via numerical calculations. *Advances in Difference Equations*, *2020*, 452. <https://doi.org/10.1186/s13662-020-02913-5>

ICOMAA 2023

On the basis property of eigenfunctions of the indefinite Sturm-Liouville problem in Lebesgue spaces

Telman Gasymov¹ and Ulker Hashimova¹

¹Baku State University, Baku, Azerbaijan,
telmankasumov@rambler.ru

Abstract

We consider the spectral problem

$$\begin{aligned} -y'' + p(x)y &= \lambda r(x)y, \quad x \in \bigcup_{s=1}^l (x_{s-1}, x_s), \quad (1) \\ y(0) = y(1) &= 0 \\ \begin{cases} y(x_s + 0) - a_s y(x_s - 0) = 0 \\ y'(x_s + 0) - b_s y'(x_s - 0) = c_s y(x_s - 0), \end{cases} \quad s = \overline{1, l-1}, \quad (3) \end{aligned} \quad (2)$$

where $0 = x_0 < x_1 < \dots < x_l = 1$, $p(x) \in L_1(0,1)$, $r(x) = r_s$ for $x \in (x_{s-1}, x_s)$, $r_s \in \mathbb{R} \setminus \{0\}$, $a_s, b_s, c_s \in \mathbb{C}$, $s = \overline{1, l}$. Problem (1)-(3) has two series of eigenvalues $\{\lambda_n^\pm\}_{n \in \mathbb{N}}$ (see [1]), they correspond to eigenfunctions $\{y_n^\pm\}_{n \in \mathbb{N}}$. In this paper, we prove that the system $\{y_n^\pm\}_{n \in \mathbb{N}}$ forms a basis in the space $L_p(0,1)$, $1 < p < \infty$, equivalent to the system $\{\sin \pi n x\}_{n \in \mathbb{N}}$. In particular, for $p = 2$ this basis is a Riesz basis in $L_2(0,1)$.

Note that in the case when instead of conjugation conditions (3) the conditions of continuity of the functions themselves and their derivatives at the points x_s are taken, similar results were obtained in [2-4].

Keywords: indefinite Sturm-Liouville problems, eigenfunctions, basicity, Riesz basis.

References:

1. Gasymov T. B., Hashimova U. G., Gurbanova N. J. On the Distribution of the Eigenvalues Discontinuous Sturm-Liouville Operators with Indefinite Weight Functions. 5th International Conference ICOMAA2022, May 11-14 2022, Yildiz Technical University, Turkey. P.186.
2. Benzinger H.E. Sturm-Liouville Problem with Indefinite Weight Functions in Banach Spaces. Proceedings of the 1984 Workshop Spectral Theory of Sturm-Liouville Differential Operators. Held at Argonne National Laboratory, May 14- June 15, 1984. ANL-84-73. P.47-55.
3. Eberhard W., Freiling G., Schneider A., Expansion Theorems for a Class of Regular Indefinite Eigenvalue Problems, Differential and Integral Equations, Vol. 3, No 6 (1990), 1181-1200.
4. Freiling G. and Kaufmann F.J., On uniform and L_p -convergence of eigenfunction expansions for indefinite eigenvalue problems, Integral Equations and Operator Theory Vol. 13(1990), 193-215.

Description of domains of definition of fractional powers of a second –order differential operator with integral boundary conditions

Telman Gasymov¹ and Reyhan Taghiyeva¹

¹Department of Mechanics- Mathematics, Baku State University,
telmankasumov@rambler.ru
reyhanabasli2015@gmail.com

Abstract

We consider a differential operator L generated in the space $L_p(0,1)$, $1 < p < \infty$, differential expression

$$l(y) = p_0(x)y'' + p_1(x)y' + p_2(x)y, \quad x \in (0,1), \quad (1)$$

and integral boundary conditions

$$U_\nu(y) = \sum_{j=0}^2 \int_0^1 g_{\nu j}(x)y^{(2-j)}(x)dx = 0, \quad \nu = 1,2, \quad (2)$$

where $p_0(x) > 0$, $p_0(x) \in W_1^1(0,1)$, $p_1(x), p_2(x) \in L_1(0,1)$, $g_{\nu j}(x) \in C[0,1]$, $\nu = 1,2, j = \overline{0,2}$.

Under the assumption of regularity of any conditions (see. [1]), the positivity of the operator L is proved and an estimate for the complex powers of the operator L is obtained. Depending on the regularity condition, Banach space X is chosen, which either coincides with the space $L_p(0,1)$ or is a subspace of finite codimension. Using the results of [2], we describe the domains of definition of the fractional powers of the operator L with the help of interpolation spaces corresponding to the complex method. Similar results in some special cases were obtained in [3].

Keywords: domains of fractional power, regularity conditions, integral boundary conditions, interpolation spaces, complex method.

References:

1. Gasymov T.B., Taghiyeva R.J. On fractional powers of the second-order differential operators. Materials of the National Scientific Conference "Theory of Functions, Functional Analysis and their Applications", November 28-29, 2022 Baku. P.199.
2. R.Seeley, Norms and domains of complex powers A_B^Z , Amer.J.Math., V.93 (1971), 299-309.
3. J.Gallardo, Second-Order Differential Operators with Integral Boundary Conditions and Generation of Analytic Semigroups, Rocky Mountain J. Math., v.30, no4 (2000), 1265-1291.

Weighted Hardy's Inequality Revisited

Tuğçe Ünver Yıldız¹, Uğur Can Yıldırım²

^{1,2}Department of Mathematics, Kirikkale University

tugceunver@kku.edu.tr

ucy9292@gmail.com

Abstract

The characterization of weighted Hardy inequalities is well known. There are numerous proofs that combine Hölder's inequality, Fubini Theorem, Minkowski's integral inequality with discretization and duality techniques. We will provide a straightforward and basic proof of an equivalent condition that is motivated by some recent results on the proof of the two-weight Hardy inequality.

Keywords: two-weight Hardy inequality, equivalent conditions

References:

1. A. Gogatishvili and L. Pick, The two-weight Hardy inequality: a new elementary and universal proof, arXiv preprint arXiv:2109.15011 (2021) (to appear in Proc. Amer. Math. Soc.)
2. A. Kufner, L.E. Persson and N. Samko, Weighted inequalities of Hardy type. Second edition. *World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ*, 2017. xx+459 pp.
3. B. Opic and A. Kufner, Hardy-type inequalities. Pitman Research Notes in Mathematics Series, 219. Longman Scientific & Technical, Harlow, 1990. xii+333 pp

ICOMAA 2023

Sequential T_3 (regular Hausdorff) Spaces

Tunçar ŞAHAN¹, Elanur ÖZCAN²

^{1,2}*Department of Mathematics, Aksaray University,*

tuncarsahan@gmail.com

ozcanelanur55@gmail.com

Abstract

The notion of sequential open set is a generalization of the notion open set in topological spaces since every open set is in fact a sequential open set. According to this, in 2019, Akız and Koçak introduced the notion of sequentially Hausdorff space which also a generalization of Hausdorff spaces. That is, every Hausdorff space is a sequentially Hausdorff space. It is very interesting a significant part of the properties provided by Hausdorff spaces are also provided for sequential Hausdorff spaces.

In this study, we introduce the notion of a sequentially T_3 -space and investigate its properties. We also give the relation between sequentially T_3 property and other well-known separation properties.

Keywords: Sequential open set, Regular Hausdorff (T_3) space, Sequential space.

References:

1. H. F. Akız and L. Koçak, Sequentially Hausdorff and full sequentially Hausdorff spaces, Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics, 68(2) (2019) 1724-1732.
2. R. Brown, Topology and groupoids, Booksurge PLC, 2006.
3. H. Çakallı, Sequential definitions of compactness, Applied Mathematics Letters., 21(6) (2008) 594-598.

ICOMAA 2023

Some Basic Results on Groups up to Congruence Relations

Tunçar ŞAHAN¹, Ahmet Eren KILIÇ²
^{1,2}Department of Mathematics, Aksaray University,
tuncarsahan@gmail.com
kilicahmet383@gmail.com

Abstract

It is a well-known fact crossed modules and group-groupoids, i.e. strict categorical groups, are categorically equivalent algebraic structures. In 2020, Datuashvili et al. aim to obtain for categorical groups an analogous description in terms of certain crossed module type objects as strict categorical groups. Thus, they introduced the notion of c-group or also known as group up to congruence relation.

In this study, we explore most significant properties of groups up to congruence relations. Moreover, we give different definitions for homomorphisms, kernels, and images up to congruence relations.

Keywords: Group, Congruence relation, Kernel, Image.

References:

1. R. Brown and C.B. Spencer, G-groupoids, crossed modules and the fundamental groupoid of a topological group, Proc. Konn. Ned. Akad. v. Wet., 79 (1976) 296–302.
2. T. Datuashvili, O. Mucuk and T. Şahan, Groups up to congruence relation and from categorical groups to c-crossed modules, J. Homotopy Relat. Struct., 15 (2020) 625–640.
3. A. Joyal and R. Street, Braided tensor categories, Adv. Math. 102 (1993) 20–78.

ICOMAA 2023

On boundedness of the Riesz potential and its commutators in local Orlicz-Morrey type spaces

Mendybay Turgumbayev¹ and Aidos Adilkhanov²
¹Department of Mathematics, Karaganda University, Kazakhstan

mentur60@mail.ru

²Eurasian national University, Astana, Kazakhstan

adilkhanov_kz@mail.ru

Abstract

In this paper, we shall give sufficient conditions for the boundedness of the Riesz potential and its commutator in local Orlicz-Morrey type spaces. We present Adams-type results for the Riesz potential and its commutator in local Orlicz-Morrey type spaces. Characterisation for the Riesz potential and its commutators on generalized Orlicz-Morrey spaces were obtained in [1].

Keywords: Local Orlicz-Morrey type spaces, Riesz potential, commutator, boundedness, Adams-type results.

References:

1. Deringoz F., Guliyev V.S., & Hasanov S.G. (2016). Characterizations for the Riesz potential and its commutators on generalized Orlicz-Morrey spaces. *Journal of Inequalities and Applications*, 248, 1-22.
2. Guliyev V.S., Deringoz F. (2014). On the Riesz Potential and Its Commutators on Generalized Orlicz-Morrey Spaces. *Journal of Function Spaces* Vol. 2014, Article ID 617414, 1- 11.
3. Guliyev V.S., Deringoz F. (2020). Riesz potential and its commutators on generalized weighted Orlicz-Morrey spaces. *Mathematische nachrichten*, 1-20.

ICOMAA 2023

On a class of nonlocal porous medium equations of Kirchhoff type

Uğur Sert

*Department of Mathematics, Hacettepe University,
usert@hacettepe.edu.tr*

Abstract

We study Dirichlet problem for a class of porous medium equations of Kirchhoff type. The equation under consideration provides a nonlocal coefficient and forcing term whose argument depends on $|u|$ with variable nonlinearity. We prove existence of generalized solutions of the considered problem under appropriate and general conditions. Also sufficient conditions for uniqueness are found and the decay rates for $\|u\|_2$ are obtained.

Keywords: Nonlocal diffusion, Kirchhoff-type problem, variable nonlinearity, porous medium equation

References:

1. U. Sert and S. Shmarev, On a degenerate nonlocal parabolic equation with variable source. *J. Math. Anal. Appl.* 2020; 484 (1): 123695.
2. K. Soltanov, On some modification Navier-Stokes equations, *Nonlinear Analysis*, 2003; 52 (3): 769-793.
3. J.L. Vazquez, *The porous medium equation: mathematical theory*, Oxford University Press, 2007.
4. M. Chipot, B. Lovat , On the asymptotic behaviour of some nonlocal problems, *Positivity* 1999; 3 (1): 65-81.



ICOMAA 2023

Construction of the asymptotics of the system of fundamental solutions for a class of unusual equations

Vagif Mastaliyev^{1,2} and Ulviya Iskandarova¹

¹Department of Mathematical Analysis, Azerbaijan State Pedagogical University,

²Department of Control for Intelligent Systems,

Academy of Public Administration under the President of the Republic of Azerbaijan

uliskenderova88@gmail.com

Abstract

In the present paper, the problem of construction of the asymptotics of the system of fundamental solutions of the equation of the mixed problem placed for a class of second order partial derivative differential equation is studied. First, after the equation is converted by means of definite substitutions into a matrix equation, the problem of construction of the asymptotics of the system of fundamental solutions of this equation is brought to the construction of the asymptotics of the solution of the corresponding integral equation. One of the main features of this equation is that the arguments of the roots of the characteristic equation corresponding to the equation in the meaning of Birkhof are not constant.

Keywords: Mixed problem, differential equation, system of fundamental solutions, matrix, integral equation, characteristic equation

References:

1. Rasulov M.L. Contour integral method, M. Nauka, 1964, 462 p.
2. Fedoryuk M.V. Asymptotic methods for linear ordinary differential equations, M., Nauka, 1984, 352 p.
3. Mamedov Yu.A. On Sturm-Liouville problem in the case of complex plane, Vestnik BSU, 1998, №1, p.133-142.
4. Mastaliyev V.Yu. Asymptotics of solutions of a Sturm-Liouville boundary value problem. Proceedings of the Scientific Conference "Problems of Applied Mathematics" . Baku 2001, p.64-69.

ICOMAA 2023

A new type algorithm for finding a common solution of Cayley inclusion and J -fixed point problem in Hilbert Spaces

Uqba Rafat¹

¹ Department of Mathematics and Statistics, Integral University Lucknow,
uqba.1709@gmail.com

Abstract

The objective of the article is to find the common solution of Cayley Inclusion problem and J -fixed point problems in the real Hilbert spaces. In addition, we support all our claims with numerical examples.

Keywords: Cayley Operator, Nonexpansive Mappings, Resolvent Operator, Inclusion Problem

References:

1. S. S. Chang, W. Takahashi, M. Toyoda, Some problems and results in the study of non-linear analysis, *Nonlinear Anal. Theory Methods Appl.* 29(7) (1997), 4197–4208
2. Q. H. Ansari, F. Babu, D. R. Sahu, Iterative Algorithms for System of Variational Inclusions in Hadamard Manifolds, *Acta Math. Sci.* 42 (2022), 1333—1356
3. M. Akram, M. Dilshad, A. K. Rajpoot, F. Babu, R. Ahmad, J. C. Yao, Modified Iterative Schemes for a Fixed Point Problem and a Split Variational Inclusion Problem, *Mathematics* 10 (2022), 1–17.
4. D. V. Hieu, P. K. Anh, L. D. Muu, J. C. Strodiot, Iterative regularization methods with new stepsize rules for solving variational inclusions, *J. Appl. Math. Comput.* 68 (2022), 571—599.

ICOMAA 2023

On the solution of the mixed problem posed for a class of equations degenerated by the type

Vagif Mastaliyev^{1, 2} and Bahruz Agarzayev³

¹Department of Mathematical Analysis, Azerbaijan State Pedagogical University,

²Department of Control for Intelligent Systems,

Academy of Public Administration under the President of the Republic of Azerbaijan,

³Azerbaijan State Maritime Academy

vagiftrk1@rambler.ru

Abstract

In the present paper by the contour integral method, we study the existence and uniqueness of the solution of a mixed problem for a class of equations with complex-valued coefficients that behave as a parabolic, despite the fact that over “time” can change from parabolic type to Schrodinger type, or even to antiparabolic type. Note that in the paper [4], [5] it was shown that mixed problems can be both ill-posed for I.G. Petrovsky well-conditioned equations and well-posed for ill-conditioned equations.

Note that one of characteristic properties of these equations is the fact that for the equations of corresponding spectral problems, the argument of roots of characteristic polynomial J.Birkhoff are not constant.

Keywords: Cauchy problem, parabolic equation, classical solution, existence, uniqueness.

References:

1. Mamedov Yu.A. On Sturm-Liouville problem in the case of complex plane, Vestnik BSU, 1998, №1, p.133-142.
2. Mamedov Yu. A., Mastaliyev V.Yu. On Growth of Green's function of Sturm-Liouville problem with complex density at the parameter. Proceedings of IMM of NASA. VOL. XVII (XXV), Baku- 2002, pp.122-127.
3. Mastaliyev V.Yu. Asymptotics of solutions of a Sturm-Liouville boundary value problem. Proceedings of the Scientific Conference “Problems of Applied Mathematics”. Baku 2001, p.64-69.
4. Mamedov Yu.A. Investigation of the well-defined of linear one-dimensional mixed problems for general systems of partial differential equations in particular derivatives with constant coefficients. - Baku, 1988. (Preprint of the Institute of Physics AN Azerbaijan. SSR. 20. -67 p.).
5. Mamedov Yu.A. On the well-defined of general mixed problems. Differential. equations. -1990. –vol. 26, no 3. p. 534-537.

ICOMAA 2023

Blow up of solutions for a logarithmic Petrovsky equation with damping term

Yavuz Dinç¹, Erhan Pişkin², Cemil Tunç³

¹Department of Electronics Technical, Mardin Artuklu University, Mardin

²Department of Mathematics, Dicle University, Diyarbakır

³Department of Mathematics, Van Yüzüncü Yıl University, Van

yavuzdinc@artuklu.edu.tr

episkin@dicle.edu.tr

cemtunc@yahoo.com

Abstract

In this presentation, we consider a logarithmic Petrovsky equation with damping term. We prove the blow up of solutions. The equation with the logarithmic term is related with many branches of sciences. Cause of this is interest in it occurs naturally in inflation cosmology, nuclear physics, supersymmetric field theories and quantum mechanics.

Keywords: Blow up, Petrovsky equation, Damping term.

References:

1. Y. Dinç, E. Pişkin, C. Tunç, Upper bounds for the blow up time for the Kirchhoff-type equation, Communications Faculty of Sciences University of Ankara Series A1: Mathematics and Statistics, (in press).
2. H. Di, Y. Shang, Z. Song, Initial boundary value problem for a class of strongly damped semilinear wave equations with logarithmic nonlinearity, Nonlinear Analysis, 51 (2020), 1-22.
3. E. Pişkin, S. Boulaaras, N. İrkıl, Qualitative analysis of solutions for the p-Laplacian hyperbolic equation with logarithmic nonlinearity, Mathematical Methods in the Applied Sciences, 44 (2021) 5654-4672.
4. E. Pişkin, Z. Çalışır, Decay and blow up at infinite time of solutions for a logarithmic Petrovsky equation, Tbilisi Mathematical Journal, 13(4) (2020) 113-127.
5. G. Zu, B. Guo, Bounds for lifespan of solutions to strongly damped semilinear wave equations with logarithmic sources and arbitrary initial energy, Evolution equations and control theory, 10 (2) (2021), 250-270.

ICOMAA 2023

Regular and Semiregular Hypermodules

Yıldız Aydın¹, Burcu Nişancı Türkmen²
Department of Management and Information Systems,
Istanbul Gelisim University
Department of Mathematics, Amasya University
yaydin@gelisim.edu.tr
burcu.turkmen@amasya.edu.tr

Abstract

Our main goal is to study regular and semiregular hypermodules which associated with small subhypermodules for a Krasner hypermodule. We present some results that give a connection between regular hyperrings and these hypermodules.

Keywords: Hyperring, Hypermodule, Regular hypermodule, Semiregular hypermodule.

References:

1. Krasner M., A class of hyperrings and hyperfields, *Int. J. Math. Math. Sci.* 6(2) (1983) 307-312.
2. Massouros C.G., Free and cyclic hypermodules, *Ann. Math. Pura. Appl.* 150(1) (1988) 153-166.
3. Hamzakolaee A. R.M., Norouzi M., & Leoreanu-Fotea V., A New Approach to Smallness in Hypermodules. *Algebraic Structures and Their Applications*, 8(1)(2021) 131-145.
4. Talae B., Small subhypermodules and their applications, *Romanian Journal of Mathematics and Computer Science*. 3(1) (2013) 5-14.
5. Nişancı Türkmen B., Bordbar H. & Cristea I., Supplements related to normal π -projective hypermodules, *Mathematics*. 10(11) (2022) 1-15.

ICOMAA 2023

Some Fixed-Point Theorems for Continuous Functions

YunusAtalan¹ and Samet Maldar²

^{1,2}*Department of Mathematics, Aksaray University,*

yunusatalan@aksaray.edu.tr

mmaldar@aksaray.edu.tr

Abstract

In this study, an iteration method that is constructed by using any continuous mapping defined on an arbitrary interval is discussed. The necessary and sufficient conditions for the convergence of this iteration method are analyzed. Finally, the comparison of the convergence rates of some iteration methods is examined through a numerical example.

Keywords: Iteration method, Continuous mapping, Convergence, Rate of convergence.

References:

1. Karakaya, V., Atalan, Y., Dogan, K., and Bouzara, N. (2017). Some Fixed-Point Results for A New Three Steps Iteration Process in Banach Spaces. *Fixed Point Theory*, 18(2), 625-640.
2. Dogan K., Gursoy F., and Karakaya V. (2020). Some Fixed-Point Results for Continuous Functions on An Arbitrary Intervals. *Sigma Journal of Engineering and Natural Sciences*, 37(3), 755-767.
3. Qing, Y., Qihou, L. (2006). The Necessary and Sufficient Condition for The Convergence of Ishikawa Iteration on An Arbitrary Interval, *Journal of Mathematical Analysis and Applications*, 323, 1383-1386.
4. Phuengrattana, W, Suantai, S. (2011) On the rate of convergence of Mann, Ishikawa, Noor and SP-iterations for continuous functions on an arbitrary interval. *Journal of Computational and Applied Mathematics*. 235, 3006-3014.

ICOMAA 2023

On Some Different Representations of q -Catalan Numbers and Their Sums

Serpil Halıcı¹, Zehra Betül Gür²

^{1,2}Department of Mathematics, Pamukkale University,

shalici@pau.edu.tr

gurbetul35@gmail.com

Abstract

In this study, we give some representations of q -Catalan numbers using classical definition of q -Catalan numbers denoted by $C_n(q)$. We examine the various sums of the sequence of $C_n(q)$ related to q -Central binomial coefficients and give some formulas for this sums. Furthermore, we give a new formula of the finite sum of $C_n(q)$.

Keywords: Catalan numbers, q -identities, binomial coefficients.

References:

1. G.E. Andrews, Catalan numbers, q -Catalan numbers and hypergeometric series, J. Combin. Theory Ser. A, 44 (2) (1987), 267-273.
2. S.B. Ekhad, M. Yang, D. Zeilberger, Automated Proofs of Many Conjectured Recurrences in the OEIS made by R.J. Mathar, arXiv:1707.04654, (2017).
3. J. Furlinger, J. Hofbauer, q -Catalan numbers, J. Combin. Theory Ser. A, 40 (2) (1985), 248-264.
4. V.J. Guo, J. Zeng, Factors of binomial sums from the Catalan triangle, Journal of Number Theory, 130(1) (2010), 172-186.
5. T. Koshy, Catalan numbers with applications, Oxford University Press, 2008.



ICOMAA 2023

A Tauberian Theorem for the Weighted Mean Summability Method of Double Sequences in Ordered Spaces

Zerrin Önder Şentürk

Department of Mathematics, Uşak University,
zerrin.onder11@gmail.com

Abstract

In this study, we are interested in the relation between the weighted mean summability method, or briefly the (\overline{N}, p, q) method, and convergence for double sequences. Accordingly, we extend a Tauberian theorem, given for the Cesàro summable single sequences by Maddox [4] and for the (\overline{N}, p) summable single sequences by Çanak [3] in ordered spaces, to the (\overline{N}, p, q) summability method.

Keywords: Double sequences, ordered linear space, the slow decrease condition, the weighted mean summability method.

References:

1. F. Móricz, Tauberian theorems for Cesàro summable double sequences, *Studia Math.* 110(1) (1994) 83–96.
2. F. Móricz and B. E. Rhoades, Necessary and sufficient Tauberian conditions for certain weighted mean methods of summability II, *Acta Math. Hungar.* 102(4) (2004) 279–285.
3. I. Çanak, A Tauberian theorem for a weighted mean method of summability in ordered spaces, *Nat. Acad. Sci. Lett.* 43(6) (2020) 553–555.
4. I. J. Maddox, A Tauberian theorem for ordered spaces, *Analysis* 9(3) (1989) 297–302.
5. R. Schmidt, Über divergente Folgen und lineare Mittelbildungen, *Math. Z.* 22(1) (1925) 89–152.

ICOMAA 2023

Fundamental Concepts in Variable Lebesgue Spaces Associated with Laplace-Bessel Differential Operator

Zeynep Tercan¹, Esra Kaya²

^{1,2}Department of Mathematics, Bilecik Şeyh Edebali University,
zeynep271100@gmail.com
esra.kaya@bilecik.edu.tr

Abstract

In this paper, we consider the concepts of convergence in $L_{p(\cdot),\gamma}(\mathbb{R}_{k,+}^n)$. In variable Lebesgue spaces, there are three types of convergence: convergence in modular, convergence in norm and convergence in measure. We investigate the relationship between these convergences. Furthermore, we prove that $L_{p(\cdot),\gamma}(\mathbb{R}_{k,+}^n)$ are Banach spaces.

Keywords: Completeness, convergence, variable Lebesgue spaces.

References:

1. Cruz-Urbe, D., Fiorenza, A., Variable Lebesgue spaces, Foundations and Harmonic Analysis, Springer Science & Business Media (2004).
2. Klyuchantsev, M. I., On singular integrals generated by the generalized shift operator I, Sibirsk. Math. Zh., 11 (1970) 810-821; translation in Siberian Math. J., 11 (1970) 612-620.
3. Levitan, B. M., Bessel function expansions in series and Fourier integrals, Uspekhi Mat. Nauk 6, 42(2) (1951) 102-143.
4. Orlicz, W., Über konjugierte Exponentenfolgen, Studia Math., 3 (1931) 200-211.

ICOMAA 2023

The Exact Solutions of Some Difference Equations via Narayana Numbers

Zeynep Akmeşe¹, Melih Göcen²

^{1,2}Department of Mathematics, Zonguldak Bülent Ecevit University,

albayrak_z@yahoo.com

gocenm@hotmail.com

Abstract

In this work, we obtain the form of the solutions of the following rational difference equations

$$x_{n+1} = \frac{1}{x_n(x_{n-1} - 1)}$$

associated with Narayana numbers. Furthermore, we deal with the asymptotic behavior of the equilibrium point of this equation. We find an interesting relation between the exact solution of the equations and the Narayana sequence.

Keywords: Narayana Numbers, difference equations, equilibrium point, asymptotic behavior, global asymptotic stability.

References:

1. Camouzis E and Ladas G (2007) Dynamics of Third-Order Rational Difference Equations with Open Problems and Conjectures. Vol. 5 Chapman & Hall/CRC, Boca Raton, Fla, USA.
2. Elaydi S (2005) An Introduction to Difference Equations. 3rd ed., Springer-Verlag, New York.
3. Kulenovi'c M R S and Nurkanovic M (2003) Global Asymptotic Behavior of Two-dimensional System of Difference Equations Modeling Cooperation. Journal of Difference Equations and Applications, 9 (1): 149-159.
4. Okumuş İ and Soykan Y (2018) Some Technique to Show the Boundedness of Rational Difference Equations. Journal of Progressive Research in Mathematics, 13(2): 2246-2258.

Statistical Convergence of Asymmetric Metric Valued Sequences with Deferred Cesàro Mean

Zeynep Hande Toyganözü

*Department of Mathematics, Suleyman Demirel University,
handetoyganozu@sdu.edu.tr*

Abstract

The aim of this study is to generalize the concept of statistical convergence of sequences with deferred Cesàro mean and deferred Cesàro summability of a sequence to asymmetric metric spaces which was defined without symmetry property. Since, lackness of symmetry causes to be given some of the classical statements differently from the metric case, it can be interesting to give some properties of these concepts and investigate the inclusion relations between them in such spaces.

Keywords: statistical convergence, deferred statistical convergence, asymmetric metric space.

References:

1. R.P. Agnew, On deferred Cesàro mean, *Ann. Math.* 33 (1932), 413-421.
2. J. Collins, J. Zimmer, An asymmetric Arzelà-Ascoli theorem, *Topology and its Applications*, 154 (2007), 2312-2322.
3. M. Et, M. Çınar, H.Şengül, Deferred Statistical Convergence in Metric Spaces, *Conference Proceedings of Science and Technology* 2(3) (2019), 189-193.
4. H. Fast, Sur la convergence statistique, *Colloquium Mathematicum* 2 (1951), 241-244.
5. M. Küçükaslan, M. Yılmaztürk, On deferred statistical convergence of sequences, *Kyungpook Math. J.* 56 (2016), 357-366.

ICOMAA 2023

Three weighted inequality for a superposition of some operators

Yerniyashova Zhadyra


L.N. Gumilyov Eurasian National University,
yerniyashova.z@gmail.com

Weighted inequalities play an important role in solving various theoretical and practical problems of function theory and functional analysis. The validity of the three-weight inequality in Lebesgue spaces for certain values of the parameters p and q and characterizing weighted functions for this inequality holds for all nonnegative measurable functions are defined in this work. The result obtained is of interest to specialists in function theory and functional analysis.

Keywords: Three weighted inequality, superposition of operators, Hardy operator, Copson operator, Tandory operator.

References:

1. Gogatishvili A., Mustafayev R.CH., Persson L.E., Some new iterated Hardy-type inequalities, *J.Funct.Spaces Appl.*30, 134194, 2012, 30c.
2. W.D. Evans, A. Gogatishvili, and B. Opic, The reverse Hardy inequality with measures V_{11} (2008), no. 1, 43-74.
3. Gogatishvili A., Pick L.: Discretization and anti-discretization of rearrangement-invariant norms. *Publ. Mat.* 47(2), 311-358 (2003).



ICOMAA 2023

On the bilinear Riesz potential and its commutators on generalized Orlicz-Morrey spaces

Zhanat Zhuldassov¹, Talgat Akhazhanov², Dauren Matin²

¹Faculty of Mathematics and Information Technology, Buketov University, Karaganda, Kazakhstan
zhanzhan85@mail.ru

²L.N. Gumilyov Eurasian National University, Astana, Kazakhstan
D.matin@mail.ru
Talgat-a2008@mail.ru

Abstract

In this paper, the authors give a characterization of the compactness for the bilinear Riesz potential commutator. More precisely, the authors prove that the commutator of the bilinear Riesz potential is a compact operator on generalized Morrey spaces.

Keywords: Riesz potential, Commutators, Compactness, VMO, Generalized Morrey space.

This work is funded by the Science Committee of the Ministry of Education and Science of the Republic of Kazakhstan (grant No. AP15473253).

References:

1. V. I. Burenkov, H. V. Guliyev, and V. S. Guliyev, "Necessary and sufficient conditions for the boundedness of fractional maximal operators in local Morrey-type spaces," *Journal of Computational and Applied Mathematics*, vol. 208, no. 1(2007) 208-301.
2. Guliyev, VS, Deringoz, F: On the Riesz potential and its commutators on generalized Orlicz-Morrey spaces. *J. Funct. Spaces* 2014, Article ID 617414 (2014).
3. E.M. Stein and G. Weiss, Fractional integrals on n-dimensional Euclidean space, *J.Math. Mech.*, (1958) 503-514.
4. Y. Chen, Y. Ding, X.Wang, Compactness for Commutators of Marcinkiewicz Integral in Morrey Spaces, *Taiwanese Jour. Math.* 15 (2011) 633-658.
5. Guliyev, V. S., Hasanov, J. J., & Zeren, Y. Necessary And Sufficient Conditions for The Boundedness of The Riesz Potential In Modified Morrey Spaces. *Journal of Mathematical Inequalities*, 5(4), (2011) 491-506.

ICOMAA 2023

Recursive Estimation of the Conditional Distribution Function on Riemannian Manifolds

BOUADJEMI Abdelkader¹

¹ *Department of Mathematics,
Amine Elokka El Hadj Moussa University of Tamanerasset
Tamanerasset,
Algeria.
bouadjemi@gmail.com*

Abstract

The main of this paper is to treat the estimation of conditional distribution function on Riemannian manifolds. We defined the recursive version of the Nadaraya–Watson estimator. Under some assumptions in Riemannian Manifolds data analysis, we study the properties of a recursive family kernels regression, the asymptotic normality of estimator is established.

Keywords: Riemannian Manifold, asymptotic normality, conditional cumulative distribution function.

References:

5. Bouadjemi, A. Asymptotic normality of the recursive kernel estimate of conditional cumulative distribution function. Journal of probability and statistical sciences, (2014). 12, Pages 117-126.
6. Salah khardani ; Wiem NefziAnne ; Françoise Yao. Nonparametric Kernel Spatial Density Estimation on Riemannian manifolds. January 2022. SSRN Electronic Journal; DOI: 10.2139/ssrn.4222942
7. Salah KhardaniAnne Françoise Yao. Nonparametric recursive regression estimation on Riemannian Manifolds. Statistics & Probability Letters October 2021,182(2):109274
8. Bruno Pelletier. (2006). Non-parametric regression estimation on closed Riemannian manifolds. Journal of Nonparametric Statistics 18(1):57-67.

ICOMAA 2023

Existence and Uniqueness of the strong Solution of the Time fractional integro-differential equation with integral boundary conditions

Ahcene Merad¹, Ouarda Benmanseur²

^{1,2}Department of Mathematics, Oum El Bouaghi University, Algeria

merad_ahcene@yahoo.fr

obenmanseur@yahoo.com

Abstract

In this work, we study the existence and uniqueness of a solution for the fractional order partial fractional differential equations with integral conditions. Using the method of energy inequalities, we find a priori estimate and the density of the range of the operator generated by given the problem.

Keywords: Nonlocal problem; Time fractional differential equation; Nonlocal integral conditions; Strong solution.

References:

1. J. Martin-Vaquero, A. Merad, Existence, uniqueness and numerical solution of a fractional PDE with integral conditions, *Nonlinear Analysis: Modelling and Control*, 24 (3), (2019)368-386.
2. T.E. Oussaeif, A. Bouziani, Existence and uniqueness of solution to parabolic fractional differential equations with integral conditions, *Electron. J. Differ. Equ.*, (2014)179.
3. A. Merad, J. Martín-Vaquero, A Galerkin method for two-dimensional hyperbolic integrodifferential equation with purely integral conditions, *Appl. Math. Comput.*, 291:386--394, 2016.
4. A. Bouziani, On a class of nonlinear reaction-diffusion systems with nonlocal boundary conditions, *Abstr. Appl. Anal.*, 9: (2004)793-813.

ICOMAA 2023

Banach Hardy classes and some Properties

Vafa Alili

*Baku Slavic University, 33 S. Rustam street, Baku, Azerbaijan,
alilivefa@mail.ru*

Abstract

In this work, rearrangement invariant space is considered and the norm generated by it of the Hardy classes of analytic functions inside and outside the unit ball, respectively. Using the shift operator, the subspace is distinguished, it is proved that infinitely differentiable and compactly supported functions are dense in this subspace. Some properties of functions from Hardy classes are studied. The classical Lebesgue spaces, the grand-Lebesgue spaces, the Orlicz spaces, and many other spaces are rearrangement invariant spaces.

Keywords: Rearrangement invariant spaces, Hardy classes, Boyd indices.

References:

1. C. Bennett, R. Sharpley, Interpolation of Operators (Academic Press, 1988.)
2. B.T. Bilalov, Z.G. Guseynov, Basicity criterion for perturbed systems of exponents in Lebesgue spaces with variable summability, Dokl. RAN, 436 (5), 586-589 (2011)
3. B.T. Bilalov, Z.G. Guseynov, Basicity of a system of exponents with a piece-wise linear phase in variable spaces, Mediterr. J. Math., 9 (3), 487-498 (2012)



ICOMAA 2023

An Inverse Coefficient Problem for a Two-Dimensional Hyperbolic Equation with Integral Overdetermination

Elvin Azizbayov¹ and Gulnara Shukurova²

¹Department of Control for Intelligent Systems,
Academy of Public Administration under the President of the Republic of Azerbaijan,
eeazizbayov@dia.edu.az

²Department of Higher Mathematics, Baku State University
shukurova-gulnara@rambler.ru

Abstract

The paper considers a classical solvability of a nonlinear inverse boundary value problem for a two-dimensional hyperbolic equation with an integral overdetermination condition. To investigate the solvability of the stated problem, we first consider an auxiliary inverse boundary value problem and prove its equivalence to the original problem. Then we prove the existence and uniqueness theorem for the auxiliary problem by the contraction mappings principle. Furthermore, based on the equivalency of these problems, the existence and uniqueness theorem for the classical solution of the original inverse problem is proved.

Note that the related inverse problems for one-dimensional hyperbolic equation was studied in the papers [1-3] and the references therein.

Keywords: Inverse problem, hyperbolic equation, classical solution, existence, uniqueness.

References:

1. Azizbayov E.I. Inverse coefficient identification problem for a hyperbolic equation with nonlocal integral condition. Turk. J. Math. Vol. 46, No.4 (2022), 1243-1255.
2. Mehraliyev Y.T., Azizbayov E.I. A time-nonlocal inverse problem for a hyperbolic equation with an integral overdetermination condition. Electron. J. Qual. Theory Differ. Equ. Vol. 2021, No.29 (2021), 1-12.
3. Mehraliyev Ya.T., Isgenderova Q.N. Inverse boundary value problem for a second-order hyperbolic equation with integral condition of the first kind. PFMT. No.1(26) (2016), 42-47.

ICOMAA 2023

An Inverse Coefficient Problem for a Two-Dimensional Hyperbolic Equation with Integral Overdetermination

Elvin Azizbayov¹ and Gulnara Shukurova²

¹Department of Control for Intelligent Systems,
Academy of Public Administration under the President of the Republic of Azerbaijan,
eeazizbayov@dia.edu.az

²Department of Higher Mathematics, Baku State University
shukurova-gulnara@rambler.ru

Abstract

The paper considers a classical solvability of a nonlinear inverse boundary value problem for a two-dimensional hyperbolic equation with an integral overdetermination condition. To investigate the solvability of the stated problem, we first consider an auxiliary inverse boundary value problem and prove its equivalence to the original problem. Then we prove the existence and uniqueness theorem for the auxiliary problem by the contraction mappings principle. Furthermore, based on the equivalency of these problems, the existence and uniqueness theorem for the classical solution of the original inverse problem is proved.

Note that the related inverse problems for one-dimensional hyperbolic equation was studied in the papers [1-3] and the references therein.

Keywords: Inverse problem, hyperbolic equation, classical solution, existence, uniqueness.

References:

1. Azizbayov E.I. Inverse coefficient identification problem for a hyperbolic equation with nonlocal integral condition. Turk. J. Math. Vol. 46, No.4 (2022), 1243-1255.
2. Mehraliyev Y.T., Azizbayov E.I. A time-nonlocal inverse problem for a hyperbolic equation with an integral overdetermination condition. Electron. J. Qual. Theory Differ. Equ. Vol. 2021, No.29 (2021), 1-12.
3. Mehraliyev Ya.T., Isgenderova Q.N. Inverse boundary value problem for a second-order hyperbolic equation with integral condition of the first kind. PFMT. No.1(26) (2016), 42-47.

ICOMAA 2023

A Compact Finite Difference Method for Fractional Equation Based on Caputo Derivative: Codes Matlab

BENDIB ISSAM¹, DALAH MOHAMED², BOUDJEDOUR ALLAOUA³

*1,2,3University Frères Mentouri Constantine 1
P.O. Box, 325 Ain El Bey Way, Constantine, Algeria, 25017.*

isambendib97@gmail.com

mdalah.ufmc@gmail.com

haboudjedour@yahoo.fr

Abstract

Abstract. In this new work, we start by some definitions and notations are prepared about caputo derivative operator and their properties. First, we give a compact finite difference scheme is described and then the stability and convergence rates are rigorously analyzed for the. Finally, we give some numerical experiments are carried out.

Keywords: A time-fractional, Caputo operator, Compact finite difference scheme, Stability, Convergence, Codes Matlab.

References:

1. M. Dehghan, M. Abbaszadeh, and W. Deng, "Fourth-order numerical method for the space-time tempered fractional diffusion-wave equation," *Applied Mathematics Letters*, vol. 73, no. 11, pp. 120–127, 2017.
2. H. Sun, Z.-Z. Sun, and G.-H. Gao, "Some high order difference schemes for the space and time fractional Bloch- Torrey equations," *Applied Mathematics and Computation*, vol. 281, pp. 356–380, 2016.
3. S. Arshad, J. Huang, A. Q. M. Khaliq, and Y. Tang, "Trapezoidal scheme for time-space fractional diffusion equation with Riesz derivative," *Journal of Computational Physics*, vol. 350, pp. 1–15, 2017.
4. H. Wang, K. Wang, and T. Sircar, "A direct $O(N \log 2N)$ finite difference method for fractional diffusion equations," *Journal of Computational Physics*, vol. 229, no. 21, pp. 8095–8104, 2010.

ICOMAA 2023

Recursive Estimation of the Conditional Distribution Function on Riemannian Manifolds

BOUADJEMI Abdelkader¹

¹ *Department of Mathematics,
Amine Elokkel El Hadj Moussa University of Tamanerasset
Tamanerasset,
Algeria.
bouadjemi@gmail.com*

Abstract

The main of this paper is to treat the estimation of conditional distribution function on Riemannian manifolds. We defined the recursive version of the Nadaraya–Watson estimator. Under some assumptions in Riemannian Manifolds data analysis, we study the properties of a recursive family kernels regression, the asymptotic normality of estimator is established.

Keywords: Riemannian Manifold, asymptotic normality, conditional cumulative distribution function.

References:

1. Bouadjemi, A. Asymptotic normality of the recursive kernel estimate of conditional cumulative distribution function. *Journal of probability and statistical sciences*, (2014). 12, Pages 117-126.
2. Salah khardani ; Wiem NefziAnne ; Françoise Yao. Nonparametric Kernel Spatial Density Estimation on Riemannian manifolds. January 2022. SSRN Electronic Journal; DOI: 10.2139/ssrn.4222942
3. Salah KhardaniAnne Françoise Yao. Nonparametric recursive regression estimation on Riemannian Manifolds. *Statistics & Probability Letters* October 2021,182(2):109274

Bruno Pelletier. (2006). Non-parametric regression estimation on closed Riemannian manifolds. *Journal of Nonparametric Statistics* 18(1):57-67.

ICOMAA 2023

Numerical Analysis Applied to Stiffened Shell Structures

Hamadi Djamal¹, Legouirah Maria² and Abdulrahman Al-Nadhari³
^{1, 2, 3} ARGHYDE Laboratory, Civil Engineering and Hydraulics Department, Biskra University,
d.hamadi@univ-biskra.dz , legouirah2018@gmail.com, a.alnadhari@univ-biskra.dz

Abstract

To avoid large deformations caused by static or dynamic external loads on shell structures in civil and mechanical engineering; longitudinal or circumferential beams as stiffeners must reinforce the shells. The finite element method is the most powerful numerical method used for the numerical analysis of structures. In the presence of stiffeners; modeling has become a complicated task to solve, because of the use of two finite elements with different behavior (shell element – beam element). Moreover, the problem of incompatibility of the degrees of freedom at the junction when using two different finite elements poses a numerical problem on the programming side. To overcome these difficulties, the analysis can be carried out by a three-dimensional finite element (solid element). In this paper, the static response of a cantilever cylindrical shell with and without stiffeners under a concentrated static load is studied using the three-dimensional solid element based on quadratic formulation “C3D8IH” of the ABAQUS software. The results obtained are compared with those obtained by the recently developed finite elements; the triangular element based on the method of discrete shear deviations (DSG3), the finite element of triangular shell (MITC3), the union of finite elements Edge-based smoothed finite element union with the triangular shell (ES-MITC3) and the SAP2000 element used as a reference solution. The numerical results obtained with different elements are discussed and commented, and the presented approach to modeling this type of structure with the three-dimensional finite element of the ABAQUS software is confirmed.

Keywords: Numerical analysis, ABAQUS software, Incompatibility degrees of freedom, Shell shell structures, Solid finite element.

References:

1. Prusty, B.G. and S.J.O.E. Satsangi, Analysis of stiffened shell for ships and ocean structures by finite element method. 2001. **28**(6): p. 621-638
2. Assan, A.J.T.-w.s., Analysis of multiple stiffened barrel shell structures by strain-based finite elements. 1999. **35**(4): p. 233-253.
3. Hamadi, D., A. Ayoub, and O.J.E.J.o.C.M. Abdelhafid, A new flat shell finite element for the linear analysis of thin shell structures. 2015. **24**(6): p. 232-255.
4. Labiodh, B., D. Hamadi, and A.J.E.S.M. Zatar, Development of a rotation free shell finite element for modeling shell structures. 2020. **8**(2): p. 151-162.

AN INTERIOR-POINT ALGORITHM FOR CQP BASED ON TRIGONOMETRIC KERNEL FUNCTION

El Amir Djeffal¹ and Randa Chalekh²

¹*Department of Mathematics, University of Batna 2, Batna Algeria,
l.djeffal@univ-batna2.dz*

²*Department of Mathematics, University of Batna 2, Batna Algeria
r.chalekh@univ-batna2.dz*

Abstract

In this paper, we present a primal-dual interior-point algorithm for convex quadratic programming problem based on a new parametric kernel function with a hiperbolic–logarithmic barrier term. Using the proposed kernel function we show some basic properties that are essential to study the complexity analysis of the correspondent algorithm which we find it coincides with the best know iteration bounds for large-update method, namely, $O(\sqrt{n} \log n \log \frac{n}{\epsilon})$ by a special choice of the parameter $\rho > 1$

Keywords:

Convex quadratic programming; kernel function; interior-point methods; iteration bound.

References:

1. N.Karmarkar, A new polynomial-time algorithm for linear programming. In Proceeding of the 16th Annual ACM. Symposium on theory of computing, pages 303-311, (1984).
2. C.Roos, T.Terlaky, J.-Ph.Vial, Theory and Algorithms for Linear Optimization. An Interior-Point Approach, John Wiley & Sons, Chichester, UK, (1997), 2nd Ed, Springer, (2006)
3. Y.Q.Bai, M.El.Ghami and C.Roos, A comparative study of kernel functions for primal-dual interior-point algorithms in linear optimization, SIAM J. Optim, 101-128 (2004)
4. M.R.Peyghami, S.F.Hafshejani, An interior point algorithm for solving convex quadratic semidefinite optimization problems using a new kernel function. Iranian J Math Sci Inform. 12 : 131–152, (2017)
5. El.A.Djeffal, M.Laouar, A primal-dual interior-point method based on a new kernel function for linear complementarity problem, Asian-European, (2018)

ICOMAA 2023

On qualitative properties of some nonuniform elliptic equations

Farman Mamedov¹

¹Institute of Mathematics and Mechanics of NAS, Azerbaijan
farman-m@mail.ru

Abstract

A talk on the qualitative properties of the weak solutions of a class of non-uniform elliptic equations

$$\operatorname{div}_x(b(x, y) \nabla_x u) + \operatorname{div}_y(\nabla_y u) = f$$

with a positive degeneration $b(x, y)$ satisfying some conditions. The proper function space, Poincare-Sobolev type inequalities, existence of the weak solution, Harnack inequality, regularity of solutions is discussed. The motivation for the proposals comes from our early studies [1--4].

Keywords: non-uniform elliptic equation, Harnack inequality, regularity of solutions, Poincare-Sobolev inequality, weak solution, Dirichlet problem.

References:

1. F.I. Mamedov and Ya. Shukurov, A Sawyer-type sufficient condition for the weighted Poincare inequality, *Positivity*, **22**, 687--699, 2018.
2. F.I. Mamedov, A Poincare's inequality with non-uniformly degenerating gradient, *Monatsh. fur Math.* **194**, 151--165, 2021.
3. F.I. Mamedov and S. Monsurro, Sobolev inequality with non-uniformly degenerating gradient, *Electr. J. Qual. Theory of Diff. Equ.* **2022** (24), 1--19, 2022.
4. F.I. Mamedov, On Harnack inequality and Holder continuity to the degenerate parabolic equations, *J. Diff. Equ.*, **340**, 521--556, 2022.

ICOMAA 2023

Chelyshkov Polynomials with Applications in Approximation Theory

Imran Talib

Department of Mathematics, Virtual University of Pakistan
imrantaalib@gmail.com, imrantalib@vu.edu.pk

Abstract

In this talk, our main motivation is to introduce a new modified and efficient numerical method to solve the fractional order derivative differential equations that are difficult to solve analytically when the source term is generalized and the boundary conditions are non-homogeneous. The framework of the proposed method is based on the newly derived fractional-order integral operational matrix of Chelyshkov polynomials which are the newest class of orthogonal polynomials. The numerical accuracy and the stability of the proposed method is demonstrated by solving various examples and comparing the results obtained otherwise in the literature.

Keywords: Orthogonal Polynomials, Square-integrable functions, Chelyshkov polynomials, Operational matrices, Fractional order differential equations, Caputo fractional order derivative.

References:

1. I. Podlubny, Fractional differential equations, volume 198 of Mathematics in Science and Engineering. Academic Press, Inc., San Diego, CA, 1999. An introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications.
2. Vladimir S Chelyshkov. Alternative orthogonal polynomials and quadratures. Electron. Trans. Numer. Anal, 25(7) (2006) 17–26.
3. A. A. Zafar, G. Kudra, and J. Awrejcewicz, An investigation of fractional Bagley–Torvik equation. Entropy, 22(1) (2019) <https://doi.org/10.3390/e22010028>.

ICOMAA 2023

On Λ -fractional variational calculus

K.A.Lazopoulos
independent researcher
Theatrou Str., Rafina, 19009 Greece
kolazop@mail.ntua.gr

Abstract: Pointing out that Λ -fractional analysis is the unique fractional calculus theory including mathematically acceptable fractional derivatives, variational calculus for Λ -fractional analysis is established. Since Λ -fractional analysis is a non-local procedure, global extremals are only accepted. That means the extremals should satisfy not only the Euler–Lagrange equation, but also the additional Weierstrass-Erdmann corner conditions. Hence non-local stability criteria are introduced. The proposed variational procedure is applied to any branch of physics, mechanics, biomechanics, etc. The present analysis is applied to Λ -fractional refraction of light and in the Λ -fractional deformation of an axially extended bar and upon the bending problem of a cantilever beam.

Keywords: Λ -fractional derivative; Λ -fractional space; initial space; fractional stress; fractional deformation; fractional strain; local stability; global stability; coexistence of phases.

References:

1. Lazopoulos, K. A. & Lazopoulos, A.K., On the Mathematical Formulation of Fractional Derivatives. *Prog. Fract. Diff. Appl.* **5(4)**, 2019, pp.261-267.
2. Lazopoulos K. A. and Lazopoulos A. On Λ -fractional variational calculus, *Annals of Mathematical Physics*, doi:<https://dx.doi.org/10.17352/amp>, 2023.
3. Lazopoulos, K.A., Lazopoulos A.K., On Λ -fractional Elastic Solid Mechanics, *Meccanica*, online, doi.org/10.1007/s11012-021-01370-y(0.12345), 2021.
4. Lazopoulos K.A., Stability criteria and Λ -fractional mechanics, *Fractals & Fractionals*. accepted, 2023.

ICOMAA 2023

On a boundary value problem for nonlinear diffusion equation with integral condition of the second kind

Matlab Salimov, Afarim Salimova
Azerbaijan State Oil and Industry University
email: m.y.salimov@gmail.com; afarims@mail.ru

Abstract

This work deals with the solvability of boundary value problem for nonlinear diffusion equation with integral condition of the second kind. Definition of classical solution for the considered boundary value problem is introduced. By the Fourier method, the problem is reduced to the system of integral equations. Using the contraction mappings method, the existence and uniqueness of the solution of the system of integral equations are proved. Finally, the existence and uniqueness of the classical solution of original problem are proved.

Keywords: boundary value problem, nonlinear diffusion equation, Fourier method, classical solution

Consider the following nonlocal boundary value problem in the rectangle $Q_T = \{(x, t) : 0 < x < 1, 0 \leq t \leq T\}$: find a function $u(x, t) \in C^{(2,1)}(\overline{Q_T})$, which satisfies in Q_T the equation [1,2]

$$a(t)u_t(x, t) = u_{xx}(x, t) + p(t)u(x, t)(1 - u(x, t)) + f(x, t), \quad (1)$$

the nonlocal initial conditions
$$u(x, 0) + \int_0^T q(t)u(x, t)dx = \varphi(x), 0 \leq x \leq 1, \quad (2)$$

and the boundary conditions $u(0, t) = 0, u_x(1, t) = 0 \quad (0 \leq t \leq T), \quad (3)$ where $\delta \geq 0$ is a given number, and $p(t), q(t), f(x, t), \varphi(x)$ are the given functions. The aim of this work is to prove the existence and uniqueness of solutions of the boundary value problem (1)-(3).

References

1. Davis, P. L. A quasilinear parabolic and a related third order problem // J. Math. Anal. and Appl. – 1972. – V. 40, №2, pp. 327-335.
2. Fisher, R.A. The Wave of Advance of Advantageous Genes / R.A.Fisher // Annals of Eugenics. -1937.-№ 7.-P.355-369.
3. Kolmogorov, A.N. Study of diffusion equation related to the increase of the amount of substance and its application to one biological problem / A.N. Kolmogorov, I.G. Petrovski, I.S. Piskunov // Bulletin MGU. Seksia A.- 1937. - V. 1, № 6, - P.1-25.

Compilation of Method for Solving Partial Differential Equations of the First-Order by the Adomian Decomposition Method

Mubark Raana¹, Yahya Q. Hasan²

¹ Department of Mathematics, Al Baydha University, Yemen,

² Department of Mathematics, Saba Region University, Yemen,

ranhbark@gmail.com

yahya217@yahoo.com

Abstract

In this paper, we present the compilation of Adomian analytical methods used in solving linear and nonlinear first-order partial differential equations (PDEs), and nonlinear single initial value questions. Moreover, some new examples have been shown.

Keywords: Adomian decomposition method, first-order PDE, Singular nonlinear first-order PDEs.

References:

1. O.G.alez-Gaxiola (2022), solution of nonlinear partial differential equations by Adomian decomposition method, *Studies in Engineering and Exact sciences*, Curitiba, v.3, n.1, p. 61-78.
2. M.A. Hassan and T.M. Elzaki (2020), Double Elzaki Transform Decomposition Method for Solving Non-Linear Partial Differential Equations, *Journal of Applied Mathematics and Physics*, 8, 1463-1471.
3. O.H.Lerma (2014), Modified Decomposition Method with New Inverse Differential Operators for Solving Singular Nonlinear IVPs in First- and Second-Order PDEs Arising in Fluid Mechanics, *International Journal of Partial Differential Equations Dep of Mech Eng*, Vol. 7.
4. A.M. Wazwaz and S. M. El-Sayed (2001), A new modification of the Adomian decomposition method for linear and nonlinear operators, *Applied Mathematics and Computation*, vol. 122, no. 3, pp. 393-405.
5. A.M. Wazwaz (2006), The modified decomposition method for analytic treatment of differential equations, *Applied Mathematics and Computation*, vol. 173, no. 1, pp. 165-176.

ICOMAA 2023

Blow up of Solutions for A Viscoelastic Wave Equations with Balakrishnan-Taylor Damping and a Delay Term

Nouri Boumaza¹, Billel Gheraibia²

¹Department of Mathematics and Computer Science, Echahid Larbi Tebessi University, Tebessa, Algeria

²Department of Mathematics and Computer Science, Larbi Ben M'Hidi University, Oum El-Bouaghi, Algeria

nouri.boumaza@univ-tebessa.dz

gheraibia.billel@univ-ueb.dz

Abstract

In this paper, we consider a viscoelastic wave equation with Balakrishnan Taylor damping and a delay term

$$u_{tt} - \left(\alpha + b \|\nabla u\|_2^2 + \alpha \int_{\Omega} \nabla u \nabla u_t dx \right) \Delta u + \sigma(t) \int_0^t g(t-s) \Delta u(s) ds + \mu_1 |u_t|^{m-2} u_t + \mu_2 |u_t(t-\tau)|^{m-2} u_t(t-\tau) = |u|^{p-2} u$$

The best of our knowledge, there is no blow-up results of solutions for the above equation. In the present paper, we study the global existence and a finite time blow-up results of solutions with negative initial energy.

Keywords: Viscoelastic wave equation, Balakrishnan-Taylor damping, delay term, Global existence, Blow up.

References:

1. B. Feng, A. Soufyane, Existence and decay rates for a coupled Balakrishnan-Taylor viscoelastic system with dynamic boundary conditions. *Math Meth Appl Sci.* 43 (6) (2020) 3375-3391.
2. B. Gheraibia, N. Boumaza, General decay result of solutions for viscoelastic wave equation with Balakrishnan-Taylor damping and a delay term. *Z. Angew. Math. Phys.* 71, 198 (2020).
3. M.J. Lee, J.Y. Park, Y.H. Kang, Asymptotic stability of a problem with Balakrishnan-Taylor damping and a time delay, *Comput. Math. Appl.* 70 (2015) 478-487.
4. S. Nicaise, C. Pignotti, Interior feedback stabilization of wave equations with time dependence delay. *Electron J Differ Equ.* 41 (2011) 1-20.
5. Z. Yang, Blow-up and Lifespan of Solutions for a Nonlinear Viscoelastic Kirchhoff Equation. *Results Math* 75, 84 (2020).

ICOMAA 2023

Dynamics of Discretized Rosenzweig–MacArthur predator–prey model with a Holling type II functional response for constant immigration

Pradeep Malik¹

¹Department of Mathematics, Faculty of Science, SGT University, Gurugram, Haryana
pradeepmaths@gmail.com

Abstract

In this paper, a discretized model for spatially interacting populations, adding constant immigration of prey to the Rosenzweig–MacArthur predator–prey model with a Holling type II functional response will be investigated. The analysis is started by determining the equilibrium points, existence, and conditions of the stability. Furthermore, it is shown that the model exhibits bifurcations of codimension 1 viz. flip bifurcation and Neimark–Sacker bifurcation on varying one parameter. Bifurcation theory and center manifold theory are used to establish the conditions for the existence of these bifurcations. An extensive numerical simulation is performed to demonstrate the analytical findings.

Keywords:

Rosenzweig–MacArthur predator–prey model, constant immigration, equilibrium points, Flip bifurcation, Neimark–Sacker bifurcation

References:

1. Sugie, J.; Saito, Y. Uniqueness of limit cycles in a Rosenzweig–MacArthur model with prey immigration. *SIAM J. Appl. Math.*, 72, 299–316, (2012).
2. Huang, J.: Bifurcations and chaos in a discrete predator–prey system with Holling type-IV functional response. *Acta Math. Appl. Sin. Engl. Ser.* 21, 157–176 (2005).
3. Singh, A., Malik, P. Bifurcations in a modified Leslie–Gower predator–prey discrete model with Michaelis–Menten prey harvesting. *J. Appl. Math. Comput.* 67, 143–174 (2021).
4. Haderler, K.P., Gerstmann, I.: The discrete Rosenzweig model. *Math. Biosci.* 98, 49–72 (1990).
5. Huang, J., Xiao, D.: Analyses of bifurcations and stability in a predator–prey system with Holling type-IV functional response. *Acta Math. Appl. Sin. Eng. Ser.* 20, 167–178 (2004).

ICOMAA 2023

Approximation on Bivariate of Durrmeyer operator based on beta function

Mohd Raiz¹, Ruchi Singh Rajawat¹, Lakshmi Narayan Mishra³, Vishnu Narayan Mishra¹

¹Department of Mathematics, Faculty of Science, Indira Gandhi National Tribal University, Lalpur Amarkantak-484887, Madhya Pradesh, India

²Department of Mathematics, School of Advanced Science, Vellore Institute of Technology, 632 014, Tamil Nadu, India

mohdraizp@gmail.com, ruchirajawat6795@gmail.com, lakshminarayan.mishra@vit.ac.in
vishunumarayanmishra@gmail.com

Abstract

In this study, we construct a new sequence of bivariate Summation-Integral type hybrid operators and their approximation behavior. Moreover, the rate of convergence of these

operators are given by using the modulus of continuity. Further, Lipschitz-maximal, Peeter's K-functional and global approximation results are investigated using weight functions.

Furthermore, approximation behavior in Boşgel functional space are studied. Lastly, the Summation-Integral type hybrid operators for the function of two variables are used to validate the numerical results and obtain the graphical illustration of the convergence behaviour of the operators univariate and bivariate case separately.

Keywords: Durrmeyer operator; Peeter's K-functional; Lipschitz-maximal; Global approximation; Boşgel space.

References:

1. S.N. Bernstein, Demonstration du théorème de Weierstrass fondée sur le calcul des probabilités, Commun. Soc. Math. Kharkov, 13(1912), 1-2.
2. J.L. Durrmeyer, Une formule d'inversion de la transformée de Laplace, Applications à la théorie des moments, Thèse de 3e cycle, Faculté des Sciences de l'Université de Paris (1967).
3. M.M. Derriennic, Sur l'approximation de fonctions intégrable sur $[0,1]$ par des polynômes de Bernstein modifiés, J. Approx. Theory, 31(1981), 32-343.
4. S.M. Mazhar and V. Totik, Approximation by modified Szász operators, Acta Sci. Math., 49 (1985), 257-269.
5. G. Prasad, P.N. Agrawal and H.S. Kasana, Approximation of functions on $[0,\infty)$ by a new sequence of modified Szász operators, Math. Forum, 6(2)(1983), 1-11.

Solution in the Small and Interior Schauder-type Estimate for the m -th Order Elliptic Operator in Morrey-Sobolev Spaces

Bilal Bilalov^{1,2}, Yusuf Zeren², Sabina Sadigova^{1,3}

¹*Institute of Mathematics and Mechanics, Baku, Azerbaijan,*

²*Department of Mathematics, Yildiz Technical University, Istanbul, Turkey,*

³*Department of Mathematics, Khazar University, Baku, Azerbaijan,*

bilal.bilalov@yildiz.edu.tr; b_bilalov@mail.ru

yzeren@yildiz.edu.tr

s_sadigova@mail.ru

Abstract

A higher order elliptic operator with non-smooth coefficients in Morrey-Sobolev spaces on a bounded domain in R^n is considered. These spaces are non separable and infinite differentiable functions are not dense in them. For this reason the classical methods for establishing interior estimates (and others including) with respect to these operators, the possibility of extending functions (with a bounded norm), determining the trace and results associated with this concept, etc. in Morrey-Sobolev spaces are not applicable, and to establish these facts a different research approach should be chosen. This work focuses on these issues. An extension theorem is proved, the trace of a function in a Morrey-Sobolev space on a $n-1$ dimensional smooth surface is defined, a theorem on the existence of a strong solution in a small is proved, an interior Schauder-type estimate in Morrey-Sobolev spaces is established. A constructive characterization of the space of traces of functions from the Morrey-Sobolev space is given, which differs from the characterization given earlier by S. Campanato. Earlier, S. Campanato [1] gave a different characterization of the space of traces based on different considerations. It should be noted that the approach proposed in this work differs from the classical approach to determining the trace.

Keywords: elliptic operator, Morrey-Sobolev spaces, solution in the small, Schauder-type estimates.

References:

1. S. Campanato, Caratterizzazione delle tracce di funzioni appartenenti ad una classe di Morrey insieme con le loro derivate prime, Annali della Scuola Normale Superiore di Pisa - Classe di Scienze, Serie 3, 15(3) (1961) 263-281.
2. B.T. Bilalov, S.R. Sadigova, On solvability in the small of higher order elliptic equations in grand-Sobolev spaces. Complex Variables and Elliptic Equations, 66:12, 2021, 2117-2130
3. B.T. Bilalov, S.R. Sadigova, Interior Schauder-type estimates for higher-order elliptic operators in grand-Sobolev spaces, Sahand Comm. in Math. Analysis, 2021, 18(2): 129-148

Properties of solutions to linear differential equations near a singular point

MAZOUZ SAID¹, SAADA HAMOUDA²

^{1,2}Departement of Mathematics Laboratory of Pure and Applied and Mathematics University of Mostaganem (UMAB) University
Mostaganem, Algeria

said.mazouz.etu@univ-mosta.dz

Abstract

In this paper, we investigate the local growth and oscillation, near the singular point $z = 0$, of solutions to the differential equation

$$f'' + \{A(z)\exp\{\frac{a}{z}\} + A_0(z)\}f' + \{B(z)\exp\{\frac{b}{z}\} + B_0(z)\}f = H(z),$$

where $A(z), A_0(z), B(z), B_0(z), H(z)$ are analytic functions in

$$D(0, R) = \{z \in \mathbb{C}: 0 < |z| < R\}$$

and a, b are non-zero complex constants.

Keywords – Linear differential equations, growth and oscillation of solutions, finite singular point, Nevanlinna theory.

References:

1. S. Cherief and S. Hamouda, Growth of solutions of a class of linear differential equations near a singular point, Kragujevac J. Math. Vol. 47(2) (2023), 187-201.
2. S. Cherief and S. Hamouda, Linear differential equations with analytic coefficients having the same order near a singular point, Bull. Iranian Math. Soc. Vol 47 (2021), 1737–1749.
3. S. Cherief and S. Hamouda, Finite and infinite order of growth of solutions to linear differential equations near a singular point, Math. Bohem., Vol. 145 (3) (2021), 1-18.
4. S. Cherief and S. Hamouda, Exponent of convergence of solutions to linear differential equations near a singular point, Graduate J. Math. (Accepted).
5. H. Fettouch and S. Hamouda; Growth of local solutions to linear differential equations around an isolated essential singularity, Electron. J. Differential Equations, Vol 2016 (2016), No. 226, 1-10.

Study of onedimensional mixed problem for one class of third order nonlinear pseudoparabolic equations

Samed Aliyev and Arzu Aliyeva

¹*Baku State University, Baku, Azerbaijan.*

samed59@bk.ru

²*Institute of Mathematics and Mechanics, National Academy of Sciences, Baku, Azerbaijan.*

arzualiyeva1966@mail.ru

Abstract

This work is dedicated to the study of existence in small of almost everywhere solution of onedimensional mixed problem for one class of third order nonlinear pseudoparabolic equations. Conception of almost everywhere solution for mixed problem under consideration is introduced. After applying Fourier method, the solution of original problem is reduced to the solution of some countable system of nonlinear integro-differential equations in unknown Fourier coefficients of the sought solution. Besides, existence theorem in small of almost everywhere solution of the mixed problem is proved by contracted mappings principle.

Keywords: nonlinear, pseudoparabolic equations, mixed problem.

References:

1. S.J.Aliyev, A.Aliyeva, G.Abdullayeva; On the existence of solution to multidimensional third order nonlinear equations, European Journal Pure and Applied Mathematics, Vol.12, 2(2019), 577-589.
2. S.J.Aliyev; On global existence of almost everywhere solution of a multidimensional mixed problem for one class third order nonlinear differential equations, Report of Baku State University, Phys. Mat. Ser. 4 (2003), 1-10 (in Russian).
3. K.I.Khudaverdiyev, Y.I.Museibli; The investigation of a onedimensional mixed problem for one class of third order nonlinear pseudoparabolic equations, Report of Baku State University, Phys.Mat. Ser.4(2002), 105-110(in Russian).
4. K.I. Khudaverdiyev; Multidimensional mixed problem for nonlinear hyperbolic equations, Az. gostekn. University publ. Baku, (2011), 611p. (in Russian).

ICOMAA 2023

Some Questions of Harmonic Analysis in Weighted Morrey Type Spaces

Samia BiBi

School of mathematical Sciences, Universiti Sains Malaysia, Malaysia.

samiaawan94@gmail.com

Abstract

In the manufacturing process of free-form complex surfaces, various problems occur which degraded the performance and can be resolved by using an optimization technique. Thousands of real-world problems can be converted into optimization techniques with distinct objective functions to acquire the optimal solution. In this paper, an assembly of GHT-Bézier developable surfaces by using a nature-inspired particle swarm optimization technique combined with optimal parameters are constructed from two GHT-Bézier boundary curves to improve the efficiency of complex engineering products. For this purpose, the control points of GHT-Bézier surface are chosen as design variables. The objective function is defined as the developability degree of the ruled surface and the shape parameters are considered to be optimization variables. So, we search for the optimum shape control parameters within the value range of shape parameters by using the PSO technique, then the developable surfaces with highly accurate developability can be visualized. The modeling examples demonstrate the effectiveness of the proposed method with the fairness of the surfaces and its approximation to the original surface.

Keywords:

GHT-Bézier developable surface, Shape control parameters, Developability degree, Particle swarm optimization.

References:

1. BiBi S, Abbas M, Misro MY, Hu G. (2019). A novel approach of hybrid trigonometric Bézier curve to the modeling of symmetric revolutionary curves and symmetric rotation surfaces. *IEEE ACCESS*. 7(1):165779–165792.
2. Aumann G. (2003). A simple algorithm for designing developable Bézier surfaces. *Comput Aided Geom Des*. 20(8):601–619.
3. Adi DIS, Shamsuddin SM, Ali A. (2009). Particle swarm optimization for NURBS curve fitting. *IEEE Xplore*. 259-253.
4. Hu G, Wu J, Li H, Hu X. (2020). Shape optimization of generalized developable H-Bézier surfaces using adaptive cuckoo search algorithm. *Adv Eng Soft*. 149:102889.

A Mathematical Model for the Breast Cancer Diagnosis and its Prognosis using Fuzzy Expert System

Sanjeev Kumar

Professor, Dept. of Mathematics, Dr. Bhimrao Ambedkar University, Agra 282002 INDIA

e-mail: sanjeevibs@yahoo.co.in

Abstract

Nowadays decision-making applications with complex issues and knowledge involving imprecision and uncertainty are very much important. Therefore, soft computing methods including Artificial Neural networks or Fuzzy Inference systems have been used broadly to model expert behaviour. Fuzzy soft computing applications are unexpectedly rising in numerous fields along with medical prognosis and diagnosis. There are several studies reported for breast cancer diagnosis, and a few types of research have also been completed on the prognosis of breast cancer. However, breast cancer prognosis suffers from vague input factors and incompleteness of information on the problem in addition to analysis. This study presents a fuzzy expert system for the diagnosis and prognosis of breast cancer, which is capable enough to record the ambiguity and imprecision regularly occurring in the interpretation of breast cancers. In this work, the Mamdani fuzzy inference model has been used as it is more practical and highly interpretable when interacting with professionals and experts during the prognosis process. This model aims to identify the risk factor for developing breast cancer in females of the age group 20-50 years. Furthermore, the fuzzy model is evaluated on a real dataset. This approach is promising for the prediction of growing the risk of breast cancer and early diagnosis of cancer, consequently, this will enhance the survival rate

Keywords: Fuzzy inference system, prognosis, diagnosis, breast cancer and Mamdani.

References:

1. Berkan, R. C., & Trubatch, S. L. (1997). Fuzzy Systems Design Principles. Building Fuzzy *if-then* rule bases. Wiley-IEEE Press, ISBN: 0780311515.
2. Zadeh, K. S. (2000). Fuzzy Health, Illness and Disease. *Journal of Medicine and Philosophy: A Forum for Bioethics and Philosophy of Medicine*, 25, 605-638.
3. Kumar, S., Hndoosh, R. W., & Sarora, M. S. (2013). A proposition for using mathematical models based on a fuzzy system with the application, *International Journal of Mathematical Sciences*, 33, 1356-1373.
4. Thani, I., & Kasbe, T. (2022). Expert system based on fuzzy rules for diagnosing breast cancer, *Health and Technology*, 12, 473-489.

Some numerical presentation for a quadratic map

Selmani Wissame¹, Djellit Ilhem²

¹*Mathematics Dynamics and modelisation laboratory, 20 Aout 1955 Skikda University, Algeria*

²*Mathematics Dynamics and modelisation laboratory, Badji Mokhtar Annaba University, Algeria*

w.selmani@univ-skikda.dz

ilhem.djelit@univ-annaba.org

Abstract

This document we base our work on polynomials of degree p and q greater than two, we generalize to move to a transformation in order to detect the attractors that exist, then we draw the bifurcation curves in different parametric planes, big bang bifurcation will appear.

Keywords: *attractor, logistic, bifurcation, fixed point, transformation*

References:

1. C.Mira, I Gumowski *Dynamique Chaotique : transformation ponctuelles Transition Ordre-désordre*, Publisher : Toulouse, 1980.
2. L. Gardini, *Homoclinic Bifurcations in n-Dimensional Endomorphisms Due to Expanding Periodic Points*, *Nonlinear Analysis TM and A*, 23 (8), (1994), 1039—1089
3. M. Henon, *A two-dimensional mapping with a strange attractor*, *Commun. Math. Phys.*, 50, (1976), 69--77.



ICOMAA 2023

Probability in Mathematical Physics: Theory and Applications

Seyyed Ali Mohammadiyeh¹ and Hamid Purhasani²

¹ *Department of Pure Mathematics, Faculty of Mathematical Sciences,
University of Kashan, Kashan, I. R. Iran*
alim@kashanu.ac.ir

² *Department of Computer Engineering, Faculty of Engineering,
University of Yazd, Yazd, I. R. Iran*
9424303@yazd.ac.ir

Abstract

Probability theory plays a crucial role in mathematical physics, providing a framework for modeling and understanding random phenomena in physical systems. This extended abstract will provide an overview of the use of probability in mathematical physics, including both theoretical foundations and practical applications. The aim is to highlight the importance of probability in this field and to demonstrate how it has been applied to solve real-world problems.

Keywords: Probability, Mathematical Physics, Quantum Mechanics, Statistical Mechanics, Stochastic Processes, Monte Carlo Methods, Data Analysis.

References:

1. F. Goodenough, "Probability and Statistics in Particle Physics," Cambridge University Press, 2005.
2. N. G. Van Kampen, "Stochastic Processes in Physics and Chemistry," North-Holland, 1992.
3. R. J. Barlow, "Statistics: An Introduction," Wiley, 1989.
4. J. W. Kleinig, "Probability and Random Processes for Engineers," Cambridge University Press, 2008.
5. S. M. Ross, "Introduction to Probability Models," Academic Press, 2010.

ICOMAA 2023

Problems of mathematical education in the curriculum in the Caucasian countries

Shahin Aghazade¹, Gunel Novruzova², Atilla Bingol³

¹*Baku State University, Baku, Azerbaijan*

²*Ganja State University, Ganja, Azerbaijan*

³*Firat University, Elazig, Turkey*

shahinaghazade@bsu.edu.az

Abstract

In this research study, teaching of mathematics in the countries where both the teaching principles of the world are advanced and backward is analyzed in a comparative manner, also problems which related to the effect of creating the necessary conditions on the development of the educational program were investigated. One of the points of focus is the comparison of the differences between countries that are best in mathematics education and developing countries.

For example, the absence of the concept of "mean(average)" in mathematics (we are not talking about arithmetic or geometric mean) should be taken as an issue that needs to be investigated.

In short, the main purpose of this research is that during the explanation of any topic in secondary education, providing completeness of topics and teaching parts that are more difficult to understand with using curriculum principles to create interest in the subject and ensure that it will be easier to pass to the next level.

We focused on principles that should be re-established in our own country - Azerbaijan, for admission to the universities of developing countries.

Keywords: principles of education, comparison of teaching process, difference in education, "mean" concept in mathematics, curriculum.

References:

1. Reynolds, D., Sammons, P., De Fraine, B., Van Damme, J., Townsend, T., Teddlie, C., et al. (2014). Educational effectiveness research (EER): a state-of-the-art review. *School Effect. School Improvement* 25, 197–230. doi: 10.1080/09243453.2014.885450
2. Mullis, I. V. S., Martin, M. O., Foy, P., and Hooper, M. (2016a). TIMSS 2015 International Results in Mathematics. Boston College, TIMSS & PIRLS International Study Center. Retrieved from: <http://timssandpirls.bc.edu/timss2015/international-results/> (accessed June 24, 2019).
3. [https://stats.libretexts.org/Bookshelves/Applied_Statistics/Introductory_Business_Statistics_\(OpenStax\)/02%3A_Descriptive_Statistics/2.05%3A_Geometric_Mean](https://stats.libretexts.org/Bookshelves/Applied_Statistics/Introductory_Business_Statistics_(OpenStax)/02%3A_Descriptive_Statistics/2.05%3A_Geometric_Mean)

On strong solvability of the Dirichlet problem for a class of semilinear elliptic equations

Sh. Yu. Salmanova
Azerbaijan State Oil and Industry University
shehla.shukurova@ mail.ru

Abstract

We study a strong solvability of the Dirichlet problem for a class of second order of the non-divergent form semilinear elliptic equations, when the leading coefficients are discontinuous and satisfy Cordes's condition, the subordinate part $g(x, u_x)$ is a Charatheodory condition function satisfying some growth condition. We get an existence result for this problem in $W_2^1(\Omega)$ Sobolev spaces, whenever the norm $\|f\|_{L_2(\Omega)}$ of the right hand side function of the equation is sufficiently small. For the proofs we have applied the Schauder's fixed point theorem on a continuous into mappings of a convex set in Banach space.

Keywords: semilinear elliptic equation, Cordes's condition, Schauder's principle.

References:

1. Alkhutov Yu. A. , Mamedov I.T. The first boundary value problem for nondivergent parabolic equations of second order with discontinuous coefficients. Matem. Sb., 131(173), (1986), no. 4 (12), 477-500 (Russian).
2. Wen G.C. Initial-mixed boundary value problems for parabolic equations of second order with measurable coefficients in a higher dimensional domain. Proceedings of the second ISAAC Congress1 (2000), 185-192.
3. Mamedov F.I., Salmanova Sh.Yu. On strong solvability of the dirichlet problem for semilinear elliptic equations with discontinuous coefficients. Trans.Natl. Acad. Sci. Azerb. Ser. Phys.- Tech. Math. Sci. **27**, no 7 (2007), 117-124.

ICOMAA 2023

Parabolic Fractional Integral Operators with Rough Kernels in Parabolic Local Generalized Morrey Spaces

Shemsiyye Muradova^{1,2}

¹Baku State University, Baku, Azerbaijan

²Ministry of Science and Education of Azerbaijan Republic,
Institute of Mathematics and Mechanics, Baku, Azerbaijan

mshams01@yahoo.com

Abstract

Let P be a real $n \times n$ matrix, whose all eigenvalues have positive real part, $A_t = t^P, t > 0, \gamma = \text{tr}P$ is the homogeneous dimension on R^n and Ω is an A_t -homogeneous of degree zero function, integrable to a power $s > 1$ on the unit sphere generated by the corresponding parabolic metric. We study the parabolic integral operator $I_{\Omega, \alpha}^p, 0 < \alpha < \gamma$ with rough kernels in the parabolic local generalized Morrey space $LM_{p, \varphi_1, P}^{\{x_0\}}(R^n)$. We find conditions on the pair (φ_1, φ_2) for the boundedness of the operator $I_{\Omega, \alpha}^p$ from the space $LM_{p, \varphi_1, P}^{\{x_0\}}(R^n)$ to another one $LM_{q, \varphi_2, P}^{\{x_0\}}(R^n), 1 < p < q < \infty, \frac{1}{p} - \frac{1}{q} = \frac{\alpha}{\gamma}$, and from the space $LM_{1, \varphi_1, P}^{\{x_0\}}(R^n)$ to the weak space $WLM_{q, \varphi_2, P}^{\{x_0\}}(R^n), 1 \leq q < \infty, \frac{1}{q} - \frac{1}{p} = \frac{\alpha}{\gamma}$.

Keywords: Parabolic fractional integral operator, parabolic local generalized Morrey space.

References:

1. Sh.A. Muradova, A.S. Balakishiyev, N.Z. Orucov. Parabolic fractional integral operators with rough kernels in parabolic local generalized Morrey spaces, Caspian Journal of Applied Mathematics, Ecology and Economics, vol.4, no.1, 2016, July, pp. 59-68.
2. V.S. Guliyev, Generalized local Morrey spaces and fractional integral operators with rough kernel, J. Math. Sci., (N. Y.) 193(2), 2013, pp. 211-227.
3. A.P. Caldereon, A. Torchinsky, Parabolic maximal function associated with a distribution, I. Advances in Math., 16, 1975, 1-64.

ICOMAA 2023

Bifurcation analysis and complex dynamics of a New Fractional System

Shiva Eshaghi

Department of Basic Science, Kermanshah University of Technology, Kermanshah, Iran

sh.eshaghi@kut.ac.ir

Abstract

In this work we analyze the dynamical behaviors of a new chaotic fractional system consisting stability, Hopf bifurcation and chaotic attractors. We obtain some conditions under which a Hopf bifurcation may occur for the system and obtain a critical value of parameter for which a Hopf bifurcation emerges. By computing different orbits of the system, we support the validity of analytical results and reveal dynamical behaviors consisting stability, limit cycles, Hopf bifurcation and chaotic attractors. We also show that the order of fractional derivative plays important role as the Hopf bifurcation parameter.

Keywords: Hopf bifurcation, Caputo fractional derivative, Chaotic dynamical system.

References:

1. Z. Jiangang, C. Yandong, D. Wenju, C. Yingxiang, and A. Xinlei, Hopf Bifurcation Analysis in a New Chaotic System with Chaos Entanglement Function, *J. Appl. Math.*, vol. 2014, Article ID 371509, 13 pages, 2014.
2. I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
3. I. Petras, *Fractional-order Nonlinear Systems: Modeling, Analysis and Simulation*, Springer, London, Beijing, 2011.
4. K. Diethelm, *The Analysis of Fractional Differential Equations*, Springer, New York, 2010.

ICOMAA 2023

Basicity of sine-cosine systems in grand Lebesgue space

Tural Hagverdi

Institute of Mathematics and Mechanics of the NAS of Azerbaijan.,
turalhagverdi@gmail.com

Abstract

Basicity of trigonometric systems $\sin(n-\alpha)t$, $n=1,2,\dots$ and $\cos(n-\alpha)t$, $n=0,1,2,\dots$ in the subspace $G^p(0,\pi)$ of grand Lebesgue space $L^p(0,\pi)$ is considered in this work, where α is a real parameter. Criteria for minimality, completeness and basicity in $G^p(0,\pi)$, $1 < p < +\infty$, with respect to the parameter α are found.

Keywords: basicity, sine-cosine system, grand Lebesgue space.

Theorem 1. Let $2\alpha + \frac{1}{p} \notin \mathbb{Z}$, $1 < p < +\infty$. Sine system $\{\sin(n-\alpha)t\}_{n \geq 1}$ forms a basis for the space $G^p(0;\pi)$ if and only if $\left[\alpha + \frac{1}{2p} \right] = 0$. If $\left[\alpha + \frac{1}{2p} \right] < 0$, then the system is minimal but not complete. If $\left[\alpha + \frac{1}{2p} \right] > 0$, then the system is complete but not minimal.

References:

1. Bilalov B.T. Basis Properties of Some Systems of Exponents, Cosines and Sines, Sibirskiy matem. Jurnal, 2004, v.45, № 2, pp.264-273
2. Bilalov B.T. On basicity of perturbed system of exponents in Morrey-type spaces. Sib. mat. журн., v.45, №2, (2019), 264-273
3. M.Ismailov On the Solvability of Riemann Problems in Grand Hardy Classes. Mathematical Notes, vol. 108, p. 523–537 (2020)

Different Versions of Dirichlet Problems for Laplace equation in Non-standard Function Spaces

Bilal Bilalov^{1,2}, Nigar Ahmadzadeh¹, Tarlan Garayev^{1,3}

¹Institute of Mathematics and Mechanics, Baku, Azerbaijan,

²Department of Mathematics, Yildiz Technical University, Istanbul, Turkey,

³Department of Mathematics, Khazar University, Baku, Azerbaijan,

bilal.bilalov@yildiz.edu.tr; b_bilalov@mail.ru

nigar_sadigova11@mail.ru

qarayevtarlan@gmail.com

Abstract

Non-standard grand-Lebesgue and Morrey spaces are considered in this work, together with grand-Sobolev and Morrey-Sobolev spaces generated by them. Dirichlet problems for Laplace equation in different versions are considered in these spaces in a bounded domain of n -dimensional space with sufficiently smooth boundary. These spaces are non-separable, so there arise some questions and differences in problem statements. We conduct a corresponding research and we illustrate the differences concerning the solvability in the classical Sobolev and Hardy spaces in two-dimensional case on the example of circle.

Keywords: non-standard Lebesgue, Morrey, grand Sobolev, Morrey-Sobolev spaces, Dirichlet problem, Laplace operator, Hardy space, solvability.

References:

1. B.T. Bilalov, [N.R. Ahmedzadeh](#), [T.Z. Garayev](#), Some Remarks on Solvability of Dirichlet Problem for Laplace Equation in Non-standard Function Spaces. [Mediterranean Journal of Mathematics](#), 19(133) (2022)
2. B.T. Bilalov, S.R. Sadigova, On solvability in the small of higher order elliptic equations in grand-Sobolev spaces, *Complex Variables and Elliptic Equations*. 66(12) (2021), 2117-2130

ICOMAA 2023

INDEX

A

Amiran Gogatishvili – 22
Abdelmoumen TIAIBA – 31
Amin Saif – 33
A.Mahdi – 33
Abdon E. Choque-Rivero – 34
Abdulhamit Kucukaslan – 35
Adem Kızıltepe – 36
Agil Khanmamedov – 37
Alaa Serhan – 38
Alexander Kushpel – 39
Ali Değirmenci – 40
Ali A. Shukur – 42
Alirza Akhmedov – 43
Amina Boucenna – 44
Amira Rouaghi – 44, 45
Amina Boucenna – 45
Ani Ozbetelashvili – 46
Artion Kashuri – 47
A.Sh.Shukurov – 49
Aynur Çöl – 50, 51
Aysel Guliyeva – 52
Ayşe Yavuz Taşcı – 53
Ayşe Fidan – 54
Ayşegül Kılıç – 55
Aysenur Busra Cakay – 56
Azhar Abek – 57
Aidos Adilkhanov – 69
Ali Demirözlü – 73
Agon Mahmuti – 78
Asif Khan – 138, 147
AFTAB ALAM – 139
Abdu Alameri – 144
Ahmet ŞAHİNER – 148
Aicha Bareche – 164
Ayub Khan – 172
Arshad Khan – 172
Ahmet Eren KILIÇ – 198
Aidos Adilkhanov – 199
Ahcene Merad – 215
Abdulrahman Al-Nadhari – 221
Afarim Salimova – 226
Arzu Aliyeva – 233

Atilla BİNGÖL – 238

B

Beherchin Akhmedli – 58,
BOUADJEMI Abdelkader – 60, 214, 220, 243
Burak Oğul – 61,
Bünyamin Kurtkaya – 62,
Bendib El Ouahma – 79,
Bülent Karakaş – 96,
B. Bilalov – 165, 183, 231
Bahruz Agarzayev – 203
Burcu Nişancı Türkmen – 205
BENDIB ISSAM – 219
BOUDJEDOUR ALLAOUA – 219

C

Claudia Capone – 23
Cemil Karacam – 63, 121
CHAHRAZED HARRAT – 64
Cheima Khennaoui – 65
Canay Aykol – 90
Cesim Temel – 151
Chaman Kumar – 168
Christoph Reisinger – 168
Cemil Tunç – 204

Ç

Çisem Güneş Aktaş – 66

D

David Cruz-Urbe – 24
Dumitru Baleanu – 25
Davud Orucov – 37
Daniyal M. Israfilov – 67
Dariusz Wardowski – 68
Dauren Matin – 69, 213
Diana Audi – 70
Diğdem ŞEN – 71
Dilara Erdemir – 72
Dilek Kazıcı – 73

Dilek SÖYLEMEZ – 86
Dr Tanisha Baghwan Dale – 125
DALAH Mohamed – 137, 219
Djellit Ilhem - 236
DERBAZI Ammar – 137

E

Erhan Pişkin – 54, 105, 106, 204
Emre Kolotoğlu – 72
Ece Gulsah Colak – 74
Edanur Ergül – 75
Ekin Uğurlu – 76, 77
Ekrem Aljimi – 78
Eldar M. Mustafayev – 80
Elif Ebren Kaya – 81
Elvin Azizbayov – 83, 217, 218
Emel Kurul – 84
Emine Önal Kır – 85
Emre GÜVEN – 86
Emre Kırılı – 87
Esmehan Uçar – 88
Esra Soslu – 89
Esra Kaya – 90, 209
Esra Celik – 91
Esra Ozge Asan – 92
Enes Seyfullah Kotil – 92
Erdal Gül – 166
Elanur ÖZCAN – 197
El Amir Djeflal – 22

F

Ferihe ATALAN - 32
Feray Hacivelioglu – 38
Fusun Özen Zengin – 53
Fatma TOKMAK FEN - 71
Fares Yazid – 93
Fatima Siham Djaradi – 93
Faruk Özger – 94, 95, 128
Fatih Tuğrul – 96
Fatih YILMAZ - 97
Fatih Sirin – 63, 98
Fatiha Najm – 99
Fatma Saydam Akman -100
Fatma Nur Budak – 101
Ferit Yalaz – 102

Feyza Elif Dal - 103
Fidan Alizadeh – 103
Faouzi Haddouchi – 158
Francisco Martínez González – 188
Farman Mamedov – 223

G

Gökçe Çaylak Kayaturan – 104
Gülistan Butakın – 105, 106
Gafurjan Ibragimov – 159
Gül Karadeniz Gözeri -170
Gulnara Shukurova – 217, 218
Gunel NOVRUZOVA – 238

H

Huseyin Tunc – 84, 92, 101
Habib Djourdem – 107
Hacer Şengül Kandemir – 108, 153
Haifa Ahmed – 109
Hajar Movsumova – 110
Hamdullah Başaran – 111
Hamide Feyza Aykut – 112
Hasan Arslan – 113
Hamadı Djamal – 221

I

Imran Talib – 224

İ

Ibrahim Senturk – 114
İlknur Atasever Güvenç – 115
Ismail TEMBELO – 116
İbrahim Demir – 177

J

Jalolxon Nuritdinov – 117

K

KARAÇAM CEMİL – 48
KARTAL ARDA EREN – 48
Kader ŞİMŞİR ACAR – 118

KARACAN EFE ATAKAN - 48
Kadriye Şimşek Alan – 119, 120
Koray İbrahim Atabey – 122
Kadriye Aydemir – 136
K.A.Lazopoulos – 225
Khanlar R. Mamedov – 51

L

Lubomira Softova – 163, 165
Legouirah Maria – 221
Lakshmi Narayan Mishra – 230

M

M. Mursaleen – 28
Mustafa Taylan Şengül – 62
Migdad Ismailov – 63, 118
Makhlouf Amar – 79
Münevver Tezer-Sezgin – 81
Murat Sari – 91, 92, 101, 180
M. A. Aziz Alaoui – 99
Murat Karakaş – 108
Mahioub Shubatah – 109
Mohammed Alsharafi – 109, 144
Mehmet Gürdal – 111
Muhammed Çınar – 122
Mikail Et – 122, 153
Mahvish Ali – 124
Manal A. Alqahtany – 125
Marek T. Malinowski – 126
Md Hasanuzzaman – 127
Mohammad Imdad – 127, 139
Md. Asaduzzaman – 128
Md. Nasiruzzaman – 129
Mehmet Unlu – 130
Mehmet Öner Şakar – 131
Melek Eriş Büyükkaya – 132
Melih Cinar – 133
Merve Ak – 134
Mine Akbaş – 134
Merve Görgülü – 135
Merve Yücel – 136
Mohammad Iliyas - 138
Mohammad Mursaleen - 138
MOHAMMAD ARIF – 139
Mohammad Dilshad – 140

Mohammad Esmael Samei – 141
Mohammed Mohswn – 142, 143
Mutu Akar – 144
Mohd Raiz – 145
Mohd Kashif – 146
Mohd Shanawaz Mansoori – 147
Muhammad Wasim Awan – 148
Murat Kirişci – 149, 150
Müberra Selah – 151
Mehdi Benabdallah – 155, 156
Münevver Tezer-Sezgin – 176
Muharrem Bayrak – 181
Muhammad Zafar Iqbal – 188
Mohammed Esmael Samei – 191, 192, 193
Mendybay Turgumbayev – 199
Matlab Salimov – 226
Mubark Raana – 227
Mohd Raiz – 230
MAZOUZ SAID – 232

N

Nurzhan Bokayev – 29
Nouredine Bouteraa – 59
Nazmiye Gonul Bilgin – 74
Nor-El-Houda Beghersa – 93, 155, 156
Nazlım Deniz Aral – 108, 153
Nadeem Rao – 145
Natavan Nasibova – 152
Nazir Ismailov – 152
Nebi Yılmaz – 154
Noura Djedaidi – 157
Nourredine Houari – 158
N.Nasibova – 165
Neelima – 168
Nurcan Argac – 184

O

Onur Çınar – 100
Oktay Sh. Mukhtarov – 136
Omongul Egamberganova – 159
Ouarda Benmansour – 215

Ö

Öykü YURTTAŞ – 32
Özlem Vardar – 100
Öztürk Savaş – 160
Ömer Ünsal – 178

P

Peter Hästö - 26
Penpark Sirimark – 78
Pradeep Malik – 229

R

Rozana Liko – 47
Radouane Yafia – 99
Ramazan Ercan – 130
Risman Mat Hasim – 159
Rena Sadikhzada – 161
Rokiya Sahali – 162
Rosamaria Rescigno – 163
Refet Polat – 187
Reyhan Taghiyeva – 195
Randa Chalekh – 222
Ruchi Singh Rajawat – 230

S

S. Sadigova – 27
Sercan Kazımoğlu – 36
Seda Kızılbudak Çalışkan – 55
Selmahan Selim – 56
Salim Yüce – 75, 171
Serpil Halıcı – 82
Salih AYTAR – 116
Souad Gasmi – 157
Salih Çelik – 160
Safia Hocine – 164
S. Tramontano – 165
Samet Karasu – 166
Samet Maldar – 167
Sani Biswas – 168
Sarih Sari – 169
Seyfettin Bayraktar – 169
Selin Sarı – 170
Sena Nur Aktaş – 171

Shadab Ali – 172
Shagun Sharma – 173
Sumit Chandok – 173
Shiv Kant Tiwari – 174
Shiva Eshaghi – 175, 241
Sinem Arslan Ölçer – 176
Sinem Kübra Kösen – 177
Sukri Khareng – 178
Sunil K. Sharma – 179
Sumeyye Yilmaz – 180
Sabina Sadigova – 183, 231
Serpil Halıcı – 186, 207
Sule Curuk – 186
Saadet Eskiizmirli – 187
Seyyed Ali Mohammadiyah – 189, 237
Samet Maldar – 206
SAADA HAMOUDA – 232
Samed Aliyev – 233
Samia BiBi – 234
Sanjeev Kumar – 235
Selmani Wissame – 236
Shahin AGHAZADE – 238
Sh. Yu. Salmanova – 239
Shemsiyye Muradova – 240

Ş

Şenay Baydaş – 96
Şerif Efe Dartar – 121, 181
Şerife Nur Bozdağ – 182
Şevval Kublay – 182
Seyma Cetin – 183
Seymanur Aydin – 184
Şuayip Toprakseven – 185

T

Telman Gasyimov – 58, 194, 195
Tanisha Baghwan Dale – 125
Temel ERMIŞ – 148
Talib Hussain – 188
Tasaduk Rashid Mir – 190
Taylan Demir – 191, 192, 193
Tuğçe Ünver Yıldız – 196
Tunçar ŞAHAN – 197, 198

Talgat Akhazhanov – 213
Tural Hagverdi – 242

U

Ulker Hashimova - 194
Uğur Can Yıldırım – 196
Uğur Sert – 200
Ulviya Iskandarova – 201
Uqba Rafat – 202

V


Vishnu Narayan Mishra – 145
Vagif Mastaliyev – 201, 203
Vafa Alili – 216
Vishnu Narayan Mishra – 230

Y

Yusuf Zeren – 27, 63, 103, 118, 123, 144, 183,
231
Yoshihiro Sawano – 30
Yuksel Soykan – 74
Yousaf Anwar – 100, 142
Yahya Qaid – 143, 227
Yashar Mehraliyev – 161
Yunus Atalan – 167, 206
Yavuz Dinç - 204
Yıldız Aydın – 205
Yerniyashova Zhadyra – 212

Z

Zafer Kütüğ – 100
Zehra Betül Gür – 207
Zerrin Önder Şentürk – 208
Zeynep Tercan – 209
Zeynep Akmeşe – 210
Zeynep Hande Toyganözü – 211
Zhanat Zhuldassov - 213



ICOMAA 2023